

On Functions of Integrable Mean Oscillation

Oscar BLASCO* and M. Amparo PÉREZ

Departamento de Análisis Matemático
Universidad de Valencia
46100 Burjassot
Valencia — Spain
oblasco@uv.es

Recibido: 1 de diciembre de 2004

Aceptado: 21 de abril de 2005

ABSTRACT

Given $f \in L^1(\mathbb{T})$ we denote by $w_{\text{mo}}(f)$ the modulus of mean oscillation given by

$$w_{\text{mo}}(f)(t) = \sup_{0 < |I| \leq t} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi}$$

where I is an arc of \mathbb{T} , $|I|$ stands for the normalized length of I , and $m_I(f) = \frac{1}{|I|} \int_I f(e^{i\theta}) \frac{d\theta}{2\pi}$. Similarly we denote by $w_{\text{ho}}(f)$ the modulus of harmonic oscillation given by

$$w_{\text{ho}}(f)(t) = \sup_{1-t \leq |z| < 1} \int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)| P_z(e^{i\theta}) \frac{d\theta}{2\pi}$$

where $P_z(e^{i\theta})$ and $P(f)$ stand for the Poisson kernel and the Poisson integral of f respectively.

It is shown that, for each $0 < p < \infty$, there exists $C_p > 0$ such that

$$\int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t} \leq \int_0^1 [w_{\text{ho}}(f)(t)]^p \frac{dt}{t} \leq C_p \int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t}.$$

Key words: mean oscillation, BMO, modulus of continuity.

2000 Mathematics Subject Classification: 46B25.

*Partially supported by Proyecto BMF2002-0401 and MTN2004-21420-E

1. Introduction.

As usual we denote by BMO the space of functions $f \in L^1(\mathbb{T})$ such that

$$\|f\|_* = \sup_{I \subseteq \mathbb{T}} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} < \infty,$$

where I is an arc of the circle \mathbb{T} , $|I|$ stands for the normalized length of I and $m_I(f) = \frac{1}{|I|} \int_I f(e^{i\theta}) \frac{d\theta}{2\pi}$. We write $\|f\|_{\text{BMO}} = |\hat{f}(0)| + \|f\|_*$.

If $f \in L^1(\mathbb{T})$ and $0 < t \leq 1$, we define the modulus of mean oscillation of f at the point t as

$$w_{\text{mo}}(f)(t) = \sup_{0 < |I| \leq t} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi}.$$

Clearly, for $0 < t \leq s < 1$, one has

$$\sup_{t < |I| \leq s} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \leq \frac{2}{t} \|f\|_1.$$

Hence, for $0 < t \leq s < 1$, one has

$$w_{\text{mo}}(f)(t) \leq w_{\text{mo}}(f)(s) \leq \max \left\{ w_{\text{mo}}(f)(t), \frac{2\|f\|_1}{t} \right\}. \tag{1}$$

In particular, $f \in \text{BMO}$ if and only if $w_{\text{mo}}(f)(t) < \infty$ for some (or for all) $0 < t \leq 1$.

It is known that one can consider other equivalent moduli to define BMO. For instance, for $0 < q < \infty$,

$$w_{\text{mo},q}(f)(t) = \sup_{0 < |I| \leq t} \left(\frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)|^q \frac{d\theta}{2\pi} \right)^{1/q}.$$

It is also well known, by the John-Nirenberg lemma (see [7, 8]), that there exist $C_1, C_2 > 0$ such that for all $\lambda > 0$ and any arc I with $|I| \leq t$,

$$\frac{|\{\theta \in \mathbb{T} : |f(e^{i\theta}) - m_I| > \lambda\}|}{|I|} \leq C_1 e^{-\frac{C_2 \lambda}{w_{\text{mo}}(f)(t)}}.$$

From here one gets that, for all $t > 0$,

$$w_{\text{mo}}(f)(t) \approx w_{\text{mo},q}(f)(t). \tag{2}$$

One can also consider

$$w'_{\text{mo}}(f)(t) = \sup_{|I| \leq t} \left(\frac{1}{|I|^2} \int_I \int_I |f(e^{i\theta}) - f(e^{i\varphi})| \frac{d\theta}{2\pi} \frac{d\varphi}{2\pi} \right)$$

or

$$\tilde{w}_{\text{mo}}(f)(t) = \sup_{|I| \leq t} \left(\inf_c \left(\frac{1}{|I|} \int_I |f(e^{i\theta}) - c| \frac{d\theta}{2\pi} \right) \right)$$

Clearly one gets

$$w_{\text{mo}}(f)(t) \leq w'_{\text{mo}}(f)(t) \leq 2w_{\text{mo}}(f)(t) \tag{3}$$

and

$$\tilde{w}_{\text{mo}}(f)(t) \leq w_{\text{mo}}(f)(t) \leq 2\tilde{w}_{\text{mo}}(f)(t).$$

A function f is said to have vanishing mean oscillation, in short $f \in \text{VMO}$, if

$$\lim_{|I| \rightarrow 0} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} = 0.$$

This is a closed subspace of BMO, which can be characterized in many ways (see [7, 8, 15]).

Theorem 1.1. *Let $f \in \text{BMO}$. The following statements are equivalent:*

- (i) $f \in \text{VMO}$.
- (ii) $\lim_{t \rightarrow 0^+} \|T_t f - f\|_{\text{BMO}} = 0$, where $T_t f(e^{i\theta}) = f(e^{i(\theta-t)})$.
- (iii) $\lim_{r \rightarrow 1} \|P_r * f - f\|_{\text{BMO}} = 0$, where $P_r(e^{i\theta}) = \Re\left(\frac{1+re^{-i\theta}}{1-re^{-i\theta}}\right)$.
- (iv) f belongs to the closure of $C(\mathbb{T})$ in BMO.
- (v) $\lim_{t \rightarrow 0^+} w_{\text{mo}}(f)(t) = 0$.

A generalization of BMO is the space $\text{BMO}(\rho)$, consisting of functions $f \in L^1(\mathbb{T})$ such that $w_{\text{mo}}(f)(t) = O(\rho(t))$ for a fixed function ρ with certain properties. The space $\text{BMO}(\rho)$ has been considered by various authors (see [10, 15, 17]).

Our aim will be to analyze spaces where the function ρ is not explicitly given, but we do know its behavior at the origin in terms of certain integrability conditions.

Given $0 < p < \infty$, we shall denote by $\text{MO}^p(\mathbb{T})$ the space of integrable functions such that $\int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t} < \infty$.

Due to (2), the spaces MO_q^p of functions such that $\int_0^1 [w_{\text{mo},q}(f)(t)]^p \frac{dt}{t} < \infty$ are all the same for $0 < q < \infty$.

These spaces were considered in [12] (see page 74), under a different notation. Also some spaces $\text{MO}_{s,r}^\alpha$, which are closely related to the ones considered in this paper, were introduced in [13].

We use the notations

$$\omega_\infty(f)(t) = \sup_{|\theta-\varphi| \leq t} |f(e^{i\theta}) - f(e^{i\varphi})|$$

and

$$\omega_q(f)(t) = \sup_{|u| \leq t} \left(\int_{\mathbb{T}} |f(e^{i(\theta+u)}) - f(e^{i\theta})|^q \frac{d\theta}{2\pi} \right)^{1/q}$$

for $0 < q < \infty$.

Now, for $0 < s < 1$ and $0 < p, q \leq \infty$, the Besov space $B_{q,p}^s(\mathbb{T})$ consists of functions in $L^q(\mathbb{T})$ such that $t^{-s}\omega_q(f) \in L^p((0, 2\pi), \frac{dt}{t})$. Of course the cases $B_{q,\infty}^s$ where $0 < q \leq \infty$ correspond to Lipschitz or Hölder classes, to be denoted $\text{Lip}_s(\mathbb{T})$ instead of $B_{\infty,\infty}^s(\mathbb{T})$.

We denote by X_p the space consisting of functions $L^\infty(\mathbb{T})$ such that $\omega_\infty(f) \in L^p((0, 2\pi), \frac{dt}{t})$.

From (3) we easily obtain, for any $t > 0$,

$$w_{\text{mo}}(f)(t) \leq C\omega_\infty(f)(t).$$

Hence $X_p \subset \text{MO}^p(\mathbb{T})$ for any $0 < p < \infty$.

On the other hand, if I, J are arcs on \mathbb{T} such that $I \subset J$ then

$$|m_J(f) - m_I(f)| \leq \frac{|J|}{|I|} w_{\text{mo}}(f)(|J|). \tag{4}$$

Now, given I with $|I| \leq t$, using the Lebesgue differentiation theorem, one gets

$$f(e^{i\theta}) = \lim_n m_{I_n} f, \quad \text{for a.a. } \theta \in I,$$

where I_n is a decreasing sequence of arcs containing θ such that $|I_n| = 2|I_{n+1}|$.

Hence using (4), we have that for any $f \in \text{MO}^1(\mathbb{T})$

$$\begin{aligned} |f(e^{i\theta}) - m_I| &\leq \lim_n |m_{I_n} f - m_I f| \\ &\leq \sum_{k=1}^\infty |m_{I_k} f - m_{I_{k-1}} f| \\ &\leq C \sum_{k=1}^\infty w_{\text{mo}}(f)(2^{-k}t) \\ &\leq C \int_0^t w_{\text{mo}}(f)(s) \frac{ds}{s}. \end{aligned}$$

Therefore we obtain, for any $t > 0$,

$$\omega_\infty(f)(t) \leq 2C \int_0^t w_{\text{mo}}(f)(s) \frac{ds}{s}. \tag{5}$$

This implies that $\text{MO}^1(\mathbb{T}) \subset \text{Lip}_\phi$, where Lip_ϕ stands for the space of continuous functions such that

$$|f(e^{i\theta}) - f(e^{i\varphi})| \leq C\phi(|\theta - \varphi|),$$

for $\phi(t) = \sup\{\int_0^t w_{\text{mo}}(f)(s) \frac{ds}{s} : \int_0^1 w_{\text{mo}}(f)(t) \frac{dt}{t} \leq 1\}$.

BMO-type characterizations of these spaces have been extensively considered in the literature. The reader is referred to [2, 3, 9] for the case $p = \infty$ and to [4, 5] for the cases $0 < s$ and $1 < p < \infty$.

We shall consider a description of $\text{MO}^p(\mathbb{T})$ where the averages over arcs are replaced by averages with respect the Poisson kernel.

We denote by BMOH the space of functions $f \in L^1(\mathbb{T})$ such that

$$\|f\|_{**} = \sup_{z \in \Delta} \int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} < \infty,$$

where Δ denotes the open unit disc, $P(f)(z) = \int_{\mathbb{T}} f(e^{i\theta}) P_z(e^{i\theta}) \frac{d\theta}{2\pi}$ and $P_z(e^{i\theta}) = \Re\left(\frac{1+z e^{-i\theta}}{1-z e^{-i\theta}}\right)$. We write $\|f\|_{\text{BMOH}} = |P(f)(0)| + \|f\|_{**}$.

It is not difficult to prove (see [7, 8]) that $f \in \text{BMO}$ if and only if $f \in \text{BMOH}$ with equivalent norms.

In this situation we define the modulus of harmonic oscillation of f at the point t as

$$w_{\text{ho}}(f)(t) = \sup_{1-t \leq |z| < 1} \int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)| P_z(e^{i\theta}) \frac{d\theta}{2\pi}.$$

Hence, $f \in \text{BMO}$ (respect. BMOH) if and only if $w_{\text{mo}}(f)(1) < \infty$ (respect. $w_{\text{ho}}(f)(1) < \infty$).

For $0 < p < \infty$, we denote by $\text{HO}^p(\mathbb{T})$ the space of $f \in L^1(\mathbb{T})$ such that $\int_0^1 [w_{\text{ho}}(f)(t)]^p \frac{dt}{t} < \infty$.

Of course one can also use other moduli to define this space. For instance, for $0 < q < \infty$,

$$w_{\text{ho},q}(f)(t) = \sup_{1-t \leq |z| < 1} \left(\int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)|^q P_z(e^{i\theta}) \frac{d\theta}{2\pi} \right)^{1/q},$$

or

$$\tilde{w}_{\text{ho}}(f)(t) = \sup_{1-t \leq |z| < 1} \inf_c \int_{\mathbb{T}} |f(e^{i\theta}) - c| P_z(e^{i\theta}) \frac{d\theta}{2\pi}.$$

The main objective of the paper is to show that, for $0 < p < \infty$, we have $\text{MO}^p(\mathbb{T}) = \text{HO}^p(\mathbb{T})$ and with equivalent “norms”.

The paper is divided into two sections. The first one is devoted to introducing $\text{MO}^p(\mathbb{T})$ and proving some of its properties and the second one to introducing $\text{HO}^p(\mathbb{T})$ and to showing that $\text{HO}^p(\mathbb{T})$ coincides with $\text{MO}^p(\mathbb{T})$.

2. Integrable mean oscillation.

We will see first that the modulus of mean oscillation is continuous. We shall use the following lemma.

Lemma 2.1. *Let $f \in L^1(\mathbb{T})$. If $\{I_n\}$ is a sequence of arcs such that $\lim_{n \rightarrow \infty} I_n = I$ for some arc I with $|I| > 0$ then*

$$\lim_{n \rightarrow \infty} \frac{1}{|I_n|} \int_{I_n} |f(e^{i\theta}) - m_{I_n}(f)| \frac{d\theta}{2\pi} = \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi}.$$

Proof. Let us first estimate

$$\begin{aligned} & \frac{1}{|I_n|} \int_{I_n} |f(e^{i\theta}) - m_{I_n}(f)| \frac{d\theta}{2\pi} - \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \\ & \leq \frac{1}{|I_n|} \int_{I_n} |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} + |m_{I_n}(f) - m_I(f)| \\ & \quad - \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \\ & \leq \frac{1}{|I|} \left(\int_{I_n} |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} - \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \right) \\ & \quad + |m_{I_n}(f) - m_I(f)| + 2\|f\|_1 \left(\frac{1}{|I_n|} - \frac{1}{|I|} \right). \end{aligned}$$

Notice that $\nu(A) = \int_A f(e^{i\theta}) \frac{d\theta}{2\pi}$ and $\mu^I(A) = \int_A |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi}$ are a complex and a positive measure respectively with integrable densities. Therefore the result follows passing to the limit as n goes to ∞ . \square

Proposition 2.2. *Let $f \in \text{BMO}$. Then $w_{\text{mo}}(f)$ is increasing and continuous in $(0, 1]$.*

Proof. Monotonicity has already been proved in our formula (1).

Let $0 < t_0 \leq 1$ and let us prove that it is left continuous at t_0 . Given $\varepsilon > 0$ we find $I_{t_0} \subset \mathbb{T}$ such that $0 < |I_{t_0}| \leq t_0$ and

$$w_{\text{mo}}(f)(t_0) \leq \frac{1}{|I_{t_0}|} \int_{I_{t_0}} |f(e^{i\theta}) - m_{I_{t_0}}(f)| \frac{d\theta}{2\pi} + \frac{\varepsilon}{2}.$$

Let (t_n) be a sequence such that $t_n \leq t_0$ for all $n \in \mathbb{N}$ and converges to t_0 .

If $|I_{t_0}| = t_0$, we can find $I_n \subset I_{t_0}$ such that $\lim_{n \rightarrow \infty} I_n = I_{t_0}$. Hence

$$\begin{aligned} & w_{\text{mo}}(f)(t_0) - w_{\text{mo}}(f)(t_n) \leq \\ & \leq \frac{1}{|I_{t_0}|} \int_{I_{t_0}} |f(e^{i\theta}) - m_{I_{t_0}}(f)| \frac{d\theta}{2\pi} - \frac{1}{|I_n|} \int_{I_n} |f(e^{i\theta}) - m_{I_n}(f)| \frac{d\theta}{2\pi} + \frac{\varepsilon}{2}. \end{aligned}$$

Now use Lemma 2.1 to get $\lim_{n \rightarrow \infty} w_{\text{mo}}(f)(t_0) - w_{\text{mo}}(f)(t_n) = 0$.

If $|I_{t_0}| < t_0$ there exists n_0 such that $|I_{t_0}| \leq t_n$ for $n \geq n_0$. Hence $w_{\text{mo}}(f)(t_0) - w_{\text{mo}}(f)(t_n) < \frac{\varepsilon}{2}$ for $n \geq n_0$.

To see that it is right continuous at t_0 , we shall argue as follows: Let (t_n) be a sequence such that $t_n \geq t_0$ for all $n \in \mathbb{N}$ and converges to t_0 . We shall find a subsequence (t_{n_k}) such that $\lim_{k \rightarrow \infty} w_{\text{mo}}(f)(t_{n_k}) = w_{\text{mo}}(f)(t_0)$.

Given $\varepsilon > 0$ we find $I_n \subset \mathbb{T}$ such that $0 < |I_n| \leq t_n$ and

$$w_{\text{mo}}(f)(t_n) \leq \frac{1}{|I_n|} \int_{I_n} |f(e^{i\theta}) - m_{I_n}(f)| \frac{d\theta}{2\pi} + \varepsilon.$$

Let $\mathcal{F} = \{n \in \mathbb{N} : |I_n| > t_0\}$. If \mathcal{F} is finite then $|I_n| \leq t_0$ for $n \geq n_0$ and

$$w_{\text{mo}}(f)(t_n) - w_{\text{mo}}(f)(t_0) < \varepsilon \quad \text{for } n \geq n_0.$$

Without loss of generality we assume $|I_n| > t_0$ for all $n \in \mathbb{N}$.

Call $I_0 = \cup_{n=1}^\infty \cap_{k=n}^\infty I_k$. It is easy to see that I_0 is an arc and that $|I_0| = t_0$. Take a subsequence n_k such that (I_{n_k}) converges to I_0 . We have

$$\begin{aligned} w_{\text{mo}}(f)(t_{n_k}) - w_{\text{mo}}(f)(t_0) &\leq w_{\text{mo}}(f)(t_{n_k}) - \frac{1}{|I_0|} \int_{I_0} |f(e^{i\theta}) - m_{I_0}(f)| \frac{d\theta}{2\pi} \\ &\leq \frac{1}{|I_{n_k}|} \int_{I_{n_k}} |f(e^{i\theta}) - m_{I_{n_k}}(f)| \frac{d\theta}{2\pi} - \frac{1}{|I_0|} \int_{I_0} |f(e^{i\theta}) - m_{I_0}(f)| \frac{d\theta}{2\pi} + \varepsilon. \end{aligned}$$

The proof is complete invoking Lemma 2.1. □

Remark 2.3. Let $f \in \text{BMO}$ and take $a(f) = \lim_{t \rightarrow 0^+} w_{\text{mo}}(f)(t)$. Hence $f \in \text{VMO}$ if and only if $a(f) = 0$.

For each $0 < p < \infty$, we define the quasi-norm (norm for $p \geq 1$) on $\text{MO}^p(\mathbb{T})$ by

$$\|f\|_{\text{MO}^p} = \|f\|_{L^1(\mathbb{T})} + \left(\int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t} \right)^{1/p}.$$

Although the next result is probably known, we include a proof for the sake of completeness.

Theorem 2.4. *Let $0 < p < \infty$. Then $(\text{MO}^p(\mathbb{T}), \|\cdot\|_{\text{MO}^p})$ is a complete space.*

Proof. Let $\{f_n\}$ be a Cauchy sequence in $\text{MO}^p(\mathbb{T})$. In particular, there exists $f \in \text{BMO}$ such that $\{f_n\}$ converges to f .

Let $|I| \leq t$, $0 < t \leq 1$. Using that $f_n \rightarrow f$ in $L^1(\mathbb{T})$ we get that $m_I(f_n) \rightarrow m_I(f)$ and that there exists a subsequence (n_k) , such that $f_{n_k} \rightarrow f$ a.e.

Now

$$\begin{aligned} \frac{1}{|I|} \int_I |f_n(e^{i\theta}) - f(e^{i\theta}) - m_I(f_n - f)| \frac{d\theta}{2\pi} &= \frac{1}{|I|} \int_I |f_n(e^{i\theta}) - \lim_k f_{n_k}(e^{i\theta}) - \lim_k m_I(f_n - f_{n_k})| \frac{d\theta}{2\pi} \\ &= \frac{1}{|I|} \int_I \lim_k |f_n(e^{i\theta}) - f_{n_k}(e^{i\theta}) - m_I(f_n - f_{n_k})| \frac{d\theta}{2\pi} \\ &\leq \liminf_k \frac{1}{|I|} \int_I |f_n(e^{i\theta}) - f_{n_k}(e^{i\theta}) - m_I(f_n - f_{n_k})| \frac{d\theta}{2\pi} \\ &\leq \liminf_k w_{\text{mo}}(f_n - f_{n_k})(t). \end{aligned}$$

Therefore

$$w_{\text{mo}}(f_n - f)(t) \leq \liminf_k w_{\text{mo}}(f_n - f_{n_k})(t).$$

Hence

$$\begin{aligned} \int_0^1 \left[w_{\text{mo}}(f_n - f)(t) \frac{dt}{t} \right]^p &\leq \int_0^1 \liminf_k [w_{\text{mo}}(f_n - f_{n_k})(t)]^p \frac{dt}{t} \\ &\leq \liminf_k \int_0^1 [w_{\text{mo}}(f_n - f_{n_k})(t)]^p \frac{dt}{t} \end{aligned}$$

Finally, using that f_n is a Cauchy sequence we get $\lim_{n \rightarrow \infty} \|f_n - f\|_{\text{MO}^p} = 0$ and that $f \in \text{MO}^p$. □

Proposition 2.5. *Let $0 < p \leq q < \infty$ and $s > 0$.*

- (i) $\text{MO}^p(\mathbb{T}) \subseteq \text{MO}^q(\mathbb{T})$.
- (ii) $\text{Lip}_s(\mathbb{T}) \subset \bigcap_{p>0} \text{MO}^p(\mathbb{T}) \subset \text{MO}^1(\mathbb{T}) \subset C(\mathbb{T})$.
- (iii) $\bigcup_{p>0} \text{MO}^p(\mathbb{T}) \subset \text{VMO}$.

Proof. (i) It is a consequence of the following fact:

$$\left(\int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t} \right)^{1/p} \approx \left(\sum_{k=0}^{\infty} [w_{\text{mo}}(f)(2^{-k})]^p \right)^{1/p}.$$

(ii) Note that $f \in \text{Lip}_s$ if and only if $w_{\text{mo}}(f)(t) \leq Ct^s$. This gives the first inclusion.

The fact that $\text{MO}^1(\mathbb{T}) \subset C(\mathbb{T})$ follows by (5).

(iii) Observe that, for any $p > 0$ and $t > 0$, one has

$$w_{\text{mo}}(f)^p(t) \log \frac{1}{t} \leq \int_t^1 w_{\text{mo}}(f)^p(u) \frac{du}{u} \leq \|f\|_{\text{MO}^p}.$$

Hence $\lim_{t \rightarrow 0^+} w_{\text{mo}}(f)(t) = 0$ for $f \in \bigcup_{p>0} \text{MO}^p(\mathbb{T})$. □

Let us point out some properties of BMO that are shared by these spaces.

Proposition 2.6. *If $f \in \text{MO}^p(\mathbb{T})$ then $|f| \in \text{MO}^p(\mathbb{T})$.*

Proof. Let $t \in (0, 1)$ and $I \subset \mathbb{T}$ with $|I| \leq t$. Then

$$\begin{aligned} \frac{1}{|I|} \int_I ||f(e^{i\theta})| - m_I(|f||) \frac{d\theta}{2\pi} &\leq \frac{1}{|I|} \int_I ||f(e^{i\theta})| - |m_I(f)|| \frac{d\theta}{2\pi} \\ &\quad + |m_I(|f|) - |m_I(f)|| \\ &\leq \frac{2}{|I|} \int_I ||f(e^{i\theta})| - |m_I(f)|| \frac{d\theta}{2\pi} \\ &\leq \frac{2}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \end{aligned}$$

This shows that $w_{\text{mo}}(|f|)(t) \leq 2 w_{\text{mo}}(f)(t)$ and the proof is complete. □

Recall that T_t denotes the translation operator, that is $T_t f(e^{i\theta}) = f(e^{i(\theta-t)})$. We have the following result.

Theorem 2.7. *Let $0 < p < \infty$ and $f \in \text{MO}^p(\mathbb{T})$. Then*

$$\lim_{s \rightarrow 0^+} \|T_s f - f\|_{\text{MO}^p} = 0.$$

Proof. Due to (iii) in Proposition 2.5 $f \in \text{VMO}$. Now Theorem 1.1 gives that $\lim_{s \rightarrow 0^+} \|T_s f - f\|_{\text{BMO}} = 0$.

Note that $w_{\text{mo}}(T_s f - f)(t) \leq \|T_s f - f\|_{\text{BMO}}$ for all $0 < t \leq 1$.

On the other hand

$$\begin{aligned} w_{\text{mo}}(T_s f - f)(t) &= \sup_{|I| \leq t} \frac{1}{|I|} \int_I |(T_s f - f)(e^{i\theta}) - m_I(T_s f - f)| \frac{d\theta}{2\pi} \\ &\leq \sup_{|I| \leq t} \frac{1}{|I|} \int_I |T_s f(e^{i\theta}) - m_I(T_s f)| \frac{d\theta}{2\pi} \\ &\quad + \sup_{|I| \leq t} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \\ &= 2 w_{\text{mo}}(f)(t) \end{aligned}$$

The Lebesgue dominated convergence theorem gives $\lim_{s \rightarrow 0^+} \|T_s f - f\|_{\text{MO}^p} = 0$. □

3. Integrable harmonic oscillation.

Throughout this section, given $z \in \Delta \setminus \{0\}$, we denote by I_z the open arc in \mathbb{T} with midpoint $\frac{z}{|z|}$ and length $|I_z| = 1 - |z|$. Given an arc $I \subset \mathbb{T}$ and $\lambda \leq |I|^{-1}$ we shall write λI for the arc with the same midpoint and length $\lambda|I|$.

Let us collect several known facts to be used later on.

Lemma 3.1. *There exist constants $0 < C, C_1, C_2, C_3 < \infty$ such that*

- (i) $1 - |z| \leq |e^{i\theta} - z| \leq C(1 - |z|)$, $e^{i\theta} \in I_z$, and $z \in \Delta$.
- (ii) $C_1 \frac{1}{|I_z|} \leq P_z(e^{i\theta}) \leq C_2 \frac{1}{|I_z|}$, $e^{i\theta} \in I_z$, and $z \in \Delta$.
- (iii) $\frac{1}{4^k |I_z|} \leq P_z(e^{i\theta}) \leq C_3 \frac{1}{4^k |I_z|}$, $e^{i\theta} \in 2^k I_z \setminus 2^{k-1} I_z$, $k \in \{1, 2, \dots, N + 1\}$, where $N = \lceil \log_2 \frac{1}{|I_z|} \rceil$ and $z \in \Delta$.

Proof. All the statements follow from the estimates

$$1 - |z| \leq |e^{i\theta} - z| \leq \left| e^{i\theta} - \frac{z}{|z|} \right| + (1 - |z|)$$

and

$$\left| e^{i\theta} - \frac{z}{|z|} \right| \leq |e^{i\theta} - z| + (1 - |z|). \quad \square$$

For $0 < p < \infty$ we define

$$\|f\|_{\text{HO}^p} = \|f\|_{L^1(\mathbb{T})} + \left(\int_0^1 [w_{\text{ho}}(f)(t)]^p \frac{dt}{t} \right)^{1/p}$$

to get a quasi-norm in the space $\text{HO}^p(\mathbb{T})$.

Proposition 3.2. *If $f \in L^1(\mathbb{T})$ and $0 < t \leq 1$ then $w_{\text{mo}}(f)(t) \leq c w_{\text{ho}}(f)(t)$.*

Proof. Let $I \subseteq \mathbb{T}$ be an arc such that $|I| \leq t$. Consider $z \in \Delta$ for which $I = I_z$. From $|I_z| = 1 - |z| \leq t$ we have $1 - t \leq |z| < 1$.

Using (ii) in Lemma 3.1 we have

$$\begin{aligned} \frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} &\leq \frac{1}{|I_z|} \int_{I_z} |f(e^{i\theta}) - P(f)(z)| \frac{d\theta}{2\pi} \\ &\quad + |m_I(f) - P(f)(z)| \\ &\leq \frac{2}{|I_z|} \int_{I_z} |f(e^{i\theta}) - P(f)(z)| \frac{d\theta}{2\pi} \\ &\leq C \left(\int_{-\pi}^{\pi} |f(e^{i\theta}) - P(f)(z)| P_z(\theta) \frac{d\theta}{2\pi} \right) \\ &\leq C w_{\text{ho}}(f)(t) \end{aligned}$$

Now taking the supremum over all arcs we get $w_{\text{mo}}(f)(t) \leq C w_{\text{ho}}(f)(t)$. □

Theorem 3.3. *Let $0 < p < \infty$. Then $\text{HO}^p(\mathbb{T}) = \text{MO}^p(\mathbb{T})$ with equivalent quasi-norms.*

Proof. $\text{HO}^p(\mathbb{T}) \subseteq \text{MO}^p(\mathbb{T})$ follows from Proposition 3.2.

Assume now that $f \in \text{MO}^p(\mathbb{T})$. Let us show that $f \in \text{HO}^p(\mathbb{T})$ and $\|f\|_{\text{HO}^p} \leq C\|f\|_{\text{MO}^p}$.

Let $t \in (0, 1]$. For any $z \in \Delta$ with $|z| = 1 - t$ we consider the arc $I = I_z$, which gives $|I_z| = 1 - |z| = t$. Take N and I_k ($k = 0, 1, \dots, N$) so that $I_k = 2^k I_z$ where $I_0 = I$, $I_{N-1} \subsetneq \mathbb{T}$, and $I_N = \mathbb{T}$.

Using (iii) in Lemma 3.1 we have

$$\begin{aligned} & \int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} \\ & \leq 2 \int_{\mathbb{T}} |f(e^{i\theta}) - m_I(f)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} \\ & \leq C \left(\int_I |f(e^{i\theta}) - m_I(f)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} \right. \\ & \quad \left. + \sum_{k=1}^N \int_{I_k \setminus I_{k-1}} |f(e^{i\theta}) - m_I(f)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} \right) \\ & \leq C \left(\frac{1}{|I|} \int_I |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \right. \\ & \quad \left. + \sum_{k=1}^N \frac{1}{4^k |I|} \int_{I_k \setminus I_{k-1}} |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \right) \\ & \leq C \left(w_{\text{mo}}(f)(t) + \sum_{k=1}^N \frac{1}{2^k |I_k|} \int_{I_k} |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \right). \end{aligned}$$

On the other hand, by (4),

$$\begin{aligned} & \frac{1}{|I_k|} \int_{I_k} |f(e^{i\theta}) - m_I(f)| \frac{d\theta}{2\pi} \\ & \leq \frac{1}{|I_k|} \int_{I_k} \left| f(e^{i\theta}) - m_{I_k}(f) + \left(\sum_{j=1}^k m_{I_j}(f) - m_{I_{j-1}}(f) \right) \right| \frac{d\theta}{2\pi} \\ & \leq \frac{1}{|I_k|} \int_{I_k} |f(e^{i\theta}) - m_{I_k}(f)| \frac{d\theta}{2\pi} + \sum_{j=1}^k |m_{I_j}(f) - m_{I_{j-1}}(f)| \\ & \leq \frac{1}{|I_k|} \int_{I_k} |f(e^{i\theta}) - m_{I_k}(f)| \frac{d\theta}{2\pi} + \sum_{j=1}^k \frac{|I_j|}{|I_{j-1}|} w_{\text{mo}}(f)(|I_j|) \\ & \leq w_{\text{mo}}(f)(|I_k|) + \sum_{j=1}^k 2w_{\text{mo}}(f)(|I_j|) \\ & \leq (1 + 2k)w_{\text{mo}}(f)(|I_k|). \end{aligned}$$

Combining both estimates one gets

$$\int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} \leq C \left(w_{\text{mo}}(f)(t) + \sum_{k=1}^N \frac{1+2k}{2^k} w_{\text{mo}}(f)(|I_k|) \right).$$

Taking the supremum over $\{z : |z| = 1 - t\}$ and using $|I_k| = 2^k t$ we obtain

$$\sup_{|z|=1-t} \int_{\mathbb{T}} |f(e^{i\theta}) - P(f)(z)| P_z(e^{i\theta}) \frac{d\theta}{2\pi} \leq C \left(w_{\text{mo}}(f)(t) + \sum_{k=1}^{N_t} \frac{1+2k}{2^k} w_{\text{mo}}(f)(2^k t) \right)$$

where $N = N_t = \lceil \log_2 \frac{1}{t} \rceil + 1$. This implies that

$$w_{\text{ho}}(f)(t) \leq C \left(w_{\text{mo}}(f)(t) + \sum_{k=1}^{N_t} \frac{1+2k}{2^k} w_{\text{mo}}(f)(2^k t) \right).$$

For $0 < p < 1$ we have

$$[w_{\text{ho}}(f)(t)]^p \leq C_p \left([w_{\text{mo}}(f)(t)]^p + \sum_{k=1}^{N_t} \frac{(1+2k)^p}{2^{pk}} [w_{\text{mo}}(f)(2^k t)]^p \right).$$

For $p \geq 1$ we apply Hölder's inequality to obtain

$$[w_{\text{ho}}(f)(t)]^p \leq C_p \left([w_{\text{mo}}(f)(t)]^p + \sum_{k=1}^{N_t} \frac{(1+2k)^p}{2^k} [w_{\text{mo}}(f)(2^k t)]^p \right).$$

Now integrating, and taking into account that $1 \leq k \leq N_t = \lceil \log_2 \frac{1}{t} \rceil + 1$ is equivalent to $0 < t \leq 2^{-k}$, we get

$$\begin{aligned} \int_0^1 [w_{\text{ho}}(f)(t)]^p \frac{dt}{t} &\leq C_p \int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t} + C_p \int_0^1 \sum_{k=1}^{N_t+1} \frac{(1+2k)^p}{2^{k \min\{p,1\}}} [w_{\text{mo}}(f)(2^k t)]^p \frac{dt}{t} \\ &\leq C_p \|f\|_{MO^p}^p + C_p \sum_{k=1}^{\infty} \frac{(1+2k)^p}{2^{k \min\{p,1\}}} \int_0^{2^{-k}} [w_{\text{mo}}(f)(2^k t)]^p \frac{dt}{t} \\ &\leq C_p \|f\|_{MO^p}^p + C_p \sum_{k=1}^{\infty} \frac{(1+2k)^p}{2^{k \min\{p,1\}}} \int_0^1 [w_{\text{mo}}(f)(t)]^p \frac{dt}{t} \\ &\leq C \|f\|_{MO^p}^p. \end{aligned}$$

Putting together all the estimates we have the result. □

References

- [1] A. Baernstein II, *Analytic functions of bounded mean oscillation*, Aspects of contemporary complex analysis (Proc. NATO Adv. Study Inst., Univ. Durham, Durham, 1979), 1980, pp. 3–36.
- [2] S. Campanato, *Proprietà di hölderianità di alcune classi di funzioni*, Ann. Scuola Norm. Sup. Pisa (3) **17** (1963), 175–188.
- [3] ———, *Proprietà di una famiglia di spazi funzionali*, Ann. Scuola Norm. Sup. Pisa (3) **18** (1964), 137–160.
- [4] J. R. Dorrnsoro, *Mean oscillation and Besov spaces*, Canad. Math. Bull. **28** (1985), no. 4, 474–480.
- [5] K. M. Dyakonov, *Besov spaces and outer functions*, Michigan Math. J. **45** (1998), no. 1, 143–157.
- [6] C. Fefferman, *Characterizations of bounded mean oscillation*, Bull. Amer. Math. Soc. **77** (1971), 587–588.
- [7] J. B. Garnett, *Bounded analytic functions*, Pure and Applied Mathematics, vol. 96, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1981.
- [8] D. Girela, *Analytic functions of bounded mean oscillation*, Complex function spaces (Mekrijärvi, 1999), 2001, pp. 61–170.
- [9] H. Greenwald, *On the theory of homogeneous Lipschitz spaces and Campanato spaces*, Pacific J. Math. **106** (1983), no. 1, 87–93.
- [10] S. Janson, *Generalizations of Lipschitz spaces and an application to Hardy spaces and bounded mean oscillation*, Duke Math. J. **47** (1980), no. 4, 959–982.
- [11] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426.
- [12] J. Peetre, *On the theory of \mathcal{L}_p, λ spaces*, J. Functional Analysis **4** (1969), 71–87.
- [13] F. Ricci and M. Taibleson, *Boundary values of harmonic functions in mixed norm spaces and their atomic structure*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **10** (1983), no. 1, 1–54.
- [14] D. Sarason, *Function theory on the unit circle*, Virginia Polytechnic Institute and State University Department of Mathematics, Blacksburg, Va., 1978.
- [15] W. S. Smith, *BMO(ρ) and Carleson measures*, Trans. Amer. Math. Soc. **287** (1985), no. 1, 107–126.
- [16] S. Spanne, *Some function spaces defined using the mean oscillation over cubes*, Ann. Scuola Norm. Sup. Pisa (3) **19** (1965), 593–608.
- [17] B. E. Viviani, *An atomic decomposition of the predual of BMO(ρ)*, Rev. Mat. Iberoamericana **3** (1987), no. 3-4, 401–425.