

## ASYMPTOTIC BEHAVIOR OF ORTHOGONAL POLYNOMIALS PRIMITIVES

ALFREDO FUNDORA ROLO, HÉCTOR PIJEIRA CABRERA AND  
WILFREDO URBINA ROMERO

*En recuerdo de nuestro cariño y amistad con Chicho*

ABSTRACT. We study the zero location and the asymptotic behavior of the primitives of the standard orthogonal polynomials with respect to a finite positive Borel measure concentrate on  $[-1, 1]$ .

### 1. INTRODUCTION

Let  $\mu$  be a finite positive Borel measure with  $\text{supp}(\mu) = \Delta \subseteq [-1, 1]$ , such that it contains an infinite number of points. Let us consider  $L_n(z) = z^n + \dots$  the  $n$ th monic (i.e. its leading coefficient is equal to one) orthogonal polynomial with respect to  $\mu$ , that is

$$(1) \quad \int_{\Delta} L_n(x) x^k d\mu(x) = 0, \quad k = 0, 1, 2, \dots, n-1.$$

Let us consider a monic polynomial  $P_n(x)$  of degree  $n$  and a complex number  $\zeta$  fixed, such that

$$(2) \quad (n+1)L_n(z) = ((z-\zeta)P_n(z))' = P_n(z) + (z-\zeta)P_n'(z).$$

Note that  $\Lambda(z) = (z-\zeta)P_n(z)$  is a monic polynomial primitive of  $(n+1)L_n(z)$ , normalized by  $\Lambda(\zeta) = 0$ . A direct consequence of (1)–(2) is that  $P_n(z)$  satisfy the orthogonality relations

$$(3) \quad \int_{\Delta} [P_n(x) + (x-\zeta)P_n'(x)] x^k d\mu(x) = 0, \quad k = 0, 1, 2, \dots, n-1.$$

The location of critical points of polynomials has many physical and geometrical interpretations. Let us consider, for instance, a field of forces given by a system of  $n$  masses  $m_j$ ,  $1 \leq j \leq n$ , at the fixed points  $z_j$ ,  $1 \leq j \leq n$ , that repels a movable unit mass at  $z$  according to the law of repulsion being the inverse distance law.

---

2000 *Mathematics Subject Classification.* 42C05, 33C25.

*Key words and phrases.* orthogonal polynomials, zero location, asymptotics.

Research of second and third authors partially supported by a research grant # PI 03-14-4744-2000 from Consejo de Desarrollo Científico y Humanístico (CDCH) of the Universidad Central de Venezuela.

Let  $Q_m(z)$ , where  $m = m_1 + m_2 + \dots + m_n$ , be the polynomial  $(z - z_1)^{m_1} \cdot (z - z_2)^{m_2} \dots (z - z_n)^{m_n}$ . The logarithmic derivative of  $Q_m(z)$  is

$$(4) \quad \frac{d(\log(Q_m(z)))}{dz} = \frac{Q'_m(z)}{Q_m(z)} = \frac{m_1}{(z - z_1)} + \frac{m_2}{(z - z_2)} + \dots + \frac{m_n}{(z - z_n)}.$$

The conjugate of  $\frac{m_j}{(z - z_j)}$  is a vector whose direction (including sense) is the direction from  $z_j$  to  $z$ , so this vector represents the force at the movable unit mass  $z$  due to a single fixed particle at  $z_j$ . Every multiple zero (but no simple zero) of  $Q_m(z)$  is a zero of  $Q'_m(z)$ ; every other zero of  $Q'_m(z)$  is by (4) a position of equilibrium in the field of force; every position of equilibrium is by (4) a zero of  $Q'_m(z)$ . This result is known as Gauss's theorem (1816).

Now, we consider an inverse problem, let  $z'_1, z'_2, \dots, z'_n$  be the zeros of the orthogonal polynomial  $L_n$  and the equilibrium positions of a field of forces with  $n + 1$  units masses, one of which  $\zeta$  is given. What is the location of the remaining masses?

By (2),

$$(5) \quad \frac{(n + 1) L_n(z)}{(z - \zeta)P_n(z)} = \frac{1}{z - \zeta} + \frac{P'_n(z)}{P_n(z)} = \frac{\Lambda'(z)}{\Lambda(z)}.$$

Then, according with (5) and the above interpretation of the logarithmic derivative, the location of the remaining units masses are the zeros of the polynomial  $P_n(z)$  defined in (2).

The main purpose of this paper is to study some of the algebraic and analytic properties of the orthogonal polynomials primitives.

## 2. LOCALIZATION OF ZEROS

It is well know that the zeros of  $L_n(z)$  are simple, using (2) is easy to see that the zeros of  $P_n(z)$  have at most multiplicity two. Nevertheless the zeros of  $P_n(z)$  need not to be simple as we can see in the following example

Let  $\mu$  be the Lebesgue measure in  $[-1, 1]$  and set in (3)  $\zeta = \frac{2\sqrt{3}}{3}$  or  $\zeta = -\frac{2\sqrt{3}}{3}$ . The corresponding monic polynomials of degree two defined by (2) are  $P_2(z) = z^2 + \frac{2\sqrt{3}}{3}z + \frac{1}{3}$  or  $P_2(z) = z^2 - \frac{2\sqrt{3}}{3}z + \frac{1}{3}$  respectively. Note that  $z = -\frac{\sqrt{3}}{3}$  or  $z = \frac{\sqrt{3}}{3}$  are zeros of multiplicity two of the corresponding polynomials  $P_2(z)$ .

Our next propose is to prove that all the zeros of the polynomials of the sequence  $\{P_n(z)\}_{n=0}^\infty$  are contained in a disc which radius is independent of  $n$ . First, let us rewrite the polynomials  $P_n$  and  $L_n$  in terms of  $(z - \zeta)$ , that is

$$(6) \quad P_n(z) = \sum_{k=0}^n a_k (z - \zeta)^k, \quad L_n(z) = \sum_{k=0}^n b_k (z - \zeta)^k.$$

**Lemma 1.** *The coefficients  $a_k$  of  $P_n$  and  $b_k$  of  $L_n$  in (6) are related by*

$$(7) \quad a_k = \frac{n + 1}{k + 1} b_k.$$

*Proof.* Replacing (6) in (2). □

The proof of the next result is based in the following Szegő's theorem (see [5] or [2, page 23]).

**Lemma 2.** *Given the polynomials*

$$f(z) = \sum_{k=0}^n \alpha_k \binom{n}{k} z^k, \quad \alpha_n \neq 0 \quad \text{and} \quad g(z) = \sum_{k=0}^n \beta_k \binom{n}{k} z^k, \quad \beta_n \neq 0,$$

let us construct a third polynomial as  $h(z) = \sum_{k=0}^n \alpha_k \beta_k \binom{n}{k} z^k$ .

If all the zeros of  $f(z)$  lie in a closed disk  $\bar{D}$  and  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the zeros of  $g(z)$ . Then every zero of  $h(z)$  has the form  $\lambda_k \gamma_k$ , where  $\gamma_k \in \bar{D}$ .

Then we have that

**Theorem 1.** *All the zeros of  $P_n$  are contained in the closed disk  $\mathbf{D}$ , where*

$$(8) \quad \mathbf{D} = \{z \in \mathbb{C} : |z| \leq 2 + 3|\zeta|\}.$$

*Proof.* Let us write  $w = z - \zeta$ , hence

$$f(w) = \sum_{k=0}^n b_k w^k = L_n(z), \quad h(w) = \sum_{k=0}^n \frac{n+1}{k+1} b_k w^k = P_n(z)$$

and

$$g(w) = \sum_{k=0}^n \frac{n+1}{k+1} \binom{n}{k} w^k = \frac{(1+w)^{n+1} - 1}{w} = \frac{(1+z-\zeta)^{n+1} - 1}{z-\zeta}.$$

If  $z_0$  is a zero of  $L_n$ , it is well known that  $-1 \leq z_0 \leq 1$ , hence  $w_0 = z_0 - \zeta$  is a zero of  $f(w)$  and lie in a closed disk  $\bar{D} = \{|w + \zeta| \leq 1\}$ . On the other hand, if  $w_1$  is a zero of  $g(w)$  then  $|1 + w_1| = 1$ .

Finally, by Lemma 2, if  $h(w_3) = 0$  we have that  $|w_3| \leq 2 + 3|\zeta|$  and then the theorem is proved. □

### 3. AUXILIARY RESULTS

In order to obtain the asymptotic behaviour of the sequence  $\{P_n\}$  we need some general results that we will discuss in what follows.

If  $\{\mu_n\}_{n=1}^\infty$  is a sequence of measures on a compact set, we say that  $\mu_n$  converges weakly to the measure  $\mu$  as  $n \rightarrow \infty$  if

$$\lim_{n \rightarrow \infty} \int f d\mu_n = \int f d\mu$$

for every continuous function  $f$  on  $\mathbb{C}$  having compact support. In this case, we write  $\mu_n \xrightarrow{*} \mu$ , or  $d\mu_n \xrightarrow{*} d\mu$ , or if  $\mu$  is absolutely continuous,  $d\mu_n(x) \xrightarrow{*} \mu'(x)dx$ .

For any polynomial  $q$  of degree exactly  $n$ , we consider

$$\nu_n(q) := \frac{1}{n} \sum_{j=1}^n \delta_{z_j},$$

where  $z_1, \dots, z_n$  are the zeros of  $q$  repeated according to their multiplicity, and  $\delta_{z_j}$  is the Dirac measure with mass one at the point  $z_j$ . This is the so called *normalized zero counting measure associated with  $q$* .

Let  $\|\cdot\|_\Delta$  denotes the supremum norm on  $\Delta$  and  $\text{Cap}(\Delta)$  the logarithmic capacity of a set  $\Delta$ . Another result needed is

**Lemma 3** ([1], Theorem 2.1 and Corollary 2.1). *Let  $\Delta \subset \mathbb{C}$  be a compact set with empty interior, connected complement and positive logarithmic capacity. If  $\{P_n\}_{n=0}^\infty$  is a sequence of monic polynomials,  $\deg(P_n) = n$ , such that*

$$\overline{\lim}_{n \rightarrow \infty} \|P_n\|_\Delta^{\frac{1}{n}} \leq \text{Cap}(\Delta),$$

then

$$\nu_n(P_n) \xrightarrow{*} \omega_\Delta,$$

where  $\omega_\Delta$  is the equilibrium measure of  $\Delta$ .

Finally, we have the following useful result

**Lemma 4** ([3], Lemma 3). *Let  $\{P_n\}$  be a sequence of polynomials. Then, for all  $j \in \mathbb{Z}_+$ ,*

$$(9) \quad \overline{\lim}_{n \rightarrow \infty} \left( \frac{\|P_n^{(j)}\|_\Delta}{\|P_n\|_\Delta} \right)^{1/n} \leq 1.$$

For  $\Delta = [-1, 1]$  is well known that  $\text{Cap}(\Delta) = \frac{1}{2}$  and the equilibrium measure on  $\Delta$  is the so-called arcsin measure given by

$$(10) \quad \mu_\Delta(B) = \int_B \frac{\arcsin'(x) dx}{\pi} = \frac{1}{\pi} \int_B \frac{dx}{\sqrt{1-x^2}},$$

where  $B$  is a Borel set in  $[-1, 1]$ .

#### 4. ASYMPTOTIC BEHAVIOR

Let us set  $\varphi(z) = z + \sqrt{z^2 - 1}$ ,  $z \in \mathbb{C} \setminus [-1, 1]$ .  $\varphi$  is a conformal map of  $\mathbb{C} \setminus [-1, 1]$  onto  $\{z \in \mathbb{C} : |z| > 1\}$ . Here the branch of the square root is chosen so that  $|z + \sqrt{z^2 - 1}| > 1$  for  $z \in \mathbb{C} \setminus [-1, 1]$ . Let  $\zeta \in \mathbb{C} \setminus [-1, 1]$  be a fixed point,  $\Omega = \mathbb{C} \setminus \mathbf{D}$  and  $\Delta = [-1, 1]$ .

**Theorem 2.** *With the previous conditions it holds, for all  $j \in \mathbb{Z}_+$ ,*

- the sequence  $\{P_n^{(j)}\}_{n=0}^\infty$  verifies

$$(11) \quad \lim_{n \rightarrow \infty} \|P_n^{(j)}\|_\Delta^{\frac{1}{n}} = \frac{1}{2};$$

- $\nu_{n,j}(P_n^{(j)})$  converges to the arcsin measure in the sense of the weak-\* topology of measures, that is

$$(12) \quad \lim_{n \rightarrow \infty} \frac{1}{n-j} \sum_{k=1}^{n-j} f(x_{n,k}^{(j)}) = \frac{1}{\pi} \int_{-1}^1 f(x) \frac{dx}{\sqrt{1-x^2}},$$

for every continuous function on  $\mathbf{D}$ , where  $\{x_{n,k}^{(j)}\}_{k=1}^{n-j}$  is the set of zeros of  $P_n^{(j)}(z)$ .

*Proof.* Let us prove first that

$$(13) \quad \lim_{n \rightarrow \infty} \|P_n\|_{\Delta}^{\frac{1}{n}} = \frac{1}{2}.$$

If  $x \in \Delta$ , integrating in (2) we have

$$(n + 1) \int_{\zeta}^x L_n(t) dt = (x - \zeta)P_n(x),$$

by taking absolute values both sides, we obtain

$$M(n + 1)\|L_n(x)\|_{\Delta} \geq (n + 1) \left| \int_{\zeta}^x L_n(t) dt \right| = |x - \zeta| |P_n(x)|, \geq m|P_n(x)|,$$

where  $m = \inf_{x \in \Delta} |x - \zeta|$  and  $M = \sup_{x \in \Delta} |x - \zeta|$ . Then

$$M(n + 1)\|L_n(x)\|_{\Delta} \geq m\|P_n\|_{\Delta} \geq m\|T_n\|_{\Delta},$$

where  $T_n$  is the  $n$ -th Chebyshev polynomial in  $[-1, 1]$ .

It is well known, for general theory of orthogonal polynomials, that

$$\lim_{n \rightarrow \infty} \|L_n\|_{\Delta}^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \|T_n\|_{\Delta}^{\frac{1}{n}} = \frac{1}{2},$$

hence we have (13).

By Lemma 4 and (13),

$$(14) \quad \overline{\lim}_{n \rightarrow \infty} \|P_n^{(j)}\|_{\Delta}^{\frac{1}{n}} = \overline{\lim}_{n \rightarrow \infty} \frac{\|P_n^{(j)}\|_{\Delta}^{\frac{1}{n}}}{\|P_n\|_{\Delta}^{\frac{1}{n}}} \|P_n\|_{\Delta}^{\frac{1}{n}} \leq \frac{1}{2} = \text{Cap}(\Delta).$$

But

$$(15) \quad \underline{\lim}_{n \rightarrow \infty} \|P_n^{(j)}\|_{\Delta}^{\frac{1}{n}} \geq \overline{\lim}_{n \rightarrow \infty} \|T_{n-j}\|_{\Delta}^{\frac{1}{n}} = \frac{1}{2} = \text{Cap}(\Delta)$$

and then (14) and (15) implies (11).

Finally, by Lemma 3 we deduce that (11) implies (12). □

**Theorem 3.** *With the above assumptions, it holds:*

- For all  $j \in \mathbb{Z}_+$ ,

$$(16) \quad \frac{P_n^{(j+1)}(z)}{nP_n^{(j)}(z)} \xrightarrow[n]{\Rightarrow} \frac{1}{\sqrt{z^2 - 1}}$$

*uniformly on compact subsets of  $\Omega$ .*

- (Relative Asymptotic) For all  $j_1, j_2 \in \mathbb{Z}_+$ ,

$$(17) \quad n^{j_2-j_1} \frac{L_n^{(j_1)}(z)}{P_n^{(j_2)}(z)} \xrightarrow[n]{\Rightarrow} \frac{z - \zeta}{\sqrt{z^2 - 1}} \left( \sqrt{z^2 - 1} \right)^{j_2-j_1}$$

*uniformly on compact subsets of  $\Omega$ .*

*Proof.* Let  $x_{n,k}^j$ ,  $k = 1, \dots, n - j$ , denote the  $n - j$  zeros of the polynomial  $P_n^{(j)}$ . It is known that all the critical points of a non-constant polynomials  $P_n$  and it's derivatives lied in the convex hull of his zeros, then by theorem 1  $x_{n,k}^j \in \mathbf{D} = \{z : |z| \leq 2 + 3|\zeta|\}$ ,  $k = 1, \dots, n - j$ . Using the decomposition in simple fractions and the definition of  $\nu_{n,j}(P_n^{(j)})$ , we obtain

$$(18) \quad \frac{P_n^{(j+1)}(z)}{nP_n^{(j)}(z)} = \frac{1}{n} \sum_{k=1}^{n-j} \frac{1}{z - x_{n,k}^j} = \frac{n-j}{n} \int \frac{d\nu_{n,j}(x)}{z-x}.$$

Therefore, the family of functions

$$(19) \quad \left\{ \frac{P_n^{(j+1)}(z)}{nP_n^{(j)}(z)} \right\}, \quad n \in \mathbb{Z}_+,$$

is uniformly bounded on each compact subset of  $\Omega = \mathbb{C} \setminus \mathbf{D}$ .

On the other hand, all the measures  $\nu_{n,j}$ ,  $n \in \mathbb{Z}_+$ , are supported in  $\mathbf{D}$  and for  $z \in \Omega$  fixed, the function  $(z - x)^{-1}$  is continuous on  $\mathbf{D}$  with respect to  $x$ . Therefore, from (12) and (18), we find that any subsequence of (19) which converges uniformly on compact subsets of  $\Omega$  converges pointwise to  $\int (z - x)^{-1} d\omega_\Delta(x)$ . Finally, by (10), the Cauchy's formula and the residue Theorem,

$$\int_{-1}^1 \frac{d\omega_\Delta(x)}{(z-x)} = \frac{1}{\pi} \int_{-1}^1 \frac{1}{(z-x)} \frac{dx}{\sqrt{1-x^2}} = \frac{1}{\sqrt{z^2-1}}.$$

Thus, the whole sequence converges uniformly on compact subsets of  $\Omega$  to this function as stated in (16).

For  $j_1 = j_2 = j$ , the proof of (17) is a direct consequence of the  $j$ -th derivative of (2) and (16), that is

$$(20) \quad \frac{n+1}{n} \frac{L_n^{(j)}(z)}{P_n^{(j)}(z)} = \frac{j+1}{n} + (z-\zeta) \frac{P_n^{(j+1)}(z)}{nP_n^{(j)}(z)} \xrightarrow{\frac{1}{n}} \frac{z-\zeta}{\sqrt{z^2-1}}$$

uniformly on compact subsets of  $\Omega$ .

Assume without loss of generality that  $j_2 < j_1$ , hence

$$(21) \quad \frac{1}{n^{j_1-j_2}} \frac{L_n^{(j_1)}(z)}{P_n^{(j_2)}(z)} = \frac{L_n^{(j_1)}(z)}{P_n^{(j_1)}(z)} \frac{P_n^{(j_1)}(z)}{nP_n^{(j_1-1)}(z)} \dots \frac{P_n^{(j_2+2)}(z)}{nP_n^{(j_2+1)}(z)} \frac{P_n^{(j_2+1)}(z)}{nP_n^{(j_2)}(z)}.$$

Then we have (17) from (16), (20) and (21). □

**Theorem 4.** *With the above conditions, the following statements hold:*

- (Strong Asymptotic) *If  $\mu'(x)$  satisfy the Szegő condition*

$$\int_{-1}^1 \frac{\log \mu'(x) dx}{\sqrt{1-x^2}} > -\infty$$

*then, for all  $j \in \mathbb{Z}_+$ ,*

$$(22) \quad \frac{P_n^{(j)}(z)}{n^j \left(\frac{\varphi(z)}{2}\right)^n} \xrightarrow{\frac{1}{n}} \frac{(\sqrt{z^2-1})^{1-j}}{z-\zeta} \frac{\mathcal{D}(\mu'(\cos \theta) |\sin \theta|, 0)}{\mathcal{D}(\mu'(\cos \theta) |\sin \theta|, \varphi^{-1}(z))},$$

uniformly on compact subsets of  $\Omega$ , where  $\mathcal{D}(h, z)$  is the Szegő function of  $h$

$$\mathcal{D}(h, z) = \exp \left( \frac{1}{4\pi} \int_0^{2\pi} \log h(\theta) \frac{e^{i\theta} + z}{e^{i\theta} - z} d\theta \right), \quad |z| < 1.$$

- (Ratio Asymptotic) If  $\mu'(x) > 0$  a.e. in  $[-1, 1]$  then, for all  $j_1, j_2, k \in \mathbb{Z}_+$ ,

$$(23) \quad \frac{n^{j_2} \frac{P_{n+k}^{(j_1)}(z)}{P_n^{(j_2)}(z)}}{(n+k)^{j_1}} \xrightarrow[n]{\Rightarrow} \left(\sqrt{z^2 - 1}\right)^{j_2 - j_1} \left(\frac{\varphi(z)}{2}\right)^k,$$

uniformly on compact subsets of  $\Omega$ .

- ( $n$ -th Root Asymptotic) If the measure  $\mu$  is such that for all measurable set  $E \subset \text{supp}(\mu)$  with  $\mu(E) = \mu([-1, 1])$  it holds that  $\text{Cap}(E) = \frac{1}{2}$ , then, for all  $j \in \mathbb{Z}_+$ ,

$$(24) \quad \sqrt[n]{|P_n^{(j)}(z)|} \xrightarrow[n]{\Rightarrow} \frac{|\varphi(z)|}{2},$$

uniformly on compact subsets of  $\Omega$ , where  $\Omega = \mathbb{C} \setminus \mathbf{D}$ ,  $\varphi(z) = z + \sqrt{z^2 - 1}$  and the branch of the square root is chosen so that  $|z + \sqrt{z^2 - 1}| > 1$  for  $z \in \mathbb{C} \setminus [-1, 1]$ .

*Proof.* The theorem is a direct consequence of (17) in theorem 3 and the well known strong asymptotic, ratio asymptotic and  $n$ -th root asymptotic behavior of standard orthogonal polynomials  $L_n$ . □

### REFERENCES

- [1] H. P. Blatt, E. B. Saff and M. Simkani, Jentzsch-Szegő type theorems for the zeros of best approximants, *J. London Math. Soc.* **38** (1988), 192–204.
- [2] P. Borwein and T. Erdélyi, *Polynomials and polynomials inequalities*, Graduate Text in Mathematics **161**, Springer-Verlag, New York, 1995.
- [3] G. López and H. Pijeira, Zero location and  $n$ -th root asymptotics of Sobolev orthogonal polynomials, *J. Approx. Theory* **99** (1999), 30-43.
- [4] H. Stahl and V. Totik, *General orthogonal polynomials*, Cambridge Univ. Press, Cambridge, 1992.
- [5] G. Szegő, Bemerkungen zu einen Satz von J. H. Grace über die Wurzeln algebraischer Gleichungen, *Math. Z.* **13** (1922), 28–55.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MATANZAS, AUTOPISTA DE VARADERO  
 KM. 3, 44740 MATANZAS, CUBA  
*E-mail address:* `alfredo@quimec.umtz.edu.cu`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MATANZAS, AUTOPISTA DE VARADERO  
 KM. 3, 44740 MATANZAS, CUBA  
*E-mail address:* `pijeira@cdict.umtz.edu.cu`

ESCUELA DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD CENTRAL DE VENEZUELA,  
 APT. 47195 LOS CHAGUARAMOS, CARACAS 1041-A, VENEZUELA  
*E-mail address:* `wurbina@euler.ciens.ucv.ve`