



Motivic and ℓ -adic realizations of the category of singularities of the zero locus of a global section of a vector bundle

Massimo Pippi¹

Accepted: 27 October 2021 / Published online: 5 January 2022
 © The Author(s) 2022

Abstract

We study the motivic and ℓ -adic realizations of the dg category of singularities of the zero locus of a global section of a line bundle over a regular scheme. We will then use the formula obtained in this way together with a theorem due to D. Orlov and J. Burke–M. Walker to give a formula for the ℓ -adic realization of the dg category of singularities of the zero locus of a global section of a vector bundle. In particular, we obtain a formula for the ℓ -adic realization of the dg category of singularities of the special fiber of a scheme over a regular local ring of dimension n .

Keywords Dg categories of singularities · Vanishing cycles · Motives · Non-commutative motives · ℓ -adic realization

Mathematics Subject Classification 14F42 · 19E08 · 32S30 · 14B05 · 14F08

Contents

1	Introduction	2
2	Preliminaries and notation	6
2.1	Some notation and convention	6
2.2	Reminders on dg categories	6
2.3	Derived algebraic geometry	8
2.4	ℓ -adic sheaves	9
2.5	Stable homotopy categories	10
2.5.1	The stable homotopy category of schemes	10
2.5.2	The stable homotopy category of non-commutative spaces	12
2.6	The bridge between motives and non-commutative motives	13
2.7	ℓ -adic realization of dg categories	14
3	Motivic realization of twisted LG models	15

✉ Massimo Pippi
 m.pippi@ucl.ac.uk

¹ Department of Mathematics, University College London, Gordon Street 25, London WC1H 0AY, UK

3.1	The category of twisted LG models	16
3.2	The dg category of singularities of a twisted LG model	19
3.3	The motivic realization of $\mathbf{Sing}(X, s_X)$	31
4	Monodromy-invariant vanishing cycles	40
4.1	The formalism of vanishing cycles	40
4.2	Monodromy-invariant vanishing cycles	43
5	The comparison theorem	46
5.1	2-periodic ℓ -adic sheaves	46
5.2	The main theorem	48
6	The ℓ -adic realization of the dg category of singularities of a twisted LG model of rank r	52
6.1	Reduction of codimension	52
6.2	The ℓ -adic realization of the dg category of singularities of a twisted LG model of rank r	63
7	Towards a vanishing cycles formalism over $\mathbb{A}_S^1/\mathbb{G}_{m,S}$	64
7.1	Tame vanishing cycles over \mathbb{A}_S^1	64
7.2	Tame vanishing cycles over $\mathbb{A}_S^1/\mathbb{G}_{m,S}$	65
8	Some remarks on the regularity hypothesis	68
	References	69

1 Introduction

The connection between categories of singularities and vanishing cycles is well known, thanks to works of Dyckerhoff [23], Preygel [45], Efimov [24] and many others. Recently, an instance of this fact has been studied in [9].

We start by quickly reviewing the main theorem of [9], that served both as a model and as a motivation for the investigations presented later. The main purpose of the above mentioned paper is to identify a classical object of singularity theory, namely the ℓ -adic sheaf of (inertia invariant) vanishing cycles, with the ℓ -adic cohomology of a non-commutative space (that is defined in *loc. cit.*), the dg category of singularities of the special fiber. The main result of Blanc et al. ([9, Theorem 4.39]) reads as follows:

Let $p : X \rightarrow S$ be a proper, flat, regular scheme over an excellent strictly henselian trait S . Let $i_\sigma : \sigma \hookrightarrow S$ be the embedding of the closed point in S and let $p_\sigma : X_\sigma \rightarrow \sigma$ the pullback of p along i_σ . Fix a prime number ℓ different from the characteristic of σ . Let $\Phi_p(\mathbb{Q}_{\ell,X}(\beta))$ be the ℓ -adic sheaf of vanishing cycles associated to $\mathbb{Q}_{\ell,X}(\beta) = \bigoplus_{n \in \mathbb{Z}} \mathbb{Q}_{\ell,X}(n)[2n]$. Denote by I the inertia group (as S is strictly henselian, it coincides with the absolute Galois group of the open point in S) and by $(-)^{hI}$ the (homotopy) fixed points ∞ -functor. In [9], the authors define the ℓ -adic realization of dg categories, denoted by $\mathcal{R}_S^{\ell,\vee}$. It is an ∞ -functor that associates an ℓ -adic complex to a dg category. For every (derived) scheme Z , let $\mathbf{Sing}(Z)$ be the dg category of singularities of Z , that is the dg quotient¹

$$\mathbf{Sing}(Z) := \mathbf{Coh}^b(Z)/\mathbf{Perf}(Z),$$

where $\mathbf{Coh}^b(Z)$ denotes the dg-category of complexes of quasi coherent \mathcal{O}_Z -modules with coherent total cohomology and $\mathbf{Perf}(Z)$ that of perfect complexes of \mathcal{O}_Z -modules.

¹ Notice that if Z is derived, we need to assume that $\mathcal{O}_Z \in \mathbf{Coh}^b(Z)$.

A. Blanc, M. Robalo, B. Toën and G. Vezzosi show that there is an equivalence of $i_{\sigma}^* \mathcal{R}_S^{\ell, \vee}(\mathbf{Sing}(S, 0)) \simeq \mathbb{Q}_{\ell, \sigma}(\beta) \otimes_{\mathbb{Q}_{\ell, \sigma}} \mathbb{Q}_{\ell, \sigma}^{hI}$ -modules


$$i_{\sigma}^* \mathcal{R}_S^{\ell, \vee}(\mathbf{Sing}(X_{\sigma})) \simeq p_{\sigma*} \Phi_p(\mathbb{Q}_{\ell, X}(\beta))^{hI}[-1].$$

Remark This theorem is of major importance in B. Toën - G. Vezzosi's approach to the Bloch's conductor conjecture [8]. For more details, see [67–69].

The content of this theorem is summarized in the following mind map:

NON COMMUTATIVE SIDE

VANISHING CYCLES SIDE



We can ask whether the above diagram makes sense in more general situations. For example, one can start with the datum of a proper, flat and regular scheme over an excellent local regular ring of dimension n . One recovers the case treated in [9] when $n = 1$.

It is immediate to observe that the left hand side of the diagram makes sense without any change: assume that we are given a proper, flat morphism $p : X \rightarrow S$, with X regular and S local, regular of dimension n . Let $\underline{\pi} = (\pi_1, \dots, \pi_n)$ be a collection of generators of the closed point of S . Then we can consider the morphism $\underline{\pi} \circ p : X \rightarrow \mathbb{A}_S^n$ and its fiber X_0 along the origin $S \rightarrow \mathbb{A}_S^n$. It makes perfectly sense to consider $\mathcal{R}_S^{\ell, \vee}(\mathbf{Sing}(X_0))$.

It comes up that this generalization is related to the following one: one can consider pairs (X, s_X) where X is a regular scheme over S and $s_X : X \rightarrow \mathbb{V}(\mathcal{L}_S)$ is a morphism towards the total space of a line bundle \mathcal{L}_S over S . One recovers the situation pictured above when $\mathcal{L}_S = \mathcal{O}_S$ is the trivial line bundle. In this situation, we want to compute $\mathcal{R}_S^{\ell, \vee}(\mathbf{Sing}(X_0))$, where X_0 is the fiber of $s_X : X \rightarrow \mathbb{V}(\mathcal{L}_S)$ along the zero section $S \rightarrow \mathbb{V}(\mathcal{L}_S)$.

One can view the former generalization as a particular case of the latter thanks to a theorem of Orlov [42] and Burke and Walker [11], which tells us that the dg category of singularities of $(X, \underline{\pi} \circ p)$ is equivalent to the dg category of singularities of $(\mathbb{P}_X^{n-1}, (\pi_1 \circ p) \cdot T_1 + \dots + (\pi_n \circ p) \cdot T_n \in \mathcal{O}(1)(\mathbb{P}_X^{n-1}))$.

Thus, we only need to find the appropriate generalization of (inertia invariant) vanishing cycles. The first thing we can think of are vanishing cycles over general

bases, developed by Laumon [33] following ideas of P. Deligne and further investigated by many others, including O. Gabber, L. Illusie, F. Orgogozo (see [29,41]). However, it seems that this is not the right point of view for our purposes. Instead, we will pursue the following analogy:

TOPOLOGY	VANISHING CYCLES	OUR SETTING
\mathbb{D} unital open disk	S strictly henselian trait	$\mathbb{V}(\mathcal{L})$ total space of a line bundle \mathcal{L} over a base scheme S
$0 \hookrightarrow \mathbb{D}$ origin	$\sigma \hookrightarrow S$ special point	$S \hookrightarrow \mathbb{V}(\mathcal{L})$ zero section
$\mathbb{D}^* = \mathbb{D} - \{0\} \hookrightarrow \mathbb{D}$ punctured disk	$\eta \hookrightarrow S$ generic point	$\mathcal{U} = \mathbb{V}(\mathcal{L}) - S \hookrightarrow \mathbb{V}(\mathcal{L})$ open complementary
$\widetilde{\mathbb{D}^*} \rightarrow \mathbb{D}^*$ universal cover of the punctured disk	$\bar{\eta} \rightarrow \eta$ separable closure of the generic point	?

As we will only need to define the analogous of inertia-invariant vanishing cycles, we will not face the problem of filling the empty spot in the mental map above. Nevertheless, we will come back to this matter at the end of the article, presenting a strategy to complete the picture.

We will define an appropriate generalization of $\Phi_p(\mathbb{Q}_{\ell,X}(\beta))^{\text{hI}}$ (see Definition 4.2.3) and prove a generalization of the formula stated in [9, Theorem 4.39]. Our main theorem will then look as follows:

Theorem 5.2.2 *Let X be a regular scheme and let s_X be a regular global section of a line bundle \mathcal{L}_X . Denote X_0 the zero locus of s_X . Then*

$$\mathcal{R}_{X_0}^{\ell,\vee}(\text{Sing}(X_0)) \simeq \Phi_{(X,s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}(\beta))[-1].$$

Here $\Phi_{(X,s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}(\beta))$ is what we call the monodromy-invariant vanishing cycles ℓ -adic sheaf (see Definition 4.2.6). It coincides with inertia invariant vanishing cycles when we put ourselves in the situation considered in [9].

Using this theorem combined with the above mentioned result of D. Orlov and J. Burke–M. Walker (Theorem 6.1.16), we will deduce the following formula for the

situation in which one starts with a regular scheme X and a regular section s_X of a vector bundle.

Theorem 6.2.1 *Let $f : X \rightarrow S$ be a flat morphism, \mathcal{E}_S be a vector bundle of rank r over S and let s_X be a global section of $\mathcal{E}_X = f^*\mathcal{E}_S$. Assume that X is a regular scheme and that s_X is regular section. The following equivalence holds in $\mathbf{Mod}_{\mathcal{R}_X^{\ell,\vee}(\mathbf{Sing}(X,0))}(\mathbf{Shv}_{\mathbb{Q}_\ell}(X))$*

$$\mathcal{R}_X^{\ell,\vee}(\mathbf{Sing}(X, s_X)) \simeq p_* i_* \Phi_{(\mathbb{P}(\mathcal{E}_X), W_{s_X})}^{\mathrm{mi}}(\mathbb{Q}_\ell(\beta))[-1],$$

where W_{s_X} is the global section of $\mathcal{O}_{\mathbb{P}(\mathcal{E}_X)}(1)$ associated to s_X , $i : V(W_{s_X}) \rightarrow \mathbb{P}(\mathcal{E}_X)$ is the closed embedding of the zero locus of W_{s_X} in $\mathbb{P}(\mathcal{E}_X)$ and $p : \mathbb{P}(\mathcal{E}_X) \rightarrow X$ is the canonical projection.

In particular, the above theorem applies to the special case $\mathcal{E}_S \simeq \mathcal{O}_S^n$ and allows us to compute the ℓ -adic realisation of the special fiber of a regular scheme over a local regular noetherian ring of dimension n . More precisely:

Corollary 6.2.2 *Assume that $S = \mathrm{Spec}(A)$ is a noetherian regular local ring of dimension n and let π_1, \dots, π_n be generators of the maximal ideal. Let $p : X \rightarrow S = \mathrm{Spec}(A)$ be a regular, flat S -scheme of finite type. Let $\underline{\pi} : S \rightarrow \mathbb{A}_S^n$ be the closed embedding associated to π_1, \dots, π_n . Then $\underline{\pi} \circ p$ is a regular global section of \mathcal{O}_X^n . Then the equivalence*

$$\mathcal{R}_X^{\ell,\vee}(\mathbf{Sing}(X, \underline{\pi} \circ p)) \simeq q_* i_* \Phi_{(\mathbb{P}_X^{n-1}, W_{\underline{\pi} \circ p})}^{\mathrm{mi}}(\mathbb{Q}_\ell(\beta))[-1]$$

holds in $\mathbf{Mod}_{\mathcal{R}_X^{\ell,\vee}(\mathbf{Sing}(X,0))}(\mathbf{Shv}_{\mathbb{Q}_\ell}(X))$.

Here $q : \mathbb{P}_X^{n-1} = \mathrm{Proj}_X(\mathcal{O}_X[T_1, \dots, T_n]) \rightarrow X$ is the canonical projection and $i : V(W_{\underline{\pi} \circ p}) \rightarrow \mathbb{P}_X^{n-1}$ is the closed embedding determined by the equation

$$W_{\underline{\pi} \circ p} = p^*(\pi_1) \cdot T_1 + \dots + p^*(\pi_n) \cdot T_n = 0.$$

We will end this article with some remarks on the following two problems:

1. It seems possible to define a formalism of vanishing cycles in twisted situations, i.e. in the situation in which we have a morphism $s_X : X \rightarrow \mathbb{V}(\mathcal{L}_X)$ for $\mathcal{L}_X \in \mathbf{Pic}(X)$. This is all about completing the empty slot in the mind map above. One should then be able to find $\Phi_{(X, s_X)}^{\mathrm{mi}}(\mathbb{Q}_\ell(\beta))$ via a procedure that corresponds to taking homotopy fixed points in the usual situation. A complete account on this formalism will appear in a forthcoming paper in collaboration with D.-C. Cisinski.
2. We will comment the regularity hypothesis that appears both in A. Blanc–M. Robalo–B. Toën–G. Vezzosi’s theorem and in the generalization we provide.

Remark The main body of this article corresponds to [44, Chapter §3], while the “preliminaries” section corresponds Chapter §1 in *loc. cit.* The only difference lies in section § 6, where we have considered a more general setting in the present text.

Remark Even if the theorems are stated in the category $\mathbf{Shv}_{\mathbb{Q}_\ell}(Z)$ of ℓ -adic sheaves, they are nevertheless true (and the proofs we provide work mutatis mutandis) as $\mathrm{BU}_{Z, \mathbb{Q}}$ -modules (see Sect. 2.7).

2 Preliminaries and notation

In this preliminary section we will briefly recall some of the mathematical tools that we will need later. We will also fix the notation that we will use in the other sections.

2.1 Some notation and convention

- Even when not explicitly stated, S will always be a regular scheme of finite type over a strictly local noetherian scheme.
- \mathbf{Sm}_S denotes the category of smooth schemes of finite type over S .
- \mathbf{Sch}_S denotes the category of separated schemes of finite type over S .
- We will freely use the language of ∞ -categories which has been developed in [34, 35]. ∞ -category will always mean $(\infty, 1)$ -category for us.
- \mathcal{S} denotes the ∞ -category of spaces.
- We write dg instead of “differential graded”.
- We will use cohomological notations. In particular, the differential of a complex increases the degree.
- If we are given a morphism of (derived) schemes $f : X \rightarrow Y$ and an object $\mathcal{E}_Y \in \mathbf{QCoh}(Y)$, we will write \mathcal{E}_X instead of $f^*\mathcal{E}_Y$.

2.2 Reminders on dg categories

Remark 2.2.1 For more details on the theory of dg categories, we invite the reader to consult [32, 60] and/or [48].

Let A be a commutative ring.

Let dgCat_S be the category of small A -linear dg categories together with A -linear dg functors. This category can be endowed with a cofibrantly generated model category structure, where weak equivalences are DK equivalences (see [56]). The underlying ∞ -category of this model category coincides with the ∞ -localization of dgCat_S with respect to the class of DK equivalences. We will denote this ∞ -category by \mathbf{dgCat}_S .

Every DK equivalence is a Morita equivalence. We can therefore endow dgCat_S with a second cofibrantly generated model category structure by using the theory of Bousfield localizations. In this case weak equivalences are Morita equivalences. Similarly to the previous case, the underlying ∞ -category of this model category coincides with the ∞ -localization of dgCat_S with respect to Morita equivalences. We will label this ∞ -category by $\mathbf{dgCat}_S^{\mathrm{idm}}$.

Let $\hat{\mathcal{C}}_c$ denote the dg category of perfect $\mathcal{C}^{\mathrm{op}}$ -dg modules. Then $\mathbf{dgCat}_S^{\mathrm{idm}}$ is equivalent to the full subcategory of \mathbf{dgCat}_S spanned by dg categories \mathcal{C} for which the Yoneda embedding $\mathcal{C} \hookrightarrow \hat{\mathcal{C}}_c$ is a DK equivalence.

To summarize, we have the following pair of composable ∞ -localizations

$$\mathbf{dgCat}_S \rightarrow \mathbf{dgCat}_S \rightarrow \mathbf{dgCat}_S^{\mathrm{idm}}. \quad (2.2.1)$$

$\mathbf{dgCat}_S \rightarrow \mathbf{dgCat}_S^{\mathrm{idm}}$ is a left adjoint to the inclusion $\mathbf{dgCat}_S^{\mathrm{idm}} \hookrightarrow \mathbf{dgCat}_S$ under the identification mentioned above. At the level of objects, it is defined by the assignment $T \mapsto \hat{T}_c$.

It is possible to enhance \mathbf{dgCat}_S and $\mathbf{dgCat}_S^{\mathrm{idm}}$ with symmetric monoidal structures. Furthermore, if we restrict to the full subcategory $\mathbf{dgCat}_S^{\mathrm{lf}} \subseteq \mathbf{dgCat}_S$ of locally flat (small) dg categories, we get two composable symmetric monoidal ∞ -functors

$$\mathbf{dgCat}_S^{\mathrm{lf}, \otimes} \rightarrow \mathbf{dgCat}_S^{\otimes} \rightarrow \mathbf{dgCat}_S^{\mathrm{idm}, \otimes}. \quad (2.2.2)$$

More details on the Morita theory of dg categories can be found in [59].

One of the most recurrent operations that occur in this work is that of forming quotients of dg categories: given a dg category \mathcal{C} together with a full sub dg category \mathcal{C}' , both of them in $\mathbf{dgCat}_S^{\mathrm{idm}}$, we can consider the pushout $\mathcal{C} \amalg_{\mathcal{C}'} 0$ in $\mathbf{dgCat}_S^{\mathrm{idm}}$. Here 0 denotes the final object in $\mathbf{dgCat}_S^{\mathrm{idm}}$, i.e. the dg category with only one object whose endomorphisms are given by the zero hom-complex. We denote this pushout by \mathcal{C}/\mathcal{C}' and refer to it as the dg quotient of $\mathcal{C}' \hookrightarrow \mathcal{C}$. Equivalently, The dg category \mathcal{C}/\mathcal{C}' can also be obtained as the image in $\mathbf{dgCat}_S^{\mathrm{idm}}$ of the pushout $\mathcal{C} \amalg_{\mathcal{C}'} 0$ formed in \mathbf{dgCat}_S . Its homotopy category coincides with (the idempotent completion of) the Verdier quotient of $H^0(\mathcal{C})$ by the full subcategory $H^0(\mathcal{C}')$ (see [22]).

Compact objects in $\mathbf{dgCat}_S^{\mathrm{idm}}$ are dg categories of finite type over A , as defined in [64]. In particular,

$$\mathrm{Ind}(\mathbf{dgCat}_S^{\mathrm{ft}}) \simeq \mathbf{dgCat}_S^{\mathrm{idm}}. \quad (2.2.3)$$

Moreover, $\mathbf{dgCat}_S^{\mathrm{idm}}$ is equivalent to the ∞ -category of small, idempotent complete, A -linear stable ∞ -categories [19].

When S is a non affine scheme, the symmetric monoidal ∞ -category of dg-categories over S is defined as the limit

$$\mathbf{dgCat}_S^{\mathrm{idm}, \otimes} = \varprojlim_{\mathrm{Spec}(A) \rightarrow S} \mathbf{dgCat}_A^{\mathrm{idm}, \otimes}. \quad (2.2.4)$$

Here there are some dg categories that we will use: let X be a derived scheme (stack).

- $\mathbf{QCoh}(X)$ will denote the dg category of quasi-coherent \mathcal{O}_X -modules. It can be defined as follows. If $X = \mathrm{Spec}(B)$, then $\mathbf{QCoh}(X) = \mathbf{Mod}_B$, the dg category of dg modules over the dg algebra associated to B via the Dold-Kan equivalence. In the general case, $\mathbf{QCoh}(X) = \varprojlim_{\mathrm{Spec}(B) \rightarrow X} \mathbf{Mod}_B$ (the functoriality being that induced by base change).
- $\mathbf{Perf}(X)$ will denote the full sub dg category of $\mathbf{QCoh}(X)$ spanned by perfect complexes. If $X = \mathrm{Spec}(B)$, then $\mathbf{Perf}(B)$ is the smallest subcategory of \mathbf{Mod}_B which

contains B and that is stable under the formation of finite colimits and retracts. More generally, an object $\mathcal{E} \in \mathbf{QCoh}(X)$ is perfect if, for any $g : \mathrm{Spec}(B) \rightarrow X$, the pullback $g^*\mathcal{E} \in \mathbf{Perf}(B)$. Perfect complexes coincide with dualizable objects in $\mathbf{QCoh}(X)$ and, under some mild additional assumptions (that will always be verified in our examples), with compact objects (see [12]).

Moreover, as we assume to work in the noetherian setting, we can consider:

- $\mathbf{Coh}^b(X)$ will denote the full sub dg category of $\mathbf{QCoh}(X)$ spanned by those cohomologically bounded complexes \mathcal{E} (i.e. $H^i(\mathcal{E}) \neq 0$ only for a finite number of indexes) such that $H^*(\mathcal{E})$ is a coherent $H^0(\mathcal{O}_X)$ -module.
- $\mathbf{Coh}^-(X)$ will denote the full dg category of $\mathbf{QCoh}(X)$ spanned by those cohomologically bounded above complexes \mathcal{E} (i.e. $H^i(\mathcal{E}) = 0$ for $i \gg 0$) such that $H^*(\mathcal{E})$ is a coherent $H^0(\mathcal{O}_X)$ -modules. These are also known as pseudo-coherent complexes.
- Let $p : X \rightarrow Y$ be a proper morphism locally almost of finite type. By [28, Cpt.4 Lemma 5.1.4], we have an induced ∞ -functor $p_* : \mathbf{Coh}^b(X) \rightarrow \mathbf{Coh}^b(Y)$. We denote $\mathbf{Coh}^b(X)_{\mathbf{Perf}(Y)}$ the full subcategory of $\mathbf{Coh}^b(X)$ spanned by those objects \mathcal{E} such that $p_*\mathcal{E} \in \mathbf{Perf}(Y)$.

2.3 Derived algebraic geometry

Derived algebraic geometry is a broad generalization of algebraic geometry whose building blocks are simplicial commutative rings, rather than commutative rings. It is usually better behaved in the situations that are typically defined bad in the classical context, e.g. non-transversal intersections. The main idea is to develop algebraic geometry in an homotopical context: instead of saying that two elements are equal we rather say that they are homotopic, the homotopy being part of the data. In this article derived schemes appear exclusively as (homotopy) fiber products of ordinary schemes. It is necessary to allow certain schemes to be derived, as if we restrict ourselves to work with discrete (i.e. classical) schemes, some important characters do not appear (e.g. the algebra which acts on certain dg categories of (relative) singularities).

The ideas and motivations that led to derived algebraic geometry go back to J.-P. Serre (Serre's intersection formula, [52]), P. Deligne (algebraic geometry in a symmetric monoidal category, [21]), Illusie, Grothendieck, André, Quillen (the cotangent complex, [3,30,46,53]) ... however, the theory nowadays relies on solid roots thanks to the work of Lurie [36] and Toën–Vezzosi [65,66]. For a brief introduction to derived algebraic geometry we refer to [63]. We will denote the ∞ -category of derived schemes by \mathbf{dSch} .

By definition, each derived scheme X has an underlying scheme $t_0(X)$ (its truncation). Indeed, the assignment $X \mapsto t_0(X)$ is part of an adjunction

$$\iota : \mathbf{Sch} \rightleftarrows \mathbf{dSch} : t_0. \quad (2.3.1)$$

A derived scheme is affine if its underlying scheme is so. There is an equivalence of ∞ -categories

$$\mathbf{dSch}^{\text{aff}} \simeq \mathbf{sRing}^{\text{op}} \quad (2.3.2)$$

where $\mathbf{dSch}^{\text{aff}}$ is the full subcategory of \mathbf{dSch} spanned by affine derived schemes and \mathbf{sRing} is the ∞ -category of simplicial rings. For any simplicial commutative ring A , we denote $\text{Spec}(A)$ the associated derived scheme.

The ∞ -category \mathbf{dSch} has all finite limits. In particular, it has fiber products. For example, if we consider a diagram

$$\text{Spec}(B) \rightarrow \text{Spec}(A) \leftarrow \text{Spec}(C)$$

of affine derived schemes, the fiber product is equivalent to $\text{Spec}(B \otimes_A^{\mathbb{L}} C)$, the spectrum of the derived tensor product.

Consider two (underived) S -schemes X, Y (S being underived itself). Then the fiber product computed in \mathbf{dSch} (denoted $X \times_S^h Y$) might differ from the one computed in \mathbf{Sch} (denoted $X \times_S Y$). However, they are related by the formula

$$t_0(X \times_S^h Y) \simeq X \times_S Y. \quad (2.3.3)$$

2.4 ℓ -adic sheaves

We shall briefly introduce the ∞ -category of ℓ -adic sheaves, following [27].

Fix a prime number ℓ , which is invertible in each residue field of our base scheme S , where S is a regular scheme of finite type over a strictly local noetherian scheme.² Let $\mathbf{Shv}(X, \mathbb{Z}/\ell^d \mathbb{Z})$ be the full subcategory of the ∞ -category $\text{Fun}(\mathbf{Sch}_{\text{et}}^{\text{op}}, \mathbf{Mod}_{\mathbb{Z}/\ell^d \mathbb{Z}})$ spanned by (hypercomplete) étale sheaves. Here \mathbf{Sch}_{et} denotes the category of étale sheaves and $\mathbf{Mod}_{\mathbb{Z}/\ell^d \mathbb{Z}}$ the ∞ -category of $\mathbb{Z}/\ell^d \mathbb{Z}$ -modules. The ∞ -category $\mathbf{Shv}(X, \mathbb{Z}/\ell^d \mathbb{Z})$ is compactly-generated, its compact objects being constructible sheaves (see [9, Proposition 3.38]). In what follows, we will denote $\mathbf{Shv}^c(X, \mathbb{Z}/\ell^d \mathbb{Z})$ the full subcategory of $\mathbf{Shv}(X, \mathbb{Z}/\ell^d \mathbb{Z})$ spanned by compact objects.

The ring homomorphisms $\mathbb{Z}/\ell^d \mathbb{Z} \rightarrow \mathbb{Z}/\ell^{d-1} \mathbb{Z}$ induce a sequence of ∞ -functors

$$\mathbf{Shv}(X, \mathbb{Z}/\ell^d \mathbb{Z}) \rightarrow \mathbf{Shv}(X, \mathbb{Z}/\ell^{d-1} \mathbb{Z}) \quad (2.4.1)$$

and it follows from [27, Proposition 2.2.8.4] that the image of a constructible sheaf is again constructible, yielding

$$\mathbf{Shv}^c(X, \mathbb{Z}/\ell^d \mathbb{Z}) \rightarrow \mathbf{Shv}^c(X, \mathbb{Z}/\ell^{d-1} \mathbb{Z}). \quad (2.4.2)$$

We can then consider the limit of the diagram of ∞ -categories

² This assumption is needed in order to consider schemes of finite étale cohomological dimension, see [14, Theorem 1.1.5].

$$\mathbf{Shv}_\ell^c(X) := \varprojlim \mathbf{Shv}^c(X, \mathbb{Z}/\ell^d \mathbb{Z}). \quad (2.4.3)$$

This ∞ -category can be identified with the full subcategory of $\mathbf{Shv}(X, \mathbb{Z})$ generated by ℓ -complete constructible sheaves, i.e. by those objects $\mathcal{F} \in \mathbf{Shv}(X, \mathbb{Z})$ such that

1. $\mathcal{F} \simeq \varprojlim \mathcal{F}/\ell^d \mathcal{F}$,
2. for any $d \geq 1$, $\mathcal{F}/\ell^d \mathcal{F}$ is constructible.

We will refer to this ∞ -category with the ∞ -category of constructible ℓ -adic sheaves.

The pushforward for a morphism $f : X \rightarrow Y$ of S -schemes of finite type induces an ∞ -functors at the level of constructible ℓ -adic sheaves

$$f_* : \mathbf{Shv}_\ell^c(X) \rightarrow \mathbf{Shv}_\ell^c(Y). \quad (2.4.4)$$

It admits a left adjoint

$$f^* : \mathbf{Shv}_\ell^c(Y) \rightarrow \mathbf{Shv}_\ell^c(X) \quad (2.4.5)$$

that, at the level of objects, takes a constructible ℓ -adic sheaf to the ℓ -completion of its pullback.

We next consider the ind-completion of such categories:

$$\mathbf{Shv}_\ell(X) := \mathrm{Ind}(\mathbf{Shv}_\ell^c(X)), \quad (2.4.6)$$

the ∞ -category of ℓ -adic sheaves. It is then a formal fact that we have a couple of adjoint functors, also called the pushforward and the pullback, defined at the level of ℓ -adic sheaves.

Finally, we consider the localization of $\mathbf{Shv}_\ell(X)$ with respect to the class of morphisms $\{\mathcal{F} \rightarrow \mathcal{F}[\ell^{-1}]\}$, obtaining the ∞ -category of \mathbb{Q}_ℓ -adic sheaves $\mathbf{Shv}_{\mathbb{Q}_\ell}(X)$.

2.5 Stable homotopy categories

In this section we will briefly recall the constructions and main properties of \mathbf{SH}_S and of $\mathbf{SH}_S^{\mathrm{nc}}$, the stable ∞ -category of schemes and the stable ∞ -category of non-commutative spaces (a.k.a. dg categories).

2.5.1 The stable homotopy category of schemes

The stable homotopy category of schemes was first introduced by Morel and Voevodsky in their celebrated paper [40]. The main idea is to develop an homotopy theory for schemes, where the role of the unit interval—which is not available in the world of schemes—is played by the affine line. It was first developed using the language of model categories. We will rather use that of ∞ -categories, following [48] and [49]. The two procedures are compatible, as shown in [48] and [49].

Let $S = \mathrm{Spec}(A)$ denote an affine scheme. One can produce the unstable homotopy category of schemes as follows: one considers the ∞ -category $\mathrm{Fun}(\mathbf{Sm}_S^{\mathrm{op}}, \mathbb{S})$ of

presheaves (of spaces) on \mathbf{Sm}_S . Its full subcategory spanned by Nisnevich sheaves, $\mathrm{Sh}_{\mathrm{Nis}}(\mathbf{Sm}_S)$ is an example of an ∞ -topos (in the sense of [35]). Then one has to consider its hypercompletion $\mathrm{Sh}_{\mathrm{Nis}}(\mathbf{Sm}_S)^{\mathrm{hyp}}$, which coincides with the localization of $\mathrm{Fun}(\mathbf{Sm}_S^{\mathrm{op}}, \mathcal{S})$ spanned by objects that are local with respect to Nisnevich hypercovers. If we further localize with respect to the projections $\{\mathbb{A}_X^1 \rightarrow X\}$, we obtain the unstable homotopy ∞ -category of schemes, which we will denote by \mathbf{H}_S . It is also important that the canonical ∞ -functor $\mathbf{Sm}_S \rightarrow \mathbf{H}_S$ can be promoted to a symmetric monoidal ∞ -functor with respect to the Cartesian structures. One then considers the pointed version of \mathbf{H}_S , \mathbf{H}_{S*} : it comes equipped with a canonical symmetric monoidal structure \mathbf{H}_{S*}^\wedge and there is a symmetric monoidal ∞ -functor $\mathbf{H}_S^\times \rightarrow \mathbf{H}_{S*}^\wedge$. The final step consists in stabilization: in classical stable homotopy theory one forces stabilization by inverting S^1 . However, in this context, there exist two circles, the topological circle $S^1 := \Delta^1/\partial\Delta^1$ and the algebraic circle $\mathbb{G}_{m,S}$. One then stabilizes \mathbf{H}_{S*} by inverting $S^1 \wedge \mathbb{G}_{m,S} \simeq (\mathbb{P}_S^1, \infty) := \mathrm{cofib}(S \xrightarrow{\infty} \mathbb{P}_S^1)$ ³—this can be done using the machinery developed in [49, §2.1]. As a result, we obtain the presentable, symmetric monoidal, stable ∞ -category $\mathbf{SH}_S^\otimes := \mathbf{H}_{S*}^\wedge[(\mathbb{P}_S^1, \infty)^{-1}]$, called the stable homotopy ∞ -category of schemes. It is moreover characterized by the following universal property (see [49, Corollary 2.39]): there is a symmetric monoidal ∞ -functor $\Sigma_+^\infty : \mathbf{Sm}_S^\times \rightarrow \mathbf{SH}_S^\otimes$ and for any presentable, symmetric monoidal pointed ∞ -category \mathcal{D}^\otimes , the map

$$\mathrm{Fun}^{\otimes, L}(\mathbf{SH}_S^\otimes, \mathcal{D}^\otimes) \rightarrow \mathrm{Fun}^\otimes(\mathbf{Sm}_S^\times, \mathcal{D}^\otimes) \quad (2.5.1)$$

induced by Σ_+^∞ is fully faithful and its image coincides with those symmetric monoidal ∞ -functors $F : \mathbf{Sm}_S^\times \rightarrow \mathcal{D}^\otimes$ that satisfy

- Nisnevich descent,
- \mathbb{A}^1 -invariance,
- $\mathrm{cofib}(F(S \xrightarrow{\infty} \mathbb{P}_S^1))$ is an invertible object in \mathcal{D} .

Remark 2.5.2 It can be shown, using results of Ayoub [4, 5], Cisinski–Déglise [15] and the machinery developed by Gaitsgory–Rozenblyum [28] and Liu–Zheng [37–39], that the assignment $S \mapsto \mathbf{SH}_S$ defines a sheaf of ∞ -categories enhanced with a Grothendieck 6-functors formalism. See [48] and [49] and the appendix in [9].

Among the objects of \mathbf{SH}_S , the spectrum of homotopy invariant non-connective K-theory BU_S ⁴ will play a crucial role in what follows. Recall that it is an object in \mathbf{SH}_S such that, for every object $X \in \mathbf{Sm}_S$,

$$\mathrm{Map}_{\mathbf{SH}_S}(\Sigma_+^\infty X, \mathrm{BU}_S) \simeq \mathrm{HK}(X). \quad (2.5.2)$$

Moreover, BU_S satisfies the algebraic Bott periodicity:

$$\mathrm{BU}_S \simeq \mathrm{BU}_S(1)[2]. \quad (2.5.3)$$

³ Clearly, this cofiber is taken in \mathbf{H}_{S*} .

⁴ More commonly, this spectrum is denoted KGL_S , see [16]. We use the notation BU_S , which comes from topology, following the lead of [9], which is our main source of inspiration.

2.5.2 The stable homotopy category of non-commutative spaces

Remark 2.5.4 This section is an exposition of the ideas and results of [48] and [49]. However, a theory of non-commutative motives was also proposed by Cisinski and Tabuada in [17, 18, 57]. Their theory is dual to the one constructed by Robalo. For more details about this, we refer to [49, Appendix A] and [9, Remark 3.4].

It is possible to mimic the procedure described in the previous paragraph to construct a presentable, symmetric monoidal, stable ∞ -category $\mathbf{SH}_S^{\text{nc}}$ defined starting with non-commutative spaces rather than schemes, a.k.a. dg categories. The role of smooth schemes is played by dg categories of finite type in this context. Moreover, there exists an analog of Nisnevich square of non-commutative spaces (see [48] and [49]).

Remark 2.5.5 By convention, \emptyset is a Nisnevich covering of the zero object in $\text{dgCat}_S^{\text{ft}}$.

Remark 2.5.6 The notion on Nisnevich squares for smooth non-commutative spaces is compatible with the classical one: if

$$\begin{array}{ccc} V \times_X U & \longrightarrow & V \\ \downarrow & & \downarrow \\ U & \longrightarrow & X \end{array} \quad (2.5.4)$$

is an elementary Nisnevich square, then its image via $\mathbf{Perf}(\bullet)$ is a Nisnevich square of non-commutative spaces ([49, Proposition 3.21]).

One then considers the category of presheaves $\text{Fun}(\mathcal{N}c_S^{\text{op}}, \mathcal{S})$ and its localization with respect to the class of morphisms $\{j(U) \amalg_{j(X)} j(V) \rightarrow j(W)\}$ determined by Nisnevich squares of non-commutative spaces, where $j : \mathcal{N}c_S \rightarrow \text{Fun}(\mathcal{N}c_S^{\text{op}}, \mathcal{S})$ is the Yoneda embedding. We further localize with respect to the morphisms $\{X \otimes \mathbf{Perf}(S) \rightarrow X \otimes \mathbf{Perf}(\mathbb{A}_S^1)\}$ ⁵ and obtain a presentable symmetric monoidal ∞ -category $\mathbf{H}_S^{\text{nc}, \otimes}$. Pursuing the analogy with the commutative case, we should now force the existence of a zero object in \mathbf{H}_S^{nc} and then stabilize with respect to the topological and algebraic circles. It comes out that the situation is simpler in the non-commutative case: let

$$\psi^{\otimes} : \mathcal{N}c_S^{\otimes} \rightarrow \mathbf{H}_S^{\text{nc}, \otimes} \rightarrow \mathbf{SH}_S^{\text{nc}, \otimes} := \mathbf{H}_S^{\text{nc}, \otimes}[(S^1)^{-1}]$$

denote the ∞ -functor that we obtain if we force the topological circle to be invertible. Then

- $\mathbf{SH}_S^{\text{nc}}$ is pointed due to the convention of Remark 2.5.5.
- The non-commutative motive

$$\psi(\mathbb{P}_S^1, \infty) = \text{cofib}(\psi(\mathbf{Perf}(S) \xrightarrow{\infty^*} \mathbf{Perf}(\mathbb{P}_S^1)))$$

⁵ Since the Yoneda embedding is symmetric monoidal (when we consider the Day convolution product on the ∞ -category of presheaves), then it suffices to localize with respect to the morphism $j(\mathbf{Perf}(S)) \rightarrow j(\mathbf{Perf}(\mathbb{A}_S^1))$.

is invertible in $\mathbf{SH}_S^{\text{nc}, \otimes}$ ([49, Proposition 3.24]): indeed, it is equivalent to $\mathbf{Perf}(S) \simeq 1^{\text{nc}}$ and therefore it is not necessary to force $\mathbb{G}_{m,S}$ to be invertible.

2.6 The bridge between motives and non-commutative motives

We have now at our disposal the following picture:

$$\begin{array}{ccc}
 \mathbf{Sm}_S^\times & \xrightarrow{\mathbf{Perf}(\bullet)} & \mathcal{N}c_S^\otimes \\
 \downarrow \Sigma_+^\infty & & \downarrow \psi^\otimes \\
 \mathbf{SH}_S^\otimes & \xrightarrow[\mathcal{R}_{\mathbf{Perf}}]{\text{dotted}} & \mathbf{SH}_S^{\text{nc}, \otimes}
 \end{array} \tag{2.6.1}$$

The existence of the dotted map, which we name perfect realization, is granted by the universal property of \mathbf{SH}_S^\otimes . Indeed

- $\psi \circ \mathbf{Perf}(\bullet)$ sends (ordinary) Nisnevich squares to pushout squares in $\mathbf{SH}_S^{\text{nc}}$: this is a consequence of the compatibility of non-commutative Nisnevich squares with the classical ones and of the definition of $\mathbf{SH}_S^{\text{nc}}$,
- \mathbb{A}^1 -invariance is forced by construction,
- $\psi(\mathbb{P}_S^1, \infty)$ is an invertible object in $\mathbf{SH}_S^{\text{nc}, \otimes}$.

The fact that $\mathcal{R}_{\mathbf{Perf}}$ commutes with colimits is also guaranteed by the universal property of \mathbf{SH}_S^\otimes . As both \mathbf{SH}_S and $\mathbf{SH}_S^{\text{nc}}$ are presentable ∞ -categories, the adjoint functor theorem [35, Corollary 5.5.2.9] implies the existence of a lax-monoidal right adjoint

$$\mathcal{M}^\otimes : \mathbf{SH}_S^{\text{nc}, \otimes} \rightarrow \mathbf{SH}_S^\otimes. \tag{2.6.2}$$

For our purposes, it will be very important the following result

Theorem 2.6.1 [49, Corollary 4.10] *The image of the monoidal unit 1_S^{nc} via \mathcal{M} is equivalent to \mathbf{BU}_S .*

In particular, the ∞ -functor \mathcal{M} factors through the full subcategory of \mathbf{SH}_S spanned by modules over \mathbf{BU}_S :

$$\mathcal{M}^\otimes : \mathbf{SH}_S^{\text{nc}, \otimes} \rightarrow \mathbf{Mod}_{\mathbf{BU}_S}(\mathbf{SH}_S)^\otimes. \tag{2.6.3}$$

In [9], the authors introduced a dual version of this ∞ -functor. Consider the endomorphism of $\mathbf{SH}_S^{\text{nc}}$ induced by the internal hom:

$$\mathbb{R}\text{Hom}_{\mathbf{SH}_S^{\text{nc}}}(-, 1_S^{\text{nc}}) : \mathbf{SH}_S^{\text{nc}, \text{op}, \otimes} \rightarrow \mathbf{SH}_S^{\text{nc}, \otimes}. \tag{2.6.4}$$

Then consider

$$\begin{array}{ccccc}
 \mathbf{dgCat}_S^{\mathrm{ft}, \otimes} & \xrightarrow{\psi^\otimes} & \mathbf{SH}_S^{\mathrm{nc}, \mathrm{op}, \otimes} & \xrightarrow{\mathbb{R}\mathrm{Hom}_{\mathbf{SH}_S^{\mathrm{nc}}}(-, 1_S^{\mathrm{nc}})} & \mathbf{SH}_S^{\mathrm{nc}, \otimes} \xrightarrow{\mathcal{M}^\otimes} \mathbf{Mod}_{\mathrm{BU}_S}(\mathbf{SH}_S^\otimes) \\
 \downarrow & \nearrow \psi^\otimes & & & \uparrow \\
 \mathbf{dgCat}_S^{\mathrm{idm}, \otimes} & \xrightarrow{\quad \quad \quad \mathcal{M}^{\vee, \otimes} \quad \quad \quad} & & &
 \end{array} \quad (2.6.5)$$

where the vertical map on the left is given by the inclusion of $\mathbf{dgCat}_S^{\mathrm{ft}, \otimes}$ in its Ind-completion $\mathbf{dgCat}_S^{\mathrm{idm}, \otimes}$ and the oblique ψ^\otimes is induced by the universal property of the Ind-completion. Let $T \in \mathbf{dgCat}_S^{\mathrm{idm}}$. Then $\mathcal{M}^\vee(T)$ is the sheaf of spectra $X \in \mathbf{Sm}_S \mapsto \mathrm{HK}(\mathbf{Perf}(X) \otimes_S T)$:

$$\begin{aligned}
 \mathcal{M}^\vee(T)(X) &= \mathrm{Map}_{\mathbf{SH}_S}(\Sigma_+^\infty X, \mathcal{M}^\vee(T)) \\
 &\simeq \mathrm{Map}_{\mathbf{SH}_S^{\mathrm{nc}}}(\mathcal{R}\mathbf{Perf}(\Sigma_+^\infty X), \mathbb{R}\mathrm{Hom}_{\mathbf{SH}_S^{\mathrm{nc}}}(T, 1_S^{\mathrm{nc}})) \\
 &\simeq \mathrm{Map}_{\mathbf{SH}_S^{\mathrm{nc}}}(\mathcal{R}\mathbf{Perf}(\Sigma_+^\infty X) \otimes_S T, 1_S^{\mathrm{nc}}) \simeq \mathrm{HK}(\mathbf{Perf}(X) \otimes_S T). \quad (2.6.6)
 \end{aligned}$$

Moreover, \mathcal{M}^\vee has the following nice properties

- it is lax monoidal (it is a composition of lax monoidal ∞ -functors),
- it commutes with filtered colimits (see [9, Remark 3.4]),
- it sends exact sequences of dg categories to fiber-cofiber sequences in $\mathbf{Mod}_{\mathrm{BU}_S}(\mathbf{SH}_S)^\otimes$ (see [9, Corollary 3.3]).

We will refer to \mathcal{M}^\vee as the motivic realization of dg categories.

2.7 ℓ -adic realization of dg categories

We will need a way to associate an ℓ -adic sheaf to a dg category. We follow the construction given in [9, §3.6, §3.7], which relies on results of J. Ayoub and D.-C. Cisinski - F. Déglise. Let $\mathrm{H}\mathbb{Q}$ be the Eilenberg-MacLane spectrum of rational homotopy theory. Then we get

$$- \otimes \mathrm{H}\mathbb{Q} : \mathbf{SH}_S^\otimes \rightarrow \mathbf{Mod}_{\mathrm{H}\mathbb{Q}}(\mathbf{SH}_S)^\otimes, \quad (2.7.1)$$

which identifies the ∞ -category on the right hand side with non-torsion objects of \mathbf{SH}_S . Similarly, if one puts $\mathrm{BU}_{S, \mathbb{Q}} := \mathrm{H}\mathbb{Q} \otimes \mathrm{BU}_S$, then one gets

$$- \otimes \mathrm{H}\mathbb{Q} : \mathbf{Mod}_{\mathrm{BU}_S}(\mathbf{SH}_S)^\otimes \rightarrow \mathbf{Mod}_{\mathrm{BU}_{S, \mathbb{Q}}}(\mathbf{SH}_S)^\otimes, \quad (2.7.2)$$

where the right hand side identifies with non-torsion BU_S -modules. This ∞ -functor is strongly compatible with the 6-functors formalism and Tate twists. Moreover, we state for future reference the following crucial fact:

Theorem 2.7.1 [47, Theorem 5.3.10], [15, §14.1], [9, Proposition 3.35] *Let X be a scheme of finite Krull dimension and let $\mathbb{M}_X \in \mathbf{CAlg}(\mathbf{SH}_X)$ denote the spectrum of Beilinson motivic cohomology. Then the canonical morphism $1_X(1)[2] \rightarrow \mathbb{B}\mathbb{U}_X \otimes \mathbb{H}\mathbb{Q} = \mathbb{B}\mathbb{U}_{X,\mathbb{Q}}$ induces an equivalence of commutative algebra objects*

$$\mathbb{M}_X(\beta) := \mathrm{Sym}(\mathbb{M}_X(1)[2])[v^{-1}] \simeq \mathbb{B}\mathbb{U}_{X,\mathbb{Q}} = \mathbb{B}\mathbb{U}_X \otimes \mathbb{H}\mathbb{Q}, \quad (2.7.3)$$

where v is the free generator in degree $(1)[2]$.

By using the theory of h-motives developed by the authors in [14] one can define an ℓ -adic realization ∞ -functor

$$\mathcal{R}^\ell : \mathbf{Mod}_{\mathbb{M}}(\mathbf{SH}) \rightarrow \mathbf{Shv}_{\mathbb{Q}_\ell}(-) \quad (2.7.4)$$

strongly compatible with the 6-functors formalism and Tate twists, at least for noetherian schemes of finite Krull dimension. Then, using the equivalence

$$\mathbf{Mod}_{\mathbb{B}\mathbb{U}}(\mathbf{Mod}_{\mathbb{M}}(\mathbf{SH})) \simeq \mathbf{Mod}_{\mathbb{B}\mathbb{U}}(\mathbf{SH}) \quad (2.7.5)$$

we obtain an ℓ -adic realization ∞ -functor

$$\begin{aligned} \mathcal{R}^\ell : \mathbf{Mod}_{\mathbb{B}\mathbb{U}}(\mathbf{SH}) &\xrightarrow{- \otimes \mathbb{H}\mathbb{Q}} \mathbf{Mod}_{\mathbb{B}\mathbb{U}_{\mathbb{Q}}}(\mathbf{SH}) \longrightarrow \mathbf{Mod}_{\mathcal{R}^\ell(\mathbb{B}\mathbb{U})}(\mathbf{Shv}_{\mathbb{Q}_\ell}(-)) \\ &\simeq \mathbf{Mod}_{\mathbb{Q}_\ell(\beta)}(\mathbf{Shv}_{\mathbb{Q}_\ell}(-)) \end{aligned} \quad (2.7.6)$$

strongly compatible with the 6-functors formalism and Tate twists. The last equivalence above holds as (2.7.4) is symmetric monoidal and commutes with Tate twists (see [9, Remark 3.43]):

$$\begin{aligned} \mathcal{R}_X^\ell(\mathbb{B}\mathbb{U}_X) &\underset{\text{Theorem 2.7.1}}{\simeq} \mathcal{R}_X^\ell(\mathrm{Sym}(\mathbb{M}_X(1)[2])[v^{-1}]) \\ &\simeq \mathrm{Sym}(\mathcal{R}_X^\ell(\mathbb{M}_X(1)[2])[v^{-1}]) \simeq \mathbb{Q}_{\ell,X}(\beta) = \mathrm{Sym}(\mathbb{Q}_{\ell,X}(1)[2])[v^{-1}]. \end{aligned} \quad (2.7.7)$$

We will use the notation

$$\mathcal{R}_S^{\ell,\vee} := \mathcal{R}_S^\ell \circ \mathcal{M}_S^\vee : \mathbf{dgCat}_S^{\mathrm{idm},\otimes} \rightarrow \mathbf{Mod}_{\mathbb{Q}_{\ell,S}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_\ell}(S)) \quad (2.7.8)$$

and refer to this ∞ -functor as the ℓ -adic realization of dg categories.

3 Motivic realization of twisted LG models

Notation 3.0.1 Let S be a noetherian (not necessarily affine) regular scheme. We will label \mathbf{Sch}_S the category of separated S -schemes of finite type.

3.1 The category of twisted LG models

Consider the category \mathbf{Sch}_S and let \mathcal{L}_S be a line bundle on S . Then we can consider the category of Landau-Ginzburg models over (S, \mathcal{L}_S) , defined as follows:

- Objects consists of pairs (X, s_X) , where $p : X \rightarrow S$ is a (flat) S -scheme and s_X is a section of $\mathcal{L}_X := p^*\mathcal{L}_S$.
- Morphisms $(X, s_X) \xrightarrow{f} (Y, s_Y)$ consist of morphisms $f : X \rightarrow Y$ in \mathbf{Sch}_S such that $s_X = f^*s_Y$.
- Composition and identity morphisms are clear.

We will denote this category by $\mathbf{LG}_{(S, \mathcal{L}_S)}$.

Remark 3.1.1 If \mathcal{L}_S is the trivial line bundle, then $\mathbf{LG}_{(S, \mathcal{L}_S)}$ coincides with the category of usual Landau-Ginzburg models over S (as defined for example in [9]). Indeed, in this case $\mathbb{V}(\mathcal{L}_S) = \text{Spec}_{\mathcal{O}_S}(\mathcal{O}_S[t])$. Therefore, for any $X \in \mathbf{Sch}_S$, a section of \mathcal{L}_X consists of a morphism $\mathcal{O}_X[t] \rightarrow \mathcal{O}_X$, i.e. of a global section of \mathcal{O}_X .

Construction 3.1.2 It is possible to endow the category $\mathbf{LG}_{(S, \mathcal{L}_S)}$ with a symmetric monoidal structure

$$\begin{aligned} \boxplus : \mathbf{LG}_{(S, \mathcal{L}_S)} \times \mathbf{LG}_{(S, \mathcal{L}_S)} &\rightarrow \mathbf{LG}_{(S, \mathcal{L}_S)} \\ ((X, s_X), (Y, s_Y)) &\mapsto (X, s_X) \boxplus (Y, s_Y) = (X \times_S Y, s_X \boxplus s_Y), \end{aligned} \quad (3.1.1)$$

where $s_X \boxplus s_Y = p_X^*s_X + p_Y^*s_Y$. This is clearly (weakly) associative and the unit is given by $(S, 0)$, where 0 stands for the zero section of \mathcal{L}_S .

Notice that this tensor product is induced by the abelian group structure on $\mathbb{V}(\mathcal{L}_S) = \text{Spec}_S(\text{Sym}_{\mathcal{O}_S}(\mathcal{L}_S^\vee))$.

Remark 3.1.3 If \mathcal{L}_S is the trivial line bundle, then \boxplus coincides with the tensor product on the category of LG modules over S .

We will now exhibit twisted LG models as a fibered category.

Definition 3.1.4 Let LG be the category defined as follows:

- Objects are triplets $(f : Y \rightarrow X, \mathcal{L}_X, s_Y)$ where f is a flat morphism between S -schemes, \mathcal{L}_X is a line bundle on X and s_Y is a global section of $f^*\mathcal{L}_X$.
- Given two objects $(f_i : Y_i \rightarrow X_i, \mathcal{L}_{X_i}, s_{Y_i})$ ($i = 1, 2$), a morphism from the first to the second is the datum of a commutative diagram

$$\begin{array}{ccc} Y_1 & \xrightarrow{f_1} & X_1 \\ \downarrow g_Y & & \downarrow g_X \\ Y_2 & \xrightarrow{f_2} & X_2 \end{array} \quad (3.1.2)$$

and of an isomorphism $\alpha : g_X^* \mathcal{L}_{X_2} \rightarrow \mathcal{L}_{X_1}$ such that s_{Y_1} corresponds to s_{Y_2} under the isomorphism

$$g_Y^* f_2^* \mathcal{L}_{X_2} \simeq f_1^* g_X^* \mathcal{L}_{X_2} \underset{\alpha}{\simeq} f_1^* \mathcal{L}_{X_1}. \quad (3.1.3)$$

By abuse of notation, we will say in the future that $g_Y^*(s_{Y_2}) = s_{Y_1}$ if this condition is satisfied. We will denote such a morphism by (g, α) .

- Composition and identities are defined in an obvious way.

We will refer to this category as the category of twisted Landau-Ginzburg models of rank 1 over (S, \mathcal{L}_S) (twisted LG models for short).

Notice that there is a functor

$$\pi : \text{LG} \rightarrow \int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X) \quad (3.1.4)$$

defined as

$$\begin{array}{ccc} (f_1 : Y_1 \rightarrow X_1, \mathcal{L}_{X_1}, s_{Y_1}) & & (X_1, \mathcal{L}_{X_1}) \\ \downarrow (g, \alpha) & \mapsto & \downarrow (g_X, \alpha) \\ (f_2 : Y_2 \rightarrow X_2, \mathcal{L}_{X_2}, s_{Y_2}) & & (X_2, \mathcal{L}_{X_2}) \end{array}$$

Lemma 3.1.5 *The functor (3.1.4) exhibits LG as a fibered category over $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$.*

Proof Consider a map $(g_X, \alpha) : (X_1, \mathcal{L}_{X_1}) \rightarrow (X_2, \mathcal{L}_{X_2})$ in $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$ and let $(f_2 : Y_2 \rightarrow X_2, \mathcal{L}_{X_2}, s_{Y_2})$ be an object of LG over (X_2, \mathcal{L}_{X_2}) . Consider the morphism

$$(g, \alpha) : (f_1 : Y_1 := X_1 \times_{X_2} Y_2 \rightarrow X_1, \mathcal{L}_{X_1}, s_{Y_1}) \rightarrow (f_2 : Y_2 \rightarrow X_2, \mathcal{L}_{X_2}, s_{Y_2})$$

where g_Y is the projection $Y_1 \rightarrow Y_2$ (which is flat as it is the pullback of a flat morphism), f_1 is the projection $Y_1 \rightarrow Y_2$ and s_{Y_1} is $g_Y^*(s_{Y_2})$. It is clear that it is a morphism of LG over (g_X, α) . We need to show that it is cartesian. Consider

$$\begin{array}{ccc}
(h : Z \rightarrow W, \mathcal{L}_W, s_Z) & & \\
\downarrow & \searrow^{(q, \gamma)} & \\
& ((r, p), \beta) & \\
& \searrow & \\
& (f_1 : Y_1 \rightarrow X_1, \mathcal{L}_{X_1}, s_{Y_1}) & \xrightarrow{(g, \alpha)} (f_2 : Y_2 \rightarrow X_2, \mathcal{L}_{X_2}, s_{Y_2}) \\
& \downarrow & \downarrow \\
(W, \mathcal{L}_W) & \xrightarrow{(p, \beta)} (X_1, \mathcal{L}_{X_1}) & \xrightarrow{(g_X, \alpha)} (X_2, \mathcal{L}_{X_2})
\end{array}
\tag{3.1.5}$$

Then the universal property of Y_1 gives us an unique morphism $r : Z \rightarrow Y_1$ such that the compositions with f_1 and g_Y are $p \circ h$ and q respectively. We just need to show that $r^*(s_{Y_1}) = s_Z$. But this is clear since $s_{Y_1} = g_Y^*(s_{Y_2})$, $q_Y^*(s_{Y_2}) = s_Z$ and $g_Y \circ r = q_Y$. \square

Remark 3.1.6 Let (X, \mathcal{L}_X) be an object of $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$. Then the fiber of (X, \mathcal{L}_X) along π is $\text{LG}_{(X, \mathcal{L}_X)}$.

Definition 3.1.7 We say that a collection of maps $\{(g_i, \alpha_i) : (U_i, \mathcal{L}_{U_i}) \rightarrow (X, \mathcal{L}_X)\}_{i \in I}$ in $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$ is a Zariski covering if $\{g_i : U_i \rightarrow X\}_{i \in I}$ is so. They clearly define a pre-topology on $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$.⁶ We will refer to the corresponding topology by Zariski topology on $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$.

Lemma 3.1.8 LG is a stack over $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$ endowed with the Zariski topology.

Proof This is a simple consequence of the fact that a morphism of schemes is uniquely determined by its restriction to a Zariski covering and that line bundles are Zariski sheaves. \square

We conclude this section with the following observation:

Lemma 3.1.9 Let $\{(g_i, \alpha_i) : (U_i, \mathcal{L}_{U_i}) \rightarrow (X, \mathcal{L}_X)\}_{i \in I}$ be a Zariski covering in $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$. Then the canonical functor

$$\text{LG}_{(X, \mathcal{L}_X)}^{\boxplus} \rightarrow \varprojlim \text{LG}_{(U_i, \mathcal{L}_{U_i})}^{\boxplus} \tag{3.1.6}$$

is a symmetric monoidal equivalence.

Proof Consider the functors

$$\begin{aligned}
\text{LG}_{(X, \mathcal{L}_X)} &\rightarrow \text{LG}_{(U_i, \mathcal{L}_{U_i})} \\
(f : Y \rightarrow X, s_Y) &\mapsto (f_i : Y \times_X U_i \rightarrow U_i, s_{Y|U_i}).
\end{aligned}
\tag{3.1.7}$$

⁶ Notice that pullbacks exists in $\int_{X \in \text{Sch}_S} \text{B}\mathbb{G}_{m,S}(X)$: the pullback of $(Y_1, \mathcal{L}_{Y_1}) \xrightarrow{(f_1, \alpha_1)} (X, \mathcal{L}_X) \xleftarrow{(f_2, \alpha_2)} (Y_2, \mathcal{L}_{Y_2})$ is $(Y_1 \times_X Y_2, \mathcal{L}_{Y_1 \times_X Y_2})$ with the projections defined in an obvious way.

It is easy to see that they respect the tensor structure. Therefore, we get the desired symmetric monoidal functor (3.1.6) in the (big) category of symmetric monoidal categories and symmetric monoidal functors. Then, in order to prove that is a symmetric monoidal equivalence, it suffices to show that the underlying functor

$$Forget(\mathrm{LG}_{(X, \mathcal{L}_X)}^{\boxplus} \rightarrow \varprojlim \mathrm{LG}_{(U_i, \mathcal{L}_{U_i})}^{\boxplus}) \simeq (\mathrm{LG}_{(X, \mathcal{L}_X)} \rightarrow Forget(\varprojlim \mathrm{LG}_{(U_i, \mathcal{L}_{U_i})}^{\boxplus})). \quad (3.1.8)$$

As the functor which forgets the symmetric monoidal structure of a category is a right adjoint, it preserves limits. Thus,

$$Forget(\varprojlim \mathrm{LG}_{(U_i, \mathcal{L}_{U_i})}^{\boxplus}) \simeq \varprojlim \mathrm{LG}_{(U_i, \mathcal{L}_{U_i})}. \quad (3.1.9)$$

The assertion now follows from the previous lemma. \square

3.2 The dg category of singularities of a twisted LG model

Let (X, s_X) be a twisted LG model over (S, \mathcal{L}_S) . The section s_X defines a closed sub-scheme of X . Since we are not assuming that the section is regular, some derived structure might appear. More precisely: let $\mathcal{O}_X \rightarrow \mathcal{L}_X$ be the morphism of \mathcal{O}_X -modules associated to s_X . Then, taking duals, it defines a morphism $\mathcal{L}_X^\vee \rightarrow \mathcal{O}_X$. If we apply the relative spectrum functor, we get a morphism $\mathrm{Spec}_{\mathcal{O}_X}(\mathcal{O}_X) = X \rightarrow \mathrm{Spec}_{\mathcal{O}_X}(\mathrm{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee)) = \mathbb{V}(\mathcal{L}_X)$, i.e. a section of the vector bundle associated to \mathcal{L}_X . By abuse of notation, we will label this morphism by s_X . We can also consider the zero section $X \rightarrow \mathbb{V}(\mathcal{L}_X)$, which is the morphism associated to $0 \in \mathcal{L}_X(X)$.

Definition 3.2.1 The derived zero locus of s_X is defined as the derived pullback

$$\begin{array}{ccc} X_0 & \xrightarrow{i} & X \\ \downarrow & & \downarrow s_X \\ X & \xrightarrow{0} & \mathbb{V}(\mathcal{L}_X). \end{array} \quad (3.2.1)$$

Remark 3.2.2 Notice that the zero section $X \xrightarrow{0} \mathbb{V}(\mathcal{L}_X)$ is an lci morphism (Zariski locally, it is just the zero section a scheme Y in the affine line \mathbb{A}_Y^1). It is well known that this class of morphisms is stable under derived pullbacks. In particular, $i : X_0 \rightarrow X$ is a derived lci morphism.

Remark 3.2.3 There is a truncation morphism $t : \pi_0(X_0) \rightarrow X$, where $\pi_0(X_0)$ is the classical scheme associated to X_0 . Whenever s_X is a regular section, the truncation morphism is an equivalence in the ∞ -category of derived schemes.

Remark 3.2.4 Of major importance for our purposes is that i is an lci morphism. Indeed, by [61], if $f : Y \rightarrow Z$ is an lci morphism of derived schemes and $E \in \mathbf{Perf}(Y)$,

then $f_*E \in \mathbf{Perf}(Z)$. In particular, we get that $\mathbf{Perf}(X_0)$ is a full subcategory of $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$, the full subcategory of $\mathbf{Coh}^b(X_0)$ spanned by those objects $E \in \mathbf{Coh}^b(X_0)$ such that $f_*E \in \mathbf{Perf}(X)$.

The dg category of absolute singularities of a derived scheme is a non-commutative invariant (in the sense of Kontsevich) which captures the singularities of the scheme.

Definition 3.2.5 Let Z be a derived scheme of finite type over S whose structure sheaf is cohomologically bounded. Then the dg category of absolute singularities is the dg quotient in $\mathbf{dgCat}_S^{\mathrm{idm}}$

$$\mathbf{Sing}(Z) := \mathbf{Coh}^b(Z)/\mathbf{Perf}(Z). \quad (3.2.2)$$

Remark 3.2.6 It is a classical theorem due to Auslander–Buchsbaum ([1, Theorem 4.1]) and Serre ([50, Théorème 3]) that a noetherian ring A is regular if and only if it has finite global dimension. This means that the category of perfect complexes $\mathbf{Perf}(A)$ coincides with that of coherent bounded complexes $\mathbf{Coh}^b(A)$, i.e. $\mathbf{Sing}(A) \simeq 0$. This explains the name of this object.

Remark 3.2.7 Notice that the hypothesis on Z are crucial, as in general the structure sheaf of a derived scheme might not be cohomologically bounded. As an example of a derived scheme that doesn't sit in this class of objects, consider

$$\mathrm{Spec}(\mathbb{C}[x] \otimes_{\mathbb{C}[x,y]/(xy)}^{\mathbb{L}} \mathbb{C}[y]) \simeq \mathrm{Spec}(\mathbb{C}[x]) \times_{\mathrm{Spec}(\mathbb{C}[x,y]/(xy))}^h \mathrm{Spec}(\mathbb{C}[y]).$$

By tensoring the projective $\mathbb{C}[x, y]/(xy)$ -resolution of $\mathbb{C}[x]$

$$\dots \xrightarrow{y} \mathbb{C}[x, y]/(xy) \xrightarrow{x} \mathbb{C}[x, y]/(xy) \xrightarrow{y} \mathbb{C}[x, y]/(xy) \rightarrow 0$$

with $\mathbb{C}[y]$ over $\mathbb{C}[x, y]/(xy)$ we get that this derived scheme is the spectrum of the cdga

$$\dots \xrightarrow{y} \mathbb{C}[y] \xrightarrow{0} \mathbb{C}[y] \xrightarrow{y} \mathbb{C}[y] \rightarrow 0,$$

which has nontrivial cohomology in every even negative degree.

Since i is an lci morphism of derived schemes, by [61] and by [28] the pushforward induces a morphism

$$i_* : \mathbf{Sing}(X_0) \rightarrow \mathbf{Sing}(X). \quad (3.2.3)$$

Definition 3.2.8 Let (X, s_X) be a twisted LG model over (S, \mathcal{L}_S) . Its dg category of singularities is defined as the fiber in $\mathbf{dgCat}_S^{\mathrm{idm}}$ of (3.2.3):

$$\mathbf{Sing}(X, s_X) := \mathrm{fiber}(i_* : \mathbf{Sing}(X_0) \rightarrow \mathbf{Sing}(X)). \quad (3.2.4)$$

Remark 3.2.9 It is a consequence of the functoriality properties of the pullback and that of taking quotients and fibers in $\mathbf{dgCat}_S^{\text{idm}}$ that the assignments $(X, s_X) \mapsto \mathbf{Sing}(X, s_X)$ can be organized in an ∞ -functor

$$\mathbf{Sing}(\bullet, \bullet) : \mathbf{LG}_{(S, \mathcal{L}_S)}^{\text{op}} \rightarrow \mathbf{dgCat}_S^{\text{idm}}. \quad (3.2.5)$$

We will need to endow the ∞ -functor (3.2.5) with the structure of a lax monoidal ∞ -functor. In order to do so, we will introduce a strict model for $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$, following the strategy exploited in [9].

Construction 3.2.10 Let (X, s_X) be a twisted LG model over (S, \mathcal{L}_S) . The identity morphism on \mathcal{L}_X^\vee induces a $\text{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee)$ -projective resolution of \mathcal{O}_X , namely

$$\mathcal{L}_X^\vee \otimes_{\mathcal{O}_X} \text{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee) \rightarrow \text{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee). \quad (3.2.6)$$

This is a morphism of $\text{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee)$ -modules (induced by the multiplication of $\text{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee)$). Base changing along the morphism $\text{Sym}_{\mathcal{O}_X}(\mathcal{L}_X^\vee) \rightarrow \mathcal{O}_X$ induced by s_X we obtain that the cdga associated to X_0 is the spectrum of $\mathcal{L}_X^\vee \xrightarrow{s_X} \mathcal{O}_X$.

Example 3.2.11 Consider the zero section of \mathcal{L}_X . Then the cdga associated to the structure sheaf of $X \times_{\mathbb{V}(\mathcal{L}_X)}^h X$ is the Koszul algebra in degrees $[-1, 0]$ (recall that we use cohomological conventions)

$$\mathcal{L}_X^\vee \xrightarrow{0} \mathcal{O}_X. \quad (3.2.7)$$

Construction 3.2.12 Let $S = \text{Spec}(A)$ be a noetherian, regular affine scheme and let L be a projective A -module of rank 1. Let $\mathbf{LG}_{(S, \mathcal{L}_S)}^{\text{aff}}$ be the category of affine twisted LG models over (S, L) , i.e. of pairs $(\text{Spec}(B), s)$ where B is a flat A -algebra of finite type and $s \in L_B = L \otimes_A B$. For any object $(X, s) = (\text{Spec}(B), s)$, the derived zero locus of s is given by the spectrum of the Koszul algebra $K(B, s)$. The associated cdga is

$$K(B, s) = L_B^\vee \xrightarrow{s} B$$

concentrated in degrees -1 and 0 , where we label $s : L_B^\vee \rightarrow B$ the morphism induced by $s : B \rightarrow L_B$. Then, similarly to [9, Remark 2.30], there is an equivalence between cofibrant $K(B, s)$ -dg modules (denoted $\widehat{K}(B, s)$) and $\mathbf{QCoh}(X_0)$. Under this equivalence, $\mathbf{Coh}^b(X_0)$ corresponds to the full sub-dg category of $\widehat{K}(B, s)$ spanned by those cohomologically bounded dg modules with coherent cohomology (over $B/s = \text{coker}(L_B^\vee \xrightarrow{s} B)$) and $\mathbf{Perf}(X_0)$ corresponds to homotopically finitely presented $K(B, s)$ -dg modules. Moreover, the pushforward

$$i_* : \mathbf{QCoh}(X_0) \rightarrow \mathbf{QCoh}(X)$$

corresponds to the forgetful functor

$$\widehat{K(B, s)} \rightarrow \widehat{B}.$$

Define $\mathrm{Coh}^s(B, s)$ as the dg category of $K(B, s)$ -dg modules whose underlying B -dg module is perfect. Since X is an affine scheme, every perfect B -dg module is equivalent to a strictly perfect one. Therefore, we can take $\mathrm{Coh}^s(B, s)$ as the dg category of $K(B, s)$ -dg modules whose underlying B -module is strictly perfect. More explicitly, such an object corresponds to a triplet (E, d, h) where (E, d) is a strictly perfect B -dg module and

$$h : E \rightarrow E \otimes_B L_B[-1]$$

is a morphism such that

$$(h \otimes id_{L_B}[-1]) \circ h = 0 \quad [d, h] = h \circ d + (d \otimes id[-1]) \circ h = id \otimes s.$$

Notice that $\mathrm{Coh}^s(B, s)$ is a locally flat A -linear dg category. This follows immediately from the fact that the underlying complexes of B -modules are strictly perfect and from the fact that B is a flat A -algebra.

This strict dg category is analogous to the one that has been introduced in [9]. It gives us a strict model for the ∞ -category $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$. Indeed, the following result holds:

Lemma 3.2.13 [9, Lemma 2.33] *Let $\mathrm{Coh}^s(B, s)^{\mathrm{acy}}$ be the full sub-category of $\mathrm{Coh}^s(B, s)$ spanned by acyclic dg modules. Then the cofibrant replacement induces an equivalence of dg categories*

$$\mathrm{Coh}^s(B, s)[\mathrm{q.iso}^{-1}] \simeq \mathrm{Coh}^s(B, s)/\mathrm{Coh}^s(B, s)^{\mathrm{acy}} \simeq \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}. \quad (3.2.8)$$

As a consequence, if we label $\mathbf{Perf}^s(B, s)$ the full subcategory of $\mathrm{Coh}^s(B, s)$ spanned by perfect $K(B, s)$ -dg modules, there are equivalences of dg categories

$$\mathrm{Coh}^s(B, s)/\mathbf{Perf}^s(B, s) \simeq \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}/\mathbf{Perf}(X_0) \simeq \mathbf{Sing}(X, s). \quad (3.2.9)$$

Construction 3.2.14 We can endow the assignment $(B, s) \rightarrow \mathrm{Coh}^s(B, s)$ with the structure of a pseudo-functor: let $f : (B, s) \rightarrow (C, t)$ be a morphism of affine twisted LG models, i.e. a morphism of A -algebras $f : B \rightarrow C$ such that the induced morphism $id \otimes f : L_B \rightarrow L_C$ sends s to t . Then we can define the dg functor

$$\begin{aligned} - \otimes_B C : \mathrm{Coh}^s(B, s) &\rightarrow \mathrm{Coh}^s(C, t) \\ (E, d, h) &\mapsto (E \otimes_B C, d \otimes id, h \otimes id) \end{aligned} \quad (3.2.10)$$

Indeed, it is clear that if (E, d, h) is a $K(B, s)$ -dg module whose underlying B -module is perfect, then the $K(C, t)$ -dg module $(E \otimes_B C, d \otimes id, h \otimes id)$ is levelwise C -projective and strictly bounded.

This yields a pseudo-functor

$$\mathrm{Coh}^s(\bullet, \bullet) : \mathrm{LG}_{(S,L)}^{\mathrm{aff,op}} \rightarrow \mathrm{dgCat}_S^{\mathrm{lf}}. \quad (3.2.11)$$

Construction 3.2.15 We can endow the pseudo-functor $(B, s) \mapsto \mathrm{Coh}^s(B, s)$ with a weakly associative and weakly unital lax monoidal structure. For any pair of twisted affine LG models $(B, s), (C, t)$, we need to construct a morphism

$$\mathrm{Coh}^s(B, s) \otimes \mathrm{Coh}^s(C, t) \rightarrow \mathrm{Coh}^s(B \otimes_A C, s \boxplus t) \quad (3.2.12)$$

For a pair (B, s) , let $Z(s)$ denote $\mathrm{Spec}(K(B, s))$. Moreover, let

$$\mathbb{V}(L) = \mathrm{Spec}(\mathrm{Sym}(L^\vee)).$$

Then consider the following diagram

$$\begin{array}{ccccc} Z(s) \times_S^h Z(t) & \xrightarrow{\phi} & Z(s \boxplus t) & \longrightarrow & \mathrm{Spec}(B \otimes_A C) \\ \downarrow & & \downarrow & & \downarrow s \times t \\ S & \xrightarrow{\text{zero}} & \mathbb{V}(L) & \xrightarrow{(id, -id)} & \mathbb{V}(L) \times_S \mathbb{V}(L) \\ & & \downarrow & & \downarrow + \\ & & S & \xrightarrow{\text{zero}} & \mathbb{V}(L) \end{array} \quad (3.2.13)$$

where ϕ is the morphism corresponding to

$$\begin{array}{ccccc} K(B, s) \otimes_A K(C, t) & & L_{B \otimes_A C}^\vee & \longrightarrow & L_{B \otimes_A C}^\vee \oplus L_{B \otimes_A C}^\vee & \longrightarrow & B \otimes_A C \\ \uparrow \phi & & & & \uparrow \begin{bmatrix} 1 \\ 1 \end{bmatrix} & & \uparrow 1 \\ K(B \otimes_A C, s \boxplus t) & & L_{B \otimes_A C}^\vee & \longrightarrow & B \otimes_A C & & \end{array} \quad (3.2.14)$$

Consider the two projections

$$Z(s) \xleftarrow{p_s} Z(s) \times_S^h Z(t) \xrightarrow{p_t} Z(t). \quad (3.2.15)$$

Given $(E, d, h) \in \mathrm{Coh}^s(B, s)$ and $(E', d', h') \in \mathrm{Coh}^s(C, t)$, define

$$(E, d, h) \boxtimes (E', d', h') = \phi_*(p_s^*(E, d, h) \otimes p_t^*(E', d', h')). \quad (3.2.16)$$

This is a strictly perfect complex of $B \otimes_A C$ -modules: its underlying complex of A -modules is $(E, d) \otimes_A (E', d')$, which immediately implies that it is strictly bounded. In order to see that each component $(E \otimes_A E')^n = \bigoplus_{i=0}^n E^i \otimes_A E'^{n-i}$ is a projective $B \otimes_A C$ -module, the same argument given in [9, Construction 2.35] applies. The morphism

$$E \otimes_A E' \rightarrow E \otimes_A E' \otimes_{B \otimes_A C} L_{B \otimes_A C}[-1]$$

is $h \otimes 1 + 1 \otimes h'$. In particular, $\phi_*(p_s^*(E, d, h) \otimes p_t^*(E', d', h'))$ lives in $\text{Coh}^s(B \otimes_A C, s \boxplus t)$. The map

$$\underline{A} \rightarrow \text{Coh}^s(A, 0), \quad (3.2.17)$$

where \underline{A} is the unit in $\text{dgCat}_A^{\text{lf}}$, i.e. the dg category with only one object whose endomorphism algebra is A , is defined by the assignment

$$\bullet \mapsto A.$$

It is clear that (3.2.16) and (3.2.17) satisfy the associativity and unity axioms, i.e. they enrich the pseudo-functor (3.2.11) with a lax monoidal structure

$$\text{Coh}^s(\bullet, \bullet)^{\otimes} : \text{LG}_{(S, L)}^{\text{aff, op, } \boxplus} \rightarrow \text{dgCat}_A^{\text{lf, } \otimes}. \quad (3.2.18)$$

As in [9, Construction 2.34, Construction 2.37], consider $\text{Pairs-dgCat}_A^{\text{lf}}$, the category of pairs (T, F) , where $T \in \text{dgCat}_A^{\text{lf}}$ and F is a class of morphisms in T . A morphism $(T, F) \rightarrow (T', F')$ is a dg functor $T \rightarrow T'$ sending F in F' . There is an obvious functor $\text{Pairs-dgCat}_A^{\text{lf}} \rightarrow \text{dgCat}_A^{\text{lf}}$ defined as $(T, F) \mapsto T$. We say that a morphism $(T, F) \rightarrow (T', F')$ in $\text{Pairs-dgCat}_A^{\text{lf}}$ is a DK-equivalence if the dg functor $T \rightarrow T'$ is a DK-equivalence in $\text{dgCat}_A^{\text{lf}}$. Denote this class of morphisms in $\text{Pairs-dgCat}_A^{\text{lf}}$ by W_{DK} .

Also notice that the symmetric monoidal structure on $\text{dgCat}_A^{\text{lf}}$ induces a symmetric monoidal structure on $\text{Pairs-dgCat}_A^{\text{lf}}$

$$(T, F) \otimes (T', F') := (T \otimes T', F \otimes F'). \quad (3.2.19)$$

It is clearly associative and unital, where the unit is $(\underline{A}, \{id\})$. We will denote this symmetric monoidal structure by $\text{Pairs-dgCat}_A^{\text{lf, } \otimes}$. Note that this symmetric monoidal structure is compatible to the class of morphisms in W_{DK} , as we are working with locally-flat dg categories.

Then as in [9, Construction 2.34, Construction 2.37], we can construct a symmetric monoidal ∞ -functor

$$\text{loc}_{dg}^{\otimes} : \text{Pairs-dgCat}_A^{\text{lf, } \otimes} [W_{DK}^{-1}] \rightarrow \text{dgCat}_A^{\text{lf, } \otimes} [W_{DK}^{-1}] \simeq \mathbf{dgCat}_S^{\otimes} \quad (3.2.20)$$

$$\begin{aligned} \mathrm{LG}_A^{\mathrm{aff}, \mathrm{op}, \boxplus} &\rightarrow \mathrm{Pairs}\text{-}\mathrm{dgCat}_A^{\mathrm{lf}, \boxtimes} \\ (B, s) &\mapsto (\mathrm{Coh}^s(B, s), \{q.iso\}) \end{aligned} \quad (3.2.21)$$
$$\begin{array}{c}
 \text{Pairs-dgCat}_A^{\text{lf}, \otimes} \xrightarrow{\text{loc}} \text{Pairs-dgCat}_A^{\text{lf}, \otimes} [W_{DK}^{-1}] \xrightarrow{\text{loc}_{dg}} \text{dgCat}_A^{\text{lf}, \otimes} [W_{DK}^{-1}] \\
 \downarrow \cong \quad \quad \quad \searrow \\
 \text{dgCat}_A^{\otimes} \xrightarrow{\text{loc}} \text{dgCat}_A^{\text{idm}, \otimes}
 \end{array} \tag{3.2.22}$$
$$\mathbf{Coh}^b(\bullet)^{\otimes}_{\mathbf{Perf}(\bullet)} : \mathbf{LG}_A^{\mathrm{op}, \boxplus} \rightarrow \mathbf{dgCat}_A^{\mathrm{idm}, \otimes}. \quad (3.2.23)$$
$$\mathbf{Coh}^b(S_0)_{\mathbf{Perf}(S)}^{\otimes} \simeq \mathbf{Coh}^b(S_0)^{\otimes} \xrightarrow{\sim} \mathbf{Perf}(\mathrm{Sym}_A(L[-2]))^{\otimes}. \quad (3.2.24)$$

In particular, we get the equivalence

$$\mathbf{Perf}(\mathbb{R}\mathrm{Hom}_{(L^\vee \xrightarrow{0} A)}(A, A)) \simeq \mathbf{Coh}^b(S_0).$$

$$\dots \underbrace{L^{\vee \otimes 2}}_{-3} \xrightarrow{0} \underbrace{L^{\vee}}_{-2} \xrightarrow{1} \underbrace{L^{\vee}}_{-1} \xrightarrow{0} \underbrace{A}_{0}. \quad (3.2.25)$$

Applying $\mathrm{Hom}_A(-, A)$ we obtain

$$\begin{aligned} \underbrace{A}_0 &\xrightarrow{0} \underbrace{\mathrm{Hom}_A(L^\vee, A)}_1 \xrightarrow{1} \underbrace{\mathrm{Hom}_A(L^\vee, A)}_2 \xrightarrow{0} \underbrace{\mathrm{Hom}_A(L^{\vee\otimes 2}, A)}_3 \dots \\ &\simeq \underbrace{A}_0 \xrightarrow{0} \underbrace{L}_1 \xrightarrow{1} \underbrace{L}_2 \xrightarrow{0} \underbrace{L^{\otimes 2}}_3 \dots \end{aligned} \quad (3.2.26)$$

which is quasi isomorphic to A . However, when we ask for $(L^\vee \xrightarrow{0} A)$ -linearity, the copies of $L^{\otimes n}$ in odd degree disappear, as the local generators ε in degree -1 of $(L^\vee \xrightarrow{0} A)$ act via the identity on $(L^\vee \xrightarrow{0} A)$. Therefore we find that $\mathbb{R}\mathrm{Hom}_{(L^\vee \xrightarrow{0} A)}(A, A)$ is quasi-isomorphic to

$$\underbrace{A}_0 \rightarrow \underbrace{0}_1 \rightarrow \underbrace{L}_2 \rightarrow \dots \simeq \mathrm{Sym}_A(L[-2]). \quad (3.2.27)$$

This shows that there exists an equivalence

$$\mathbb{R}\mathrm{Hom}_{(L^\vee \xrightarrow{0} A)}(A, A) \simeq \mathrm{Sym}_A(L[-2]) \quad (3.2.28)$$

as $(L^\vee \xrightarrow{0} A)$ -dg modules. Notice that both these objects carry a canonical algebra structure. In order to conclude that the two commutative algebra structures coincide, we consider the dg functor

$$\mathbb{R}\mathrm{Hom}_{(L^\vee \xrightarrow{0} A)}(A, -) : \mathrm{Coh}^s(S, 0) \rightarrow \mathrm{Sym}_A(L[-2]) - \mathrm{dgmod}. \quad (3.2.29)$$

Notice that $\mathrm{Sym}_A(L[-2])$ is a strict cdga, seen as a commutative algebra object in $\mathbf{QCoh}(A)$. Similarly to [9, Lemma 2.39], for (E, d, h) an object in $\mathrm{Coh}^s(A, 0)$, the $\mathrm{Sym}_A(L[-2])$ -dg module $\mathbb{R}\mathrm{Hom}_{(L^\vee \xrightarrow{0} A)}(A, (E, d, h))$ can be computed (in degrees i and $i + 1$) as

$$\bigoplus_{n \geq 0} E_{i-2n} \otimes_A L^{\otimes n} \xrightarrow{d+h} \bigoplus_{n \geq 0} E_{i+1-2n} \otimes_A L^{\otimes n}. \quad (3.2.30)$$

The same arguments given in *loc.cit.* hold mutatis mutandis in our situation and therefore we obtain a symmetric monoidal functor

$$\mathrm{Coh}^s(S, 0)^{\otimes} \rightarrow \mathrm{Sym}_A(L[-2]) - \mathrm{dgmod}^{\otimes} \quad (3.2.31)$$

which preserves quasi-isomorphisms. If we localize both the l.h.s. and the r.h.s. we thus obtain a symmetric monoidal ∞ -functor

$$\mathbf{Coh}^b(S_0)^{\otimes} \rightarrow \mathbf{QCoh}(\mathrm{Sym}_A(L[-2]))^{\otimes}, \quad (3.2.32)$$

from which one recovers the equivalence of symmetric monoidal dg categories

$$\mathbf{Coh}^b(S_0)^\otimes \simeq \mathbf{Perf}(\mathrm{Sym}_A(L[-2]))^\otimes. \quad (3.2.33)$$

□

Corollary 3.2.17 *Let $S = \mathrm{Spec}(A)$ be a regular noetherian affine scheme and let L be a line bundle over S . The lax monoidal ∞ -functors (3.2.23) factors as*

$$\mathbf{Coh}^b(\bullet)_{\mathbf{Perf}(\bullet)}^\otimes : \mathrm{LG}_{(S,L)}^{\mathrm{op}, \boxplus} \rightarrow \mathbf{Mod}_{\mathbf{Perf}(\mathrm{Sym}_A(L[-2]))}(\mathbf{dgCat}_S^{\mathrm{idm}})^\otimes. \quad (3.2.34)$$

Then, for a noetherian regular scheme S with a line bundle \mathcal{L}_S , we obtain a lax monoidal ∞ -functor

$$\mathbf{Coh}^b(\bullet)_{\mathbf{Perf}(\bullet)}^\otimes : \mathrm{LG}_{(S, \mathcal{L}_S)}^{\mathrm{op}, \otimes} \rightarrow \mathbf{dgCat}_S^{\mathrm{idm}, \otimes} \quad (3.2.35)$$

as the limit

$$\lim_{(Spec(A), \mathcal{L}_{S|Spec(A)})} (\mathbf{Coh}^b(\bullet)_{\mathbf{Perf}(\bullet)}^\otimes : \mathrm{LG}_{Spec(A), \mathcal{L}_{S|Spec(A)}}^{\mathrm{op}, \boxplus} \rightarrow \mathbf{dgCat}_A^{\mathrm{idm}, \otimes}) \quad (3.2.36)$$

where $Spec(A) \rightarrow S$ is a Zariski open subscheme. We have used Lemma 3.1.9 and the definition

$$\lim_{Spec(A) \rightarrow S} \mathbf{dgCat}_A^{\mathrm{idm}, \otimes} = \mathbf{dgCat}_S^{\mathrm{idm}, \otimes}. \quad (3.2.37)$$

Remark 3.2.18 The monoidal structure on $\mathbf{Coh}^b(\bullet)_{\mathbf{Perf}(\bullet)}^\otimes$ implies that each dg category $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$ is endowed with an action of $\mathbf{Coh}^b(S_0)$ (recall that we are assuming that S is regular). Similarly to [9, Remark 2.38] we can describe this action. Consider the diagram

$$\begin{array}{ccccc} X_0 \times_S^h S_0 & \xrightarrow{a} & X_0 & & \\ p_{X_0} \swarrow & \downarrow i & \swarrow i & & \downarrow \\ X_0 & \xrightarrow{\quad} & X & & S \\ \downarrow & p_{S_0} \downarrow & \downarrow & & \downarrow \\ & S_0 & \xrightarrow{\quad} & S & \\ \downarrow & \swarrow & \downarrow & \swarrow & \\ S & \xrightarrow{\text{zero}} & \mathbb{V}(\mathcal{L}_S) & & \end{array} \quad (3.2.38)$$

Notice that $X_0 \times_S^h S_0 \simeq X_0 \times_X^h X_0$. Given $\mathcal{E} \in \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$ and $\mathcal{F} \in \mathbf{Coh}^b(S_0)$, then $\mathcal{E} \boxtimes \mathcal{F} = a_*(p_{X_0}^* \mathcal{E} \otimes p_{S_0}^* \mathcal{F})$. In particular, when $\mathcal{F} = \mathcal{L}_S^\vee \xrightarrow{0} \mathcal{O}_S = \mathcal{O}_{S_0}$, then

$$\begin{aligned} \mathcal{E} \boxtimes (\mathcal{L}_S^\vee \xrightarrow{0} \mathcal{O}_S) &= a_*(p_{X_0}^* \mathcal{E} \otimes p_{S_0}^* \mathcal{O}_{S_0}) \simeq a_*(p_{X_0}^* \mathcal{E} \otimes a^* \mathcal{O}_{X_0}) \\ &\underset{\text{proj. form.}}{\simeq} a_* p_{X_0}^* \mathcal{E} \underset{\text{der. prop. base change}}{\simeq} i^* i_* \mathcal{E}. \end{aligned} \quad (3.2.39)$$

When $\mathcal{F} = \mathcal{O}_S = t_* \mathcal{O}_S$, where $t : S = \pi_0(S_0) \rightarrow S_0$ is the truncation morphism, the (homotopy) cartesian square

$$\begin{array}{ccc} X_0 \times_S S & \xrightarrow{id \times t} & X_0 \times_S^h S_0 \\ \downarrow q & & \downarrow p_{S_0} \\ S & \xrightarrow{t} & S_0 \end{array} \quad (3.2.40)$$

implies that

$$\begin{aligned} \mathcal{E} \boxtimes \mathcal{O}_S &= a_*(p_{X_0}^* \mathcal{E} \otimes p_{S_0}^* t_* \mathcal{O}_S) \underset{\text{der. prop. base change}}{\simeq} a_*(p_{X_0}^* \mathcal{E} \otimes (id \times t)_* \mathcal{O}_{X_0}) \\ &\underset{\text{proj. form.}}{\simeq} a_*(id \times t)_*((id \times t)^* p_{X_0}^* \mathcal{E}) \simeq \mathcal{E}. \end{aligned} \quad (3.2.41)$$

Finally, we can consider \mathcal{L}_S^\vee endowed with the trivial $\mathcal{L}_S^\vee \xrightarrow{0} \mathcal{O}_S$ dg module structure, namely $t_* \mathcal{L}_S^\vee$. Then, for any $\mathcal{E} \in \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$, we have

$$\mathcal{E} \boxtimes t_* \mathcal{L}_S^\vee = a_*(p_{X_0}^* \mathcal{E} \otimes p_{S_0}^* t_* \mathcal{L}_S^\vee) \underset{\text{der. prop. base change}}{\simeq} a_*(p_{X_0}^* \mathcal{E} \otimes (id \times t)_* q^* \mathcal{L}_S^\vee). \quad (3.2.42)$$

Notice that $q^* \mathcal{L}_S^\vee \simeq \mathcal{L}_X^\vee \otimes_{\mathcal{O}_X} \mathcal{O}_{X_0}$, thus we can continue

$$\simeq a_*(p_{X_0}^* \mathcal{E} \otimes (id \times t)_*(\mathcal{L}_X^\vee \otimes_{\mathcal{O}_X} \mathcal{O}_{X_0})) \simeq \mathcal{E} \otimes_{\mathcal{O}_{X_0}} \mathcal{O}_{X_0} \otimes_{\mathcal{O}_X} \mathcal{L}_X^\vee \simeq \mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{L}_X^\vee. \quad (3.2.43)$$

We will construct, using $\mathbf{Coh}^b(\bullet)_{\mathbf{Perf}(\bullet)}^{\otimes}$, a lax monoidal ∞ -functor

$$\mathbf{Sing}(\bullet, \bullet)^{\otimes} : \mathbf{LG}_{(S, \mathcal{L}_S)}^{\text{op}, \boxplus} \rightarrow \mathbf{dgCat}_S^{\text{idm}, \otimes}, \quad (3.2.44)$$

following [45] and [9].

Construction 3.2.19 As in Lemma 3.2.16, consider the strict commutative dg algebra $\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2])$ as a commutative algebra object in $\mathbf{QCoh}(S)$. Consider the $\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2])$ -algebra

$$\mathcal{R} := \mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2])[v^{-1}] \quad (3.2.45)$$

where v is the generator in degree 2. More explicitly,

$$\mathcal{R} = \cdots \rightarrow \underbrace{\mathcal{L}_S^{\vee \otimes 2}}_{-4} \rightarrow 0 \rightarrow \underbrace{\mathcal{L}_S^{\vee}}_{-2} \rightarrow 0 \rightarrow \underbrace{\mathcal{O}_S}_{0} \rightarrow 0 \rightarrow \underbrace{\mathcal{L}_S}_{2} \rightarrow 0 \rightarrow \underbrace{\mathcal{L}_S^{\otimes 2}}_{4} \rightarrow \cdots \quad (3.2.46)$$

Then we have a symmetric monoidal ∞ -functor $-\otimes_{\mathbf{Perf}(\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} \mathbf{Perf}(\mathcal{R})$

$$\mathbf{Mod}_{\mathbf{Perf}(\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))}(\mathbf{dgCat}_S^{\mathrm{idm}})^{\otimes} \rightarrow \mathbf{Mod}_{\mathbf{Perf}(\mathcal{R})}(\mathbf{dgCat}_S^{\mathrm{idm}})^{\otimes}. \quad (3.2.47)$$

Composing it with (3.2.34) we obtain a lax monoidal ∞ -functor

$$\mathrm{LG}_{(S, \mathcal{L}_S)}^{\mathrm{op}, \boxplus} \rightarrow \mathbf{Mod}_{\mathbf{Perf}(\mathcal{R})}(\mathbf{dgCat}_S^{\mathrm{idm}})^{\otimes} \quad (3.2.48)$$

which, at the level of objects, is defined by

$$(X, s_X) \mapsto \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \otimes_{\mathbf{Perf}(\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} \mathbf{Perf}(\mathcal{R}).$$

Remark 3.2.20 Let $\mathcal{U} = \{U_i = \mathrm{Spec}(A_i)\}$ be a Zariski affine covering of S , such that $\mathcal{L}_{S|U_i}$ is equivalent to A_i . Then the restriction of \mathcal{R} to U_i is equivalent to the dg algebra $A_i[u, u^{-1}]$, where u sits in cohomological degree 2.

Lemma 3.2.21 Let $\mathcal{U} = \{U_i = \mathrm{Spec}(B_i)\}$ be an affine open covering which trivializes \mathcal{L}_S and \mathcal{L}_S^{\vee} .

Under the equivalence (3.2.24), the full sub-category $\mathbf{Perf}(S_0)$ of $\mathbf{Coh}^b(S_0)$ corresponds to the full sub-category of $\mathbf{Perf}(\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))$ spanned by perfect $\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2])$ -modules whose restriction to each U_i is a u -torsion perfect $B_i[u]$ -dg module, that we will denote $\mathbf{Perf}(\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))^{\mathrm{tors}}$.

Proof Let $\mathcal{M} \in \mathbf{Perf}(S_0)$. By definitions this means that, for every Zariski open $f : \mathrm{Spec}(A) \rightarrow S_0$, the pullback $f^*\mathcal{M} \in \mathbf{Perf}(A)$. Consider our covering \mathcal{U} . Then, for every $i \in I$, $\mathcal{M}|_{U_i}$ is a perfect B_i -module. By the same argument given above, we get equivalences $\mathbf{Coh}^b(U_{i,0})^{\otimes} \simeq \mathbf{Perf}(B_i[u])^{\otimes}$ and the argument provided in [9, Proposition 2.43] guarantees that $\mathbf{Perf}(U_{i,0})$ corresponds to $\mathbf{Perf}(B_i[u])^{u-\mathrm{tors}}$ under this equivalence. This shows that every perfect complex is locally of u -torsion. The converse follows by the local characterization of perfect complexes and by the characterization given in [9, Proposition 2.43]. \square

Lemma 3.2.22 *The dg quotient $\mathbf{Coh}^b(S_0) \rightarrow \mathbf{Sing}(S_0)$ corresponds to the base-change*

$$- \otimes_{\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2])} \mathcal{R} : \mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2])) \rightarrow \mathbf{Perf}(\mathcal{R}) \quad (3.2.49)$$

under the equivalence (3.2.24)

Proof This follows by the fact that the categories involved satisfy Zariski descent, by the previous lemmas and by [9, Proposition 2.43]. \square

Lemma 3.2.23 *There is an equivalence of dg categories*

$$\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \otimes_{\mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} \mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))^{\mathrm{tors}} \simeq \mathbf{Perf}(X_0). \quad (3.2.50)$$

Proof By definition,

$$\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \otimes_{\mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} \mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))^{\mathrm{tors}} \quad (3.2.51)$$

is the subcategory spanned by locally u -torsion modules in $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$. We will show that they coincide with the subcategory of perfect complexes, using [9, Proposition 2.45]. Suppose that \mathcal{M} is in $\mathbf{Perf}(X_0)$. Then, for every affine open covering $j : \mathrm{Spec}(A) \rightarrow X_0$ which trivializes \mathcal{L}_{X_0} , $j^*\mathcal{M}$ is perfect, and thus it is a u -torsion perfect $A[u]$ module by *loc. cit.* Conversely, if \mathcal{M} is locally u -torsion, each restriction $j^*\mathcal{M}$ is u -torsion, i.e. perfect. This proves the lemma. \square

Corollary 3.2.24 *Let (X, s_X) be a twisted LG model over (S, \mathcal{L}_S) . Then the exact sequence in $\mathbf{dgCat}_S^{\mathrm{idm}}$*

$$\mathbf{Perf}(X_0) \rightarrow \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \rightarrow \mathbf{Sing}(X, s_X) \quad (3.2.52)$$

is equivalent to

$$\begin{array}{ccc} \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \otimes_{\mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} \mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))^{\mathrm{tors}} & & \\ \downarrow & & \\ \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} & & (3.2.53) \\ \downarrow & & \\ \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \otimes_{\mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} \mathbf{Perf}(\mathcal{R}). & & \end{array}$$

Proof This is an obvious consequence of the previous lemmas and of the fact that the ∞ -functor

$$T \mapsto \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \otimes_{\mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))} T$$

preserves exact sequences in $\mathbf{Mod}_{\mathbf{Perf}(\mathbf{Sym}_{\mathcal{O}_S}(\mathcal{L}_S[-2]))}(\mathbf{dgCat}_S^{\mathrm{idm}})$. \square

Construction 3.2.25 Consider the following lax monoidal ∞ -functor

$$\begin{aligned} \mathrm{LG}_{(S, \mathcal{L}_S)}^{\mathrm{op}, \boxplus} &\xrightarrow{\mathrm{Coh}^b(\bullet)^{\otimes} \mathrm{Perf}(\bullet)} \mathbf{ModPerf}(\mathrm{Sym}_{\odot_S}(\mathcal{L}_S[-2]))(\mathbf{dgCat}_S^{\mathrm{idm}})^{\otimes} \\ &\quad \downarrow - \otimes_{\mathrm{Sym}_{\odot_S}(\mathcal{L}_S[-2])} \mathcal{R} \\ &\quad \mathbf{ModPerf}(\mathcal{R})(\mathbf{dgCat}_S^{\mathrm{idm}})^{\otimes}. \end{aligned} \quad (3.2.54)$$

The previous corollary means that its underlying ∞ -functor, composed with the forgetful functor

$$\mathbf{ModPerf}(\mathcal{R})(\mathbf{dgCat}_S^{\mathrm{idm}}) \rightarrow \mathbf{dgCat}_S^{\mathrm{idm}}, \quad (3.2.55)$$

is defined on objects by the assignment

$$(X, s) \mapsto \mathbf{Sing}(X, s).$$

3.3 The motivic realization of $\mathbf{Sing}(X, s_X)$

Recall from Sect. 2.6 that there is a motivic realization lax monoidal ∞ -functor

$$\mathcal{M}_S^{\vee} : \mathbf{dgCat}_S^{\mathrm{idm}, \otimes} \rightarrow \mathbf{Mod}_{\mathrm{BU}_S}(\mathbf{SH}_S)^{\otimes} \quad (3.3.1)$$

with the following properties

- \mathcal{M}_S^{\vee} commutes with filtered colimits.
- For every dg category T , $\mathcal{M}_S^{\vee}(T)$ is the spectrum

$$Y \in \mathbf{Sm}_S \mapsto \mathrm{HK}(\mathbf{Perf}(Y) \otimes_S T).$$

- \mathcal{M}_S^{\vee} sends exact sequences of dg categories to fiber-cofiber sequences in $\mathbf{Mod}_{\mathrm{BU}_S}(\mathbf{SH}_S^{\otimes})$.

Our main scope in this section is to study the motivic realization of the dg category $\mathbf{Sing}(X, s_X)$ associated to $(X, s_X) \in \mathrm{LG}_{(S, \mathcal{L}_S)}$, under the assumption that X is regular. The first important fact is the following one:

Proposition 3.3.1 [9, Proposition 3.13] *Let $p : X \rightarrow S$ be in \mathbf{Sch}_S . Then $\mathcal{M}_X^{\vee}(\mathbf{Perf}(X)) \simeq \mathrm{BU}_X$ (see Sect. 2.5.1 for notation).*

Proof Consider the construction given in Sect. 2.6 with S replaced by X :

$$\mathbf{Sm}_X^{\times} \xrightarrow{\mathrm{Perf}(\bullet)} \mathbf{dgCat}^{\mathrm{idm}, \mathrm{op}, \otimes} \xrightarrow{\iota} \mathbf{SH}_X^{\mathrm{nc}, \mathrm{op}, \otimes} \xrightarrow{\mathbb{R}\mathrm{Hom}(-, 1_X^{\mathrm{nc}})} \mathbf{SH}_X^{\mathrm{nc}, \otimes} \xrightarrow{\mathcal{M}_X} \mathbf{SH}_X^{\otimes}. \quad (3.3.2)$$

We wish to show that $\mathcal{M}_X(\mathbb{R}\mathrm{Hom}(\iota \circ \mathbf{Perf}(X), 1_X^{nc})) \simeq \mathrm{BU}_X$. Notice that since the ∞ -functor above is right lax monoidal and $\mathbf{Perf}(X)$ is a commutative algebra in $\mathbf{dgCat}^{\mathrm{idm}, \mathrm{op}, \otimes}$, there is a canonical morphism of commutative algebras

$$\mathrm{BU}_X \rightarrow \mathcal{M}_X(\mathbb{R}\mathrm{Hom}(\iota \circ \mathbf{Perf}(X), 1_X^{nc})). \quad (3.3.3)$$

It suffices to show that they represent the same ∞ -functor. Let \mathcal{Y} be an object in \mathbf{SH}_X^\otimes . Then

$$\begin{aligned} & \mathrm{Map}_{\mathbf{SH}_X}(\mathcal{Y}, \mathcal{M}_X(\mathbb{R}\mathrm{Hom}(\iota \circ \mathbf{Perf}(X), 1_X^{nc}))) \\ & \simeq \mathrm{Map}_{\mathbf{SH}_X^{nc}}(\mathrm{RPerf}(\mathcal{Y}), \mathbb{R}\mathrm{Hom}(\iota \circ \mathbf{Perf}(X), 1_X^{nc})) \\ & \simeq \mathrm{Map}_{\mathbf{SH}_X^{nc}}(\mathrm{RPerf}(\mathcal{Y}) \otimes \mathrm{RPerf}(\Sigma_+^\infty X), 1_X^{nc}) \simeq \mathrm{Map}_{\mathbf{SH}_X^{nc}}(\mathrm{RPerf}(\mathcal{Y} \otimes \Sigma_+^\infty X), 1_X^{nc}) \\ & \simeq \mathrm{Map}_{\mathbf{SH}_X}(\mathrm{RPerf}(\mathcal{Y}), 1_X^{nc}) \simeq \mathrm{Map}_{\mathbf{SH}_X}(\mathcal{Y}, \mathcal{M}_X(1_X^{nc})) \simeq \mathrm{Map}_{\mathbf{SH}_X}(\mathcal{Y}, \mathrm{BU}_X) \end{aligned} \quad (3.3.4)$$

where we have used that $\iota \circ \mathbf{Perf}(\bullet) \simeq \mathrm{RPerf} \circ \Sigma_+^\infty$, which are all symmetric monoidal functors, $\Sigma_+^\infty(X)$ is the unit in \mathbf{SH}_X^\otimes , that RPerf is left adjoint to \mathcal{M}_X and Robalo's result $\mathcal{M}_X(1_X^{nc}) \simeq \mathrm{BU}_X$ ([49, Theorem 1.8]).

The fact that the equivalence holds in $\mathrm{CAlg}(\mathbf{SH}_X^\otimes)$ follows immediately from the conservativity of the forgetful functor

$$\mathrm{CAlg}(\mathbf{SH}_X^\otimes) \rightarrow \mathbf{SH}_X^\otimes. \quad (3.3.5)$$

□

Remark 3.3.2 Notice that since the equivalence $\mathrm{BU}_X \rightarrow \mathcal{M}_X^\vee(\mathbf{Perf}(X))$ holds in $\mathrm{CAlg}(\mathbf{SH}_X)$, if we consider a perfect complex \mathcal{E} scheme X , then the image of

$$\mathbf{Perf}(X) \xrightarrow{-\otimes \mathcal{E}} \mathbf{Perf}(X) \quad (3.3.6)$$

along \mathcal{M}_X^\vee coincides with multiplication by the class $[\mathcal{E}] \in \mathrm{HK}_0(X)$ in the commutative algebra BU_X , which we denote $m_{\mathcal{E}}$. In a formula: $\mathcal{M}_X^\vee(- \otimes \mathcal{E}) \simeq m_{\mathcal{E}} \in \mathrm{Map}_{\mathrm{BU}_X}(\mathrm{BU}_X, \mathrm{BU}_X)$. Moreover, for any exact triangle $\mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{E}''$ we get $[\mathcal{E}] = [\mathcal{E}'] + [\mathcal{E}''] \in \mathrm{HK}_0(X)$. Then $m_{\mathcal{E}} = m_{\mathcal{E}'} + m_{\mathcal{E}''}$. In particular, $m_{\mathcal{E}[1]} = -m_{\mathcal{E}}$.

The next step will be understanding the motivic realization of the category of coherent bounded complexes $\mathbf{Coh}^b(Z)$ of an S -scheme Z , at least when it is possible to regard it as a closed subscheme of a regular S -scheme X .

Proposition 3.3.3 [9, Proposition 3.17] *Let $p : X \rightarrow S$ be a regular S -scheme of finite type. Consider a closed subscheme $i : Z \rightarrow X$ and label $j : U \rightarrow X$ the embedding of the complementary open subscheme. Then there is an equivalence*

$$\mathcal{M}_X(\mathbf{Coh}^b(Z)) \simeq i_* i^! \mathrm{BU}_X. \quad (3.3.7)$$

Proof Consider the exact sequence of dg categories

$$\mathbf{Coh}^b(X)_Z \rightarrow \mathbf{Coh}^b(X) \xrightarrow{j^*} \mathbf{Coh}^b(U) \quad (3.3.8)$$

where $\mathbf{Coh}^b(X)_Z$ denotes the subcategory of objects in $\mathbf{Coh}^b(X)$ whose support is in Z .

Recall that, if X is a scheme, an \mathcal{O}_X -linear dg category is a sheaf of dg categories on the Zariski site of X . Then the above sequence of dg categories is exact by definition and if we apply \mathcal{M}_X^\vee to it we obtain a fiber cofiber sequence in \mathbf{SH}_X (see [69, Remark 2.2.2]).

The regularity hypothesis imposed on X implies that $\mathbf{Coh}^b(X) \simeq \mathbf{Perf}(X)$ and $\mathbf{Coh}^b(U) \simeq \mathbf{Perf}(U)$. If we apply \mathcal{M}_X^\vee and use the previous proposition, we obtain

$$\mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z) \rightarrow \mathbf{BU}_X \xrightarrow{\mathcal{M}_X^\vee(j^*)} \mathcal{M}_X^\vee(\mathbf{Perf}(U)), \quad (3.3.9)$$

which is a fiber-cofiber sequence in \mathbf{SH}_X (since \mathcal{M}_X^\vee sends exact triangles of dg categories to fiber-cofiber sequences). Moreover, \mathcal{M}_X^\vee is compatible with pushforwards and $\mathcal{M}_X^\vee(j^*) \sim j^*$. As the spectrum \mathbf{BU} of non-connective homotopy-invariant K-theory is compatible with pullbacks (see [15]), the previous fiber-cofiber sequence is nothing but

$$\mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z) \rightarrow \mathbf{BU}_X \rightarrow j_* j^* \mathbf{BU}_X, \quad (3.3.10)$$

where the map on the right is induced by the unit of the adjunction (j^*, j_*) . In particular, we get a canonical equivalence

$$\mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z) \simeq i_* i^! \mathbf{BU}_X \quad (3.3.11)$$

and therefore we are left to show that we have an equivalence

$$\mathcal{M}_X^\vee(\mathbf{Coh}^b(Z)) \simeq \mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z). \quad (3.3.12)$$

Here $\mathbf{Coh}^b(Z)$ is the sheaf of dg categories

$$V = \text{Spec}(R) \mapsto \mathbf{Coh}^b(Z \times_X V), \quad (3.3.13)$$

where V is an affine open subscheme of X .

Notice that there is a canonical morphism

$$\mathcal{M}_X^\vee(i_*) : \mathcal{M}_X^\vee(\mathbf{Coh}^b(Z)) \rightarrow \mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z).$$

The collection of objects $\Sigma_+^\infty Y \otimes \mathbf{BU}_X$, where $Y \in \mathbf{Sm}_X$, forms a family of compact generators of $\mathbf{Mod}_{\mathbf{BU}_X}(\mathbf{SH}_X)$. As \mathcal{M}_X^\vee commutes with filtered colimits, it suffices to

show that the morphism

$$\begin{array}{c} \mathrm{Map}_{\mathrm{BU}_X}(\Sigma_+^\infty Y \otimes \mathrm{BU}_X, \mathcal{M}_X^\vee(\mathbf{Coh}^b(Z))) \\ \downarrow \\ \mathrm{Map}_{\mathrm{BU}_X}(\Sigma_+^\infty Y \otimes \mathrm{BU}_X, \mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z)) \end{array} \quad (3.3.14)$$

is an equivalence of spectra. Notice that

$$\begin{aligned} \mathrm{Map}_{\mathrm{BU}_X}(\Sigma_+^\infty Y \otimes \mathrm{BU}_X, \mathcal{M}_X^\vee(\mathbf{Coh}^b(Z))) &\simeq \mathrm{Map}_{\mathrm{SH}_X}(\Sigma_+^\infty Y, \mathcal{M}_X^\vee(\mathbf{Coh}^b(Z))) \\ &\simeq \mathrm{Map}_{\mathrm{SH}_X^{nc}}(\mathbf{RPerf} \circ \Sigma_+^\infty Y, \mathbb{R}\mathrm{Hom}(\mathbf{Coh}^b(Z), 1_X^{nc})) \\ &\simeq \mathrm{Map}_{\mathrm{SH}_X^{nc}}(\mathbf{Perf}(Y) \otimes \mathbf{Coh}^b(Z), 1_X^{nc}). \end{aligned} \quad (3.3.15)$$

By [45], $\mathbf{Perf}(Y) \otimes \mathbf{Coh}^b(Z) \simeq \mathbf{Coh}^b(Y) \otimes \mathbf{Coh}^b(Z) \simeq \mathbf{Coh}^b(Y \times_X Z)$, and therefore the spectrum above coincides with $\mathrm{HK}(\mathbf{Coh}^b(Y \times_X Z))$, which by the homotopy-invariance of G -theory and by the theorem of the Heart ([7, Corollary 6.4.1]) coincides with the G -theory of $Y \times_X Z$. In the same way, we obtain that $\mathrm{Map}_{\mathrm{SH}_X}(\Sigma_+^\infty Y, \mathcal{M}_X^\vee(\mathbf{Coh}^b(X)_Z))$ coincides with the G -theory spectrum of the abelian category $\mathbf{Coh}(Y)_{Y \times_X Z}$ (the heart of the dg category $\mathbf{Coh}^b(Y)_{Y \times_X Z}$). The claim now follows from Quillen's dévissage. \square

Remark 3.3.4 An anonymous referee has pointed out that in order to deal with \mathcal{O}_X -linear dg categories one can use the theory of 1-affiness of Gaiitsgory [26]. One can use this to give an alternative proof of the proposition above.

As a final observation, we remark that the assignment $Z \mapsto \mathcal{M}_S^\vee(\mathbf{Coh}^b(Z))$ is insensible to (derived) thickenings.

Lemma 3.3.5 [9, Proposition 3.24] *Let Z be a derived scheme of finite type over S and let $t : \pi_0(Z) \rightarrow Z$ be the canonical closed embedding of the underlying scheme of Z . Then*

$$\mathcal{M}_S^\vee(\mathbf{Coh}^b(Z)) \simeq \mathcal{M}_S^\vee(\mathbf{Coh}^b(\pi_0(Z))). \quad (3.3.16)$$

Proof By the proof of the previous proposition, $\mathcal{M}_S^\vee(\mathbf{Coh}^b(Z))$ represents the spectra valued sheaf $Y \mapsto \mathrm{HK}(\mathbf{Coh}^b(Y \times_S Z))$. Similarly, $\mathcal{M}_S^\vee(\mathbf{Coh}^b(\pi_0(Z)))$ represents the spectra-valued sheaf $Y \mapsto \mathrm{HK}(\mathbf{Coh}^b(Y \times_S \pi_0(Z)))$. Notice that $\pi_0(Y \times_S Z) = \pi_0(Y) \times_{\pi_0(S)} \pi_0(Z) = Y \times_S \pi_0(Z)$ and that the heart of $\mathbf{Coh}^b(Y \times_S Z)$ is equivalent to the heart of $\mathbf{Coh}^b(Y \times_S \pi_0(Z))$. Then the theorem of the Heart and the computation above allows us to conclude. \square

Now consider a twisted LG model (X, s_X) over (S, \mathcal{L}_S) and assume that X is a regular scheme. As above, let $X_0 \xrightarrow{i} X$ be the derived zero section of s_X in X and let $j : X_{\mathcal{U}} = X - X_0 \rightarrow X$ be the corresponding open embedding. In this case, we

have that $\mathbf{Sing}(X, s_X) \simeq \mathbf{Sing}(X_0)$. Consider the diagram in $\mathbf{dgCat}_X^{\text{idm}}$ and its image in \mathbf{SH}_X

$$\begin{array}{ccccc} \mathbf{Perf}(X_0) & \hookrightarrow & \mathbf{Coh}^b(X_0) & \longrightarrow & \mathbf{Sing}(X_0) \\ & \searrow i_* & \downarrow i_* & & \\ & & \mathbf{Perf}(X) & & \end{array} \quad (3.3.17)$$

$$\begin{array}{ccccc} & & \downarrow j^* & & \\ & & \mathbf{Perf}(X_U) & & \\ \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) & \longrightarrow & \mathcal{M}_X^\vee(\mathbf{Coh}^b(X_0)) & \longrightarrow & \mathcal{M}_X^\vee(\mathbf{Sing}(X_0)) \\ & \searrow \mathcal{M}^\vee(i_*) & \downarrow \mathcal{M}_X^\vee(i_*) & & \\ \mathcal{M}_X^\vee & & \mathcal{M}_X^\vee(\mathbf{Perf}(X)) & & \\ \sim & & \downarrow j^* & & \\ & & \mathcal{M}_X^\vee(\mathbf{Perf}(X_U)) & & \end{array} \quad (3.3.18)$$

By the previous results, the second diagram can be rewritten and completed as follows

$$\begin{array}{ccccc} \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) & \longrightarrow & i_* i^! \mathbf{BU}_X & \longrightarrow & \mathcal{M}_X^\vee(\mathbf{Sing}(X_0)) \\ & \searrow \mathcal{M}^\vee(i_*) & \downarrow \mathcal{M}_X^\vee(i_*) & & \\ j_! j^* \mathbf{BU}_X & \longrightarrow & \mathbf{BU}_X & \longrightarrow & i_* i^* \mathbf{BU}_X \\ & \searrow \alpha & \downarrow \text{counit}(j^*, j_*) & & \\ & & j_* j^* \mathbf{BU}_X & & \end{array} \quad (3.3.19)$$

where $i = i \circ t : \pi_0(X_0) \rightarrow X$ is the closed embedding of the underlying scheme of X_0 in X .

Remark 3.3.6 The morphism $\mathbf{BU}_X \rightarrow i_* i^* \mathbf{BU}_X$ can be factored as

$$\underbrace{\mathcal{M}_X^\vee(\mathbf{Perf}(X))}_{\mathbf{BU}_X \simeq} \xrightarrow{\mathcal{M}_X^\vee(i^*)} \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) \xrightarrow{\mathcal{M}_X^\vee(t^*)} \underbrace{\mathcal{M}_X^\vee(\mathbf{Perf}(\pi_0(X_0)))}_{\simeq i_* i^* \mathbf{BU}_X}. \quad (3.3.20)$$

Lemma 3.3.7 *The endomorphism*

$$\mathcal{M}_X^\vee(i^*) \circ \mathcal{M}_X^\vee(i_*) : \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) \rightarrow \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) \quad (3.3.21)$$

is homotopic to $1 - m_{\mathcal{L}_{X_0}^\vee}$, where $m_{\mathcal{L}_{X_0}^\vee}$ is the autoequivalence induced by $-\otimes \mathcal{L}_{X_0}^\vee : \mathbf{Perf}(X_0) \rightarrow \mathbf{Perf}(X_0)$.

Proof Indeed, by Remark 3.2.18, the endomorphism i^*i_* is equivalent to $-\boxtimes (\mathcal{L}_S^\vee \xrightarrow{0} \mathcal{O}_S)$, the identity is equivalent to $-\boxtimes \mathcal{O}_S$ and $-\boxtimes \mathcal{L}_S^\vee$ corresponds to $-\otimes \mathcal{L}_{X_0}^\vee : \mathbf{Perf}(X_0) \rightarrow \mathbf{Perf}(X_0)$. Then, considering the cofiber sequence of dg functors

$$\begin{array}{ccc} -\otimes_{\mathcal{O}_S} \mathcal{L}_S^\vee & \xrightarrow{0} & -\otimes_{\mathcal{O}_S} \mathcal{O}_S \simeq id \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & i^*i_* \end{array} \quad (3.3.22)$$

we see that i^*i_* is equivalent to the dg functor

$$\begin{aligned} \mathbf{Perf}(X_0) &\rightarrow \mathbf{Perf}(X_0) \\ \mathcal{E} &\mapsto \mathcal{E} \oplus \mathcal{E} \otimes_{\mathcal{O}_{X_0}} \mathcal{L}_{X_0}^\vee[1]. \end{aligned} \quad (3.3.23)$$

By Remark 3.3.2 we conclude that

$$\mathcal{M}_X^\vee(i^*i_*) \simeq m_{\mathcal{O}_{X_0} \oplus \mathcal{L}_{X_0}^\vee[1]} \simeq 1 - m_{\mathcal{L}_{X_0}^\vee}. \quad (3.3.24)$$

□

Corollary 3.3.8 *Let $(X, s_X) \in \mathbf{LG}_{(S, \mathcal{L}_S)}$ and assume that X is regular. Then there is a fiber-cofiber sequence in $\mathbf{Mod}_{\mathbf{BU}_X}(\mathbf{SH}_X)$*

$$\begin{aligned} &\mathcal{M}_X^\vee(\mathbf{Sing}(X, s_X)) \\ &\quad \downarrow \\ &\text{cofiber}(\mathcal{M}_X^\vee(t^*) \circ (1 - m_{\mathcal{L}_{X_0}^\vee}) : \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) \rightarrow i_*i^*\mathbf{BU}_X) \\ &\quad \downarrow \\ &\text{cofiber}(\alpha : j_!j^*\mathbf{BU}_X \rightarrow j_*j^*\mathbf{BU}_X). \end{aligned} \quad (3.3.25)$$

In particular, if we apply the ∞ -functor i^* , we get the following fiber-cofiber sequence:

$$\begin{aligned} &i^*\mathcal{M}_X^\vee(\mathbf{Sing}(X, s_X)) \\ &\quad \downarrow \\ &i^*\text{cofiber}(\mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) \xrightarrow{\mathcal{M}_X^\vee(t^*) \circ (1 - m_{\mathcal{L}_{X_0}^\vee})} i_*i^*\mathbf{BU}_X) \\ &\quad \downarrow \\ &i^*j_*j^*\mathbf{BU}_X. \end{aligned} \quad (3.3.26)$$

Proof The second statement is an immediate consequence of the first and of the equivalence $i^*j_! \simeq 0$. The first fiber-cofiber sequence can be obtained by applying the

octahedron axiom to the triangle

$$\begin{array}{ccc} \mathcal{M}_X^\vee(\mathbf{Perf}(X_0)) & \longrightarrow & i_* i^! \mathbf{BU}_X \\ & \searrow & \downarrow \\ \mathcal{M}_X^\vee(i^*) \circ \mathcal{M}_X^\vee(i_*) & \longrightarrow & i_* i^* \mathbf{BU}_X \end{array}$$

which appears in diagram (3.3.19) and by the fact that

$$\mathrm{cofib}(\alpha) \simeq \mathrm{cofib}(i_* i^! \mathbf{BU}_X \rightarrow i_* i^* \mathbf{BU}_X)$$

(see [44, Lemma 4.4.2.8]). □

Let $\mathcal{M}_{S, \mathbb{Q}}^\vee$ denote the composition

$$\mathbf{dgCat}_S^{\mathrm{idm}, \otimes} \rightarrow \mathbf{Mod}_{\mathbf{BU}_S}(\mathbf{SH}_S)^\otimes \xrightarrow{- \otimes \mathbf{H}\mathbb{Q}} \mathbf{Mod}_{\mathbf{BU}_{S, \mathbb{Q}}}(\mathbf{SH}_S)^\otimes. \quad (3.3.27)$$

We shall now study the commutative algebra object

$$\mathcal{M}_{S, \mathbb{Q}}^\vee(\mathbf{Sing}(S, 0)) \underset{S \text{ regular}}{\simeq} \mathcal{M}_{S, \mathbb{Q}}^\vee(\mathbf{Sing}(S_0)). \quad (3.3.28)$$

This is a particularly important object for our purposes as, for every $(X, s_X) \in \mathbf{LG}_{(S, \mathcal{L}_S)}$, the \mathbb{Q} -linear motivic realization of $\mathbf{Sing}(X, s_X)$ lies in the category of $\mathcal{M}_{S, \mathbb{Q}}^\vee(\mathbf{Sing}(S_0))$ -modules $\mathbf{Mod}_{\mathcal{M}_{S, \mathbb{Q}}^\vee(\mathbf{Sing}(S_0))}(\mathbf{SH}_S)$.

Proposition 3.3.9 *There is an equivalence in $\mathbf{CAlg}(\mathbf{SH}_S)$:*

$$\mathcal{M}_{S, \mathbb{Q}}^\vee(\mathbf{Sing}(S, 0)) \simeq \mathrm{cofiber}(\mathbf{BU}_{S, \mathbb{Q}} \xrightarrow{1 - m_{\mathcal{L}_S^\vee}} \mathbf{BU}_{S, \mathbb{Q}}) \quad (3.3.29)$$

Here

$$\mathrm{cofiber}(\mathbf{BU}_{S, \mathbb{Q}} \xrightarrow{1 - m_{\mathcal{L}_S^\vee}} \mathbf{BU}_{S, \mathbb{Q}}) := \mathbf{BU}_{S, \mathbb{Q}} \otimes_{\mathbf{H}\mathbb{Q}[t]} \mathbf{H}\mathbb{Q} \in \mathbf{CAlg}(\mathbf{SH}_S), \quad (3.3.30)$$

where we consider the morphisms $\mathbf{H}\mathbb{Q}[t] \rightarrow \mathbf{H}\mathbb{Q}$ induced by $t \mapsto 0$ and $\mathbf{H}\mathbb{Q}[t] \rightarrow \mathbf{BU}_{S, \mathbb{Q}}$ induced by $t \mapsto 1 - m_{\mathcal{L}_S^\vee}$.

Proof The underlying object of the commutative algebra $\mathbf{BU}_{S, \mathbb{Q}} \otimes_{\mathbf{H}\mathbb{Q}[t]} \mathbf{H}\mathbb{Q}$ is $\mathrm{cofiber}(\mathbf{BU}_{S, \mathbb{Q}} \xrightarrow{1 - m_{\mathcal{L}_S^\vee}} \mathbf{BU}_{S, \mathbb{Q}})$. This follows easily from the fact that

$$\mathbf{H}\mathbb{Q}[t] \xrightarrow{t} \mathbf{H}\mathbb{Q}[t] \quad (3.3.31)$$

is a free resolution of $H\mathbb{Q}$. Consider the diagram

$$\begin{array}{ccccc}
 S_0 & \xrightarrow{i} & S & \xleftarrow{j} & \emptyset \\
 \downarrow & & \downarrow i_0 & & \downarrow \\
 S & \xrightarrow{i_0} & \mathbb{V} := \mathbb{V}(\mathcal{L}_S) & \xleftarrow{j_0} & \mathcal{U} = \mathbb{V} - S,
 \end{array} \quad (3.3.32)$$

where both squares are (homotopy) cartesian. Notice that in this case $\pi_0(S_0) = S$ and therefore $i = i \circ t = id$. Then, the fiber-cofiber sequence of the previous corollary gives us an equivalence

$$\mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Sing}(S_0)) \simeq \text{cofiber}(\mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Perf}(S_0)) \xrightarrow{\mathcal{M}_{S,\mathbb{Q}}^{\vee}(t^*) \circ (1 - m_{\mathcal{L}_S}^{\vee})} \mathbf{BU}_{S,\mathbb{Q}})). \quad (3.3.33)$$

Since the square

$$\begin{array}{ccc}
 \mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Perf}(S_0)) & \xrightarrow{1 - m_{\mathcal{L}_{S_0}}^{\vee}} & \mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Perf}(S_0)) \\
 \downarrow \mathcal{M}_S^{\vee}(t^*) & & \downarrow \mathcal{M}_S^{\vee}(t^*) \\
 \mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Perf}(S)) & \xrightarrow{1 - m_{\mathcal{L}_S}^{\vee}} & \mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Perf}(S))
 \end{array} \quad (3.3.34)$$

is commutative up to coherent homotopy, to get the first desired equivalence

$$\begin{aligned}
 \mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Sing}(S, 0)) &\simeq \mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Sing}(S_0)) \\
 &\simeq \text{cofiber}(\mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Perf}(S_0)) \xrightarrow{\mathcal{M}_{S,\mathbb{Q}}^{\vee}(t^*) \circ (1 - m_{\mathcal{L}_S}^{\vee})} \mathbf{BU}_{S,\mathbb{Q}})) \quad (3.3.35)
 \end{aligned}$$

it suffices to show that $\mathcal{M}_{S,\mathbb{Q}}^{\vee}(t^*)$ is an equivalence. This is true as $i \circ t = id$ and $t \circ i$ is homotopic to the identity (see [9, Remark 3.31]). \square

Proposition 3.3.10 *There is an equivalence in $\mathbf{CAlg}(\mathbf{SH}_S)$*

$$\mathcal{M}_{S,\mathbb{Q}}^{\vee}(\mathbf{Sing}(S, 0)) \simeq i_0^* j_{0*} \mathbf{BU}_{\mathcal{U},\mathbb{Q}}. \quad (3.3.36)$$

Here $i_0 : S \rightarrow \mathbb{V}(\mathcal{L}_S)$ is the zero section, $j_0 : \mathcal{U} = \mathbb{V}(\mathcal{L}_S) - S \rightarrow \mathbb{V}(\mathcal{L}_S)$ the open complementary.

Proof Let $\mathbb{V} := \mathbb{V}(\mathcal{L}_S)$. Consider the equivalence that we get from the localization sequence of (i_0, j_0)

$$i_0^* j_{0*} \mathbf{BU}_{\mathcal{U},\mathbb{Q}} \simeq \text{cofiber}(i_0^* i_{0*} i_0^! \mathbf{BU}_{\mathbb{V},\mathbb{Q}} \simeq i_0^! \mathbf{BU}_{\mathbb{V},\mathbb{Q}} \xrightarrow{c} \mathbf{BU}_{S,\mathbb{Q}} \simeq i_0^* \mathbf{BU}_{\mathbb{V},\mathbb{Q}}), \quad (3.3.37)$$

where the map c is the one induced by the counit $i_{0*}i_0^! \mathbf{BU}_{\mathbb{V}, \mathbb{Q}} \rightarrow \mathbf{BU}_{\mathbb{V}, \mathbb{Q}}$. Notice that $i_0 : S \rightarrow V$ is a closed embedding between regular schemes. In particular, absolute purity holds. It follows from [15, Remark 13.5.5] that the composition

$$\mathbf{BU}_{\mathbb{V}, \mathbb{Q}} \rightarrow i_{0*}i_0^! \mathbf{BU}_{\mathbb{V}, \mathbb{Q}} \xrightarrow{\eta_{i_0}} \mathbf{BU}_{\mathbb{V}, \mathbb{Q}} \quad (3.3.38)$$

corresponds to $1 - m_{\mathcal{L}_S^\vee}$, as the conormal sheaf of $i_0 : S \rightarrow \mathbb{V}$ is \mathcal{L}_S^\vee . If we apply $i^!$ we obtain

$$i^!(1 - m_{\mathcal{L}_S^\vee}) : i^! \mathbf{BU}_{\mathbb{V}, \mathbb{Q}} \xrightarrow{c} \mathbf{BU}_{S, \mathbb{Q}} \xrightarrow[\simeq \text{abs. pur.}]{i^! \eta'_{i_0}} i^! \mathbf{BU}_{\mathbb{V}, \mathbb{Q}}, \quad (3.3.39)$$

which under the equivalence $\mathbf{BU}_{S, \mathbb{Q}} \simeq i^! \mathbf{BU}_{\mathbb{V}, \mathbb{Q}}$ corresponds to $1 - m_{\mathcal{L}_S^\vee}$. \square

Construction 3.3.11 Let $(X, s_X) \in \mathbf{LG}_{(S, \mathcal{L}_S)}$. Consider the morphism

$$\mathbf{BU}_X \xrightarrow{1 - m_{\mathcal{L}_X^\vee}} \mathbf{BU}_X \quad (3.3.40)$$

Since $j : X_{\mathcal{U}} \rightarrow X$ is the complementary open subscheme to the zero locus of the section s_X , it follows that $j^* \mathcal{L}_X^\vee \simeq \mathcal{O}_{X_{\mathcal{U}}}$. In particular,

$$j^* \circ (1 - m_{\mathcal{L}_X^\vee}) \simeq 1 - m_{\mathcal{L}_{X_{\mathcal{U}}}^\vee} \sim 0. \quad (3.3.41)$$

Then we obtain a morphism

$$sp_X^{\text{mot}} : \text{cofiber}(\mathbf{BU}_X \xrightarrow{1 - m_{\mathcal{L}_X^\vee}} \mathbf{BU}_X) \rightarrow j_* j^* \mathbf{BU}_X. \quad (3.3.42)$$

Proposition 3.3.12 Let $(X, s_X) \in \mathbf{LG}_{(S, \mathcal{L}_S)}$ and assume that X is regular and that $X \xrightarrow{s_X} \mathbb{V}(\mathcal{L}_S) \xleftarrow{\text{zero}} S$ is Tor-independent (i.e. $X \times_{\mathbb{V}(\mathcal{L}_S)} S \simeq X \times_{\mathbb{V}(\mathcal{L}_S)}^h S$). Then

$$i^* \mathcal{M}_X^\vee(\mathbf{Sing}(X, s_X)) \simeq \text{fiber}(i^* sp_X^{\text{mot}}). \quad (3.3.43)$$

Proof This follows immediately from the octahedron property applied to the triangle in the following diagram

$$\begin{array}{ccccc} \mathbf{BU}_{X_0} & \longrightarrow & i^! \mathbf{BU}_X & \longrightarrow & i^* \mathcal{M}_X^\vee(\mathbf{Sing}(X_0)) \\ & \searrow 1 - m_{\mathcal{L}_{X_0}^\vee} & \downarrow & & \\ & & \mathbf{BU}_{X_0} & & \\ & & \downarrow & & \\ & & i^* j_* j^* \mathbf{BU}_X & & \end{array} \quad (3.3.44)$$

and from the compatibility of $1 - m_{\mathcal{L}_X^\vee}$ with pullbacks. \square

4 Monodromy-invariant vanishing cycles

4.1 The formalism of vanishing cycles

We shall begin with a quick review of the formalism of vanishing cycles. We hope that this will be useful to understand the analogy which led to the definition of monodromy-invariant vanishing cycles.

Notation 4.1.1 Throughout this section, A will be an excellent henselian trait and S will denote the associated affine scheme. Label k its residue field and K its fraction field. Let σ be $\text{Spec}(k)$ and η be $\text{Spec}(K)$. Fix algebraic closures k^{alg} and K^{alg} of k and K respectively. We will consider:

- The maximal separable extension k^{sep} of k inside k^{alg} . We will use the notation $\bar{\sigma} = \text{Spec}(k^{sep})$.
- The maximal unramified extension K^{unr} of K in K^{alg} . We will use the notation $\eta^{unr} = \text{Spec}(K^{unr})$.
- The maximal tamely ramified extension K^t of K inside K^{alg} . We will use the notation $\eta^t = \text{Spec}(K^t)$.
- The maximal separable extension K^{sep} of K inside K^{alg} . We will use the notation $\bar{\eta} = \text{Spec}(K^{sep})$.

Moreover, we will fix an uniformizer π .

Remark 4.1.2 It is well known that there is an equivalence between the category of separable extensions of k and that of unramified extensions of K . In particular, $\text{Gal}(k^{sep}/k) \simeq \text{Gal}(K^{unr}/K)$. This, together with the fundamental theorem of Galois theory, implies that there is an exact sequence of groups

$$1 \rightarrow \text{Gal}(K^{sep}/K^{unr}) \rightarrow \text{Gal}(K^{sep}/K) \rightarrow \text{Gal}(k^{sep}/k) \rightarrow 1. \quad (4.1.1)$$

The Galois group on the left is called the inertia group and it is usually denoted by I . The chain of extensions $K^{unr} \subseteq K^t \subseteq K^{sep}$ gives us the following decomposition of I :

$$1 \rightarrow \underbrace{\text{Gal}(K^{sep}/K^t)}_{=: I_w} \rightarrow I \rightarrow \underbrace{\text{Gal}(K^{sep}/K)}_{=: I_t} \rightarrow 1. \quad (4.1.2)$$

The Galois group I_w is called wild inertia group, while I_t is called tame inertia group. See [51] for more details.

Remark 4.1.3 Let A^{sh} be a strict henselianisation of A and let $\bar{S} = \text{Spec}(A^{sh})$. We then have the following picture

$$\begin{array}{ccccccc}
 & & & & \bar{j} & & \\
 & & & & \curvearrowright & & \\
 \bar{\sigma} & \xrightarrow{\quad \bar{i} \quad} & \bar{S} & \xleftarrow{j_{unr}} & \eta^{unr} & \xleftarrow{\quad \eta^t \quad} & \bar{\eta} \\
 \downarrow v_\sigma & & \downarrow v & & \downarrow v_\eta & & \\
 \sigma & \xrightarrow{\quad i \quad} & S & \xleftarrow{\quad j \quad} & \eta & &
 \end{array} \quad (4.1.3)$$

where both squares are cartesian. In particular, notice that (σ, η) and $(\bar{\sigma}, \eta^{unr})$ form closed-open coverings of S and \bar{S} respectively.

Let $p : X \rightarrow S$ be of finite type and let $\bar{X} = X \times_S \bar{S}$. Consider the following diagram, cartesian over the base (this is also called Grothendieck's trick in the literature)

$$\begin{array}{ccccccc}
 X_{\bar{\sigma}} & \hookrightarrow & \bar{X} & \xleftarrow{\quad} & X_{\eta^{unr}} & \xleftarrow{\quad} & X_{\eta^t} \xleftarrow{\quad} X_{\bar{\eta}} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \bar{\sigma} & \hookrightarrow & \bar{S} & \xleftarrow{\quad} & \eta^{unr} & \xleftarrow{\quad} & \eta^t \xleftarrow{\quad} \bar{\eta}.
 \end{array} \quad (4.1.4)$$

Remark 4.1.4 The ∞ -category $\mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$ is the recollement of $\mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\sigma}})$ and $\mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\eta^{unr}})$ in the sense of [34, Appendix A.8].⁷ Indeed, $\mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$ is a stable ∞ -category and therefore it admits finite limits. Moreover, both $\bar{i}_* : \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\sigma}}) \rightarrow \mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$ and $j_*^{unr} : \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\eta^{unr}}) \rightarrow \mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$ are fully faithful ∞ -functors: these are immediate consequences of the proper and smooth base change theorems applied to the cartesian squares

$$\begin{array}{ccc}
 X_{\bar{\sigma}} & \xrightarrow{id} & X_{\bar{\sigma}} \\
 \downarrow id & & \downarrow \bar{i} \\
 X_{\bar{\sigma}} & \xrightarrow{\quad \bar{i} \quad} & \bar{X}
 \end{array}
 \quad
 \begin{array}{ccc}
 X_{\bar{\sigma}} & \xrightarrow{id} & X_{\bar{\sigma}} \\
 \downarrow id & & \downarrow j_{\eta^{unr}} \\
 X_{\bar{\sigma}} & \xrightarrow{j_{\eta^{unr}}} & \bar{X}.
 \end{array} \quad (4.1.5)$$

It is known that the conditions of [34, Definition A.8.1] are verified. In particular, using the equivalence of $\mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\eta^{unr}})$ with the ∞ -category of ℓ -adic sheaves on the generic geometric fiber endowed with a continuous action of $I \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\eta}})^I$, we have that $\mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$ is the recollement of $\mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\sigma}})$ and $\mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\eta}})^I$.

⁷ Strictly speaking, one should consider the full subcategories of $\mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$ spanned by the images of \bar{i}_* and $j_{\eta^{unr}}^*$ which are closed under equivalences.

Remark 4.1.5 Notice that we have the following diagram where each arrow is an equivalence:

$$\begin{array}{ccc}
 & \bar{j}_*(-)^{\mathrm{h}I} & \\
 & \downarrow & \\
 \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\eta^{\mathrm{unr}}}) & \xrightarrow{\bar{j}^*} & \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\eta}})^I \\
 \uparrow \scriptstyle j_{I*}(-)^{\mathrm{h}I_I} \quad \searrow \scriptstyle j_I^* & & \nearrow \scriptstyle j_w^* \quad \downarrow \scriptstyle j_{w*}(-)^{\mathrm{h}I_w} \\
 & \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\eta^I})^{I_I} &
 \end{array} \tag{4.1.6}$$

Construction 4.1.6 We will now sketch the construction of Deligne's topos in the ∞ -categorical world. Notice that, by the recollement technique, the étale ∞ -topos of \bar{S} is the recollement of the étale ∞ -topos of $\bar{\sigma}$ and that of η^{unr} . Moreover, notice that the étale ∞ -topos of η^{unr} is equivalent to the ∞ -category of spaces with a continuous action of I . For a scheme Y over $\bar{\sigma}$, we consider the lax ∞ -limit $\tilde{Y}_{et} \times_{\tilde{\bar{\sigma}}_{et}} \tilde{\bar{S}}_{et}$, which exists by [35, pag. 614]. The decomposition of $\tilde{\bar{S}}_{et}$ gives us a digram

$$\begin{array}{ccccc}
 \tilde{Y}_{et} & \longrightarrow & \tilde{Y}_{et} \times_{\tilde{\bar{\sigma}}_{et}} \tilde{\bar{S}}_{et} & \longleftarrow & \tilde{Y}_{et} \times_{\tilde{\bar{\sigma}}_{et}} \eta^{\mathrm{unr}}_{et} \\
 \downarrow & & \downarrow & & \downarrow \\
 \tilde{\bar{\sigma}}_{et} & \longrightarrow & \tilde{\bar{S}}_{et} & \longleftarrow & \eta^{\mathrm{unr}}_{et}
 \end{array} \tag{4.1.7}$$

We will label the ∞ -category of ℓ -adic sheaves on $\tilde{Y}_{et} \times_{\tilde{\bar{\sigma}}_{et}} \eta^{\mathrm{unr}}_{et}$ by $\mathbf{Shv}_{\mathbb{Q}_\ell}(Y)^I$, as it is the ∞ -category of ℓ -adic sheaves on Y endowed with a continuous action of I . The ∞ -category $\mathbf{Shv}_{\mathbb{Q}_\ell}(Y)^{I_I}$ of ℓ -adic sheaves on Y endowed with a continuous action of I_I identifies with the full subcategory of $\mathbf{Shv}_{\mathbb{Q}_\ell}(Y)^I$ such that the induced action of I_w is trivial.

Definition 4.1.7 Let $p : \bar{X} \rightarrow \bar{S}$ be an \bar{S} -scheme. Let $\mathcal{E} \in \mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X}_{\eta^{\mathrm{unr}}})$. The ℓ -adic sheaf of tame nearby cycles of \mathcal{E} is defined as

$$\Psi_I(\mathcal{E}) := \bar{i}^* j_{I*} \mathcal{E}|_{X_{\eta_I}} \in \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\sigma}})^{I_I}. \tag{4.1.8}$$

Analogously, the ℓ -adic sheaf of nearby cycles of \mathcal{E} is defined as

$$\Psi(\mathcal{E}) := \bar{i}^* \bar{j}_* \mathcal{E}|_{X_{\bar{\eta}}} \in \mathbf{Shv}_{\mathbb{Q}_\ell}(X_{\bar{\sigma}})^I. \tag{4.1.9}$$

Remark 4.1.8 With the same notation as above, the following equivalence holds in $\mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X}_{\bar{\sigma}})^{I_I}$:

$$\Psi_I(\mathcal{E}) \simeq \Psi(\mathcal{E})^{\mathrm{h}I_w}. \tag{4.1.10}$$

Definition 4.1.9 Let $p : \bar{X} \rightarrow \bar{S}$ be an \bar{S} -scheme. Let $\mathcal{F} \in \mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X})$. The unit of the adjunction (j_i^*, j_{i*}) induces a morphism in $\mathbf{Shv}_{\mathbb{Q}_\ell}(\bar{X}_{\bar{\sigma}})^{I_t}$

$$\bar{i}^* \mathcal{F} \rightarrow \Psi_t(\mathcal{F}_{\bar{X}_{\eta_{unr}}}), \quad (4.1.11)$$

where we regard the object on the left endowed with the trivial I_t -action. The cofiber of this morphism is by definition the ℓ -adic sheaf of tame vanishing cycles of \mathcal{F} , which we will denote by $\Phi_t(\mathcal{F})$. In a similar way, we define the ℓ -adic sheaf of vanishing cycles $\Phi(\mathcal{F})$ as the cofiber of the morphism (called the specialization morphism in literature)

$$\bar{i}^* \mathcal{F} \rightarrow \Psi(\mathcal{F}_{\bar{X}_{\bar{\eta}}}) \quad (4.1.12)$$

induced by the unit of the adjunction (\bar{j}^*, \bar{j}_*) .

4.2 Monodromy-invariant vanishing cycles

Context 4.2.1 Assume that S is strictly henselian, i.e. that k is a separably closed field and that $p : X \rightarrow S$ is a proper morphism.

For our purposes, we are interested in the image of $\Phi(\mathbb{Q}_{\ell, X})$ via the ∞ -functor

$$(-)^{hI} : \mathbf{Shv}_{\mathbb{Q}_\ell}(X_\sigma)^I \rightarrow \mathbf{Shv}_{\mathbb{Q}_\ell}(X_\sigma).$$

It is then important to remark that it is possible to determine $p_{\sigma*} \Phi(\mathbb{Q}_{\ell, X})^{hI}$ without ever mentioning this ∞ -functor. Notice that $\Phi(\mathbb{Q}_{\ell, X})^{hI}$ is the cofiber of the image of the specialization morphism via $(-)^{hI}$:

$$\mathbb{Q}_{\ell, X_0}^{hI} \xrightarrow{(sp)^{hI}} \Psi(\mathbb{Q}_{\ell, X_{\bar{\eta}}})^{hI} \rightarrow \Phi(\mathbb{Q}_{\ell, X})^{hI} \quad (4.2.1)$$

is a fiber-cofiber sequence in $\mathbf{Shv}_{\mathbb{Q}_\ell}(X_\sigma)$.

Remark 4.2.2 There are equivalences in $\mathbf{CAlg}(\mathbf{Shv}_{\mathbb{Q}_\ell}(\sigma))$

- $(p_{\sigma*} \mathbb{Q}_{\ell, X_\sigma})^{hI} \simeq p_{\sigma*} \mathbb{Q}_{\ell, X_\sigma} \otimes_{\mathbb{Q}_{\ell, \sigma}} \mathbb{Q}_{\ell, \sigma}^{hI}$ (see [9, Proposition 4.32]),
- $(p_{\sigma*} i^* \bar{j}_* \mathbb{Q}_{\ell, X_{\bar{\eta}}})^{hI} \simeq p_{\sigma*} i^* j_* \mathbb{Q}_{\ell, X_\eta}$ (see [9, Proposition 4.31]),
- $\mathbb{Q}_{\ell, \sigma}^{hI} \simeq i_0^* j_{0*} \mathbb{Q}_{\ell, \eta}$ (see [9, Lemma 4.34]).

Consider the object $p_{\sigma*} p_\sigma^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta}$ in $\mathbf{Shv}_{\mathbb{Q}_\ell}(\sigma)$. The following chain of equivalences, together with the previous remark, identify it with $(p_{\sigma*} \mathbb{Q}_{\ell, X_\sigma})^{hI}$:

$$\begin{aligned}
& p_{\sigma*} p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} \\
& \simeq p_{\sigma*} (\mathbb{Q}_{\ell, X_{\sigma}} \otimes_{\mathbb{Q}_{\ell, X_{\sigma}}} p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta}) \underset{\text{proj. form.}}{\simeq} p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \otimes_{\mathbb{Q}_{\ell, \sigma}} i_0^* j_{0*} \mathbb{Q}_{\ell, \eta}.
\end{aligned} \tag{4.2.2}$$

Definition 4.2.3 Consider the canonical morphism in $\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X_{\sigma})$

$$sp_p^{\text{hm}} : p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} \simeq i^* p^* j_{0*} \mathbb{Q}_{\ell, \eta} \rightarrow i^* j_* p_{\eta}^* \mathbb{Q}_{\ell, \eta} \simeq i^* j_* \mathbb{Q}_{\ell, X_{\eta}} \tag{4.2.3}$$

induced by the base change natural transformation $p^* j_{0*} \rightarrow j_* p_{\eta}^*$. We will refer to it as the monodromy-invariant specialization morphism. We will refer to the cofiber of this morphism in $\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X_{\sigma})$ as monodromy-invariant vanishing cycles, which we will denote $\Phi_p^{\text{hm}}(\mathbb{Q}_{\ell})$.

In order to justify the choice of the name of this morphism, we shall prove the following:

Lemma 4.2.4 *The two arrows $p_{\sigma*}(sp)^{\text{hl}}$ and $p_{\sigma*}sp_p^{\text{hm}}$ are homotopic under the equivalences*

$$\begin{aligned}
& (p_{\sigma*} i^* \bar{j}_* \mathbb{Q}_{\ell, X_{\eta}})^{\text{hl}} \simeq p_{\sigma*} i^* j_* \mathbb{Q}_{\ell, X_{\eta}}, \\
& (p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}})^{\text{hl}} \simeq p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \otimes_{\mathbb{Q}_{\ell, \sigma}} \mathbb{Q}_{\ell, \sigma}^{\text{hl}} \simeq p_{\sigma*} p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta}.
\end{aligned}$$

Proof We shall view both arrows as maps $p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \otimes_{\mathbb{Q}_{\ell, \sigma}} \mathbb{Q}_{\ell, \sigma}^{\text{hl}} \rightarrow p_{\sigma*} i^* j_* \mathbb{Q}_{\ell, X_{\eta}}$.

Since the tensor product defines a cocartesian symmetric monoidal structure on $\text{CAlg}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(\sigma))$ (see [34]), it suffices to show that the maps of commutative algebras obtained from $p_{\sigma*}(sp)^{\text{hl}}$ and $p_{\sigma*}sp_p^{\text{hm}}$ by precomposition with $p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \rightarrow p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \otimes_{\mathbb{Q}_{\ell, \sigma}} \mathbb{Q}_{\ell, \sigma}^{\text{hl}}$ and $\mathbb{Q}_{\ell, \sigma}^{\text{hl}} \rightarrow p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \otimes_{\mathbb{Q}_{\ell, \sigma}} \mathbb{Q}_{\ell, \sigma}^{\text{hl}}$ respectively are equivalent. It is essentially the proof of the main theorem in [9] that the precompositions of $p_{\sigma*}(sp)^{\text{hl}}$ with these two maps are induced by the canonical morphisms $1 \rightarrow j_* j^*$ and $1 \rightarrow p_{\eta*} p_{\eta}^*$. We are left to show that the same holds true if we consider sp_p^{hm} instead of $(sp)^{\text{hl}}$. Recall that sp_p^{hm} is induced by the base change morphism $p^* j_{0*} \rightarrow j_* p_{\eta}^*$, i.e. the morphism which corresponds under adjunction to $j_{0*} \rightarrow j_{0*} p_{\eta*} p_{\eta}^*$, induced by the unit of the adjunction $(p_{\eta}^*, p_{\eta*})$. From this we obtain the following diagram

$$\begin{array}{ccc}
i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} & \xrightarrow{1 \rightarrow p_{\eta*} p_{\eta}^*} & i_0^* j_{0*} p_{\eta*} p_{\eta}^* \mathbb{Q}_{\ell, \eta} \simeq p_{\sigma*} i^* j_* p_{\eta}^* \mathbb{Q}_{\ell, \eta} \\
\downarrow 1 \rightarrow p_{\sigma*} p_{\sigma}^* & & \downarrow \\
p_{\sigma*} p_{\sigma}^* \mathbb{Q}_{\ell, \sigma} & \xrightarrow{1 \rightarrow j_0^* j_{0*}} & p_{\sigma*} p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} \xrightarrow{\quad} p_{\sigma*} p_{\sigma}^* p_{\sigma*} i^* j_* p_{\eta}^* \mathbb{Q}_{\ell, \eta} \xrightarrow{id} \\
& \searrow p_{\sigma*} sp_p^{\text{hm}} & \downarrow \\
& & p_{\sigma*} i^* j_* p_{\eta}^* \mathbb{Q}_{\ell, \eta}.
\end{array} \tag{4.2.4}$$

In particular the composition $i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} \rightarrow p_{\sigma*} p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} \xrightarrow{p_{\sigma*} s p_p^{\text{hm}}} p_{\sigma*} i^* j_* \mathbb{Q}_{\ell, X_{\eta}}$ is homotopic to the map induced by $1 \rightarrow p_{\eta*} p_{\eta}^*$.

The composition $p_{\sigma*} p_{\sigma}^* \mathbb{Q}_{\ell, \sigma} \rightarrow p_{\sigma*} p_{\sigma}^* i_0^* j_{0*} \mathbb{Q}_{\ell, \eta} \rightarrow p_{\sigma*} i^* j_* \mathbb{Q}_{\ell, X_{\eta}}$ is homotopic to the map $p_{\sigma*} \mathbb{Q}_{\ell, X_{\sigma}} \rightarrow p_{\sigma*} i^* j_* \mathbb{Q}_{\ell, X_{\eta}}$ induced by $1 \rightarrow j_* j^*$ by the following commutative triangle

$$\begin{array}{ccc}
 1 \rightarrow j_{0*} j_0^* & \xrightarrow{\quad} & p^* j_{0*} j_0^* \mathbb{Q}_{\ell, S} \\
 p^* \mathbb{Q}_{\ell, S} \searrow & & \downarrow \\
 1 \rightarrow j_* j^* & \xrightarrow{\quad} & j_* p_{\eta}^* j_0^* \mathbb{Q}_{\ell, S} \simeq j_* j^* p^* \mathbb{Q}_{\ell, S}.
 \end{array} \tag{4.2.5}$$

□

Remark 4.2.5 Notice that what we said above is actually true for any ℓ -adic sheaf in the image of $p^* : \mathbf{Shv}_{\mathbb{Q}_{\ell}}(S) \rightarrow \mathbf{Shv}_{\mathbb{Q}_{\ell}}(X)$.

The advantage of this reformulation is that it allows to define inertia invariant vanishing cycles without ever mention the inertia group. We will adopt it for the situation we are interested in.

Assume that S is a noetherian regular scheme.

Definition 4.2.6 Let (X, s_X) be a twisted LG model over (S, \mathcal{L}_S) . Consider the diagram obtained from the zero section of the vector bundle associated to \mathcal{L}_X :

$$\begin{array}{ccccc}
 \pi_0(X_0) & \xrightarrow{i} & X & \xleftarrow{j} & X_{\mathcal{U}} \\
 \downarrow s_0 & & \downarrow s_X & & \downarrow s_{\mathcal{U}} \\
 X & \xrightarrow{i_0} & \mathbb{V}(\mathcal{L}_X) & \xleftarrow{j_0} & \mathcal{U}_X := \mathbb{V}(\mathcal{L}_X) - X.
 \end{array} \tag{4.2.6}$$

Define the monodromy-invariant specialization morphism associated to (X, s_X) as the map in $\mathbf{Shv}_{\mathbb{Q}_{\ell}}(\pi_0(X_0))$

$$sp_{(X, s_X)}^{\text{mi}} : s_0^* i_0^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \simeq i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \rightarrow i^* j_* \mathbb{Q}_{\ell, X_{\mathcal{U}}} \simeq i^* j_* s_{\mathcal{U}}^* \mathbb{Q}_{\ell, \mathcal{U}_X} \tag{4.2.7}$$

induced by the natural transformation $s_X^* j_{0*} \rightarrow j_* s_{\mathcal{U}}^*$. We will refer to the cofiber of $sp_{(X, s_X)}^{\text{mi}}$ as monodromy invariant vanishing cycles of (X, s_X) , that we will denote $\Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell})$.

Proposition 4.2.7 Let (X, s_X) be as above and assume X regular. There is an equivalence

$$i_0^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \simeq \text{cofib}(c_1(\mathcal{L}_X) : \mathbb{Q}_{\ell, X}(-1)[-2] \rightarrow \mathbb{Q}_{\ell, X}). \tag{4.2.8}$$

Proof Consider the diagram

$$X \xrightarrow{i_0} \mathbb{V}_X := \mathbb{V}(\mathcal{L}_X) \xleftarrow{j_0} \mathcal{U}.$$

The ℓ -adic sheaf $i_0^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}}$ is the cofiber of the morphism $i_0^* (i_{0*} i_0^! \mathbb{Q}_{\ell, \mathbb{V}_X} \xrightarrow{\text{count}} \mathbb{Q}_{\ell, \mathbb{V}_X})$ in $\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X)$. Recall that $i_0^* i_{0*} \simeq id$. Therefore, we can consider the following triangle

$$\begin{array}{ccc} i_0^! \mathbb{Q}_{\ell, \mathbb{V}_X} & \longrightarrow & \mathbb{Q}_{\ell, X} \\ \text{abs. pur.} \swarrow \simeq & & \nearrow \\ \mathbb{Q}_{\ell, X}(-1)[-2] & & \end{array} \quad (4.2.9)$$

The absolute purity isomorphism is given by the class $cl(X) \in H_X^2(\mathbb{V}_X, \mathbb{Q}_{\ell}(1))$, whose image in $H^2(X, \mathbb{Q}_{\ell}(1))$ is $c_1(\mathcal{N}_{X|\mathbb{V}_X}^{\vee})$, the first Chern class of the conormal bundle (see [25]). The map $\mathbb{Q}_{\ell, X} \rightarrow \mathbb{Q}_{\ell, X}(1)[2]$ corresponding to this class is the image of the right vertical arrow in the diagram above via the ∞ -functor $-(1)[2]$. It suffices to notice that $\mathcal{N}_{X|\mathbb{V}_X}^{\vee} \simeq \mathcal{L}_X$ to conclude. \square

5 The comparison theorem

5.1 2-periodic ℓ -adic sheaves

We shall now approach the comparison between monodromy-invariant vanishing cycles and the ℓ -adic realization of the dg category of singularity $\mathbf{Sing}(X_0)$. In order to do so, we need to work with the category of $\mathbb{Q}_{\ell, S}(\beta)$ -modules in $\mathbf{Shv}_{\mathbb{Q}_{\ell}}(S)$. Recall that there is an adjunction of ∞ -functors

$$\mathbf{Shv}_{\mathbb{Q}_{\ell}}(S) \rightleftarrows \mathbf{Mod}_{\mathbb{Q}_{\ell, S}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(S)) \quad (5.1.1)$$

given by $- \otimes_{\mathbb{Q}_{\ell, S}} \mathbb{Q}_{\ell, S}(\beta)$ and by the forgetful functor.

Notice that $\mathbf{Mod}_{\mathbb{Q}_{\ell, \bullet}}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(\bullet))$ defines a fibered category over the category of schemes, which satisfies Grothendieck's six functors formalism.

Definition 5.1.1 With the same notation as above let

$$s_0^* i_0^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \simeq i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \rightarrow i^* j_* s_{\mathcal{U}}^* \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \quad (5.1.2)$$

be the arrow in $\mathbf{Mod}_{\mathbb{Q}_{\ell, X}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X))$ induced by the base change morphism $s_X^* j_{0*} \rightarrow j_* s_{\mathcal{U}}^*$. Denote by $\Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}(\beta))$ its cofiber.

Proposition 5.1.2 *The maps (5.1.2) and $sp_{(X, s_X)}^{\text{mi}} \otimes_{\mathbb{Q}_{\ell, \pi(X_0)}} \mathbb{Q}_{\ell, \pi(X_0)}(\beta)$ (see Definition 4.2.7) are homotopic. In particular, there is an equivalence*

$$\Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}) \otimes_{\mathbb{Q}_{\ell, \pi(X_0)}} \mathbb{Q}_{\ell, \pi(X_0)}(\beta) \simeq \Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell, X}(\beta)). \quad (5.1.3)$$

in $\mathbf{Mod}_{\mathbb{Q}_{\ell, \pi(X_0)}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(\pi(X_0)))$.

Proof This is analogous to [9, Proposition 4.28]. As a first step, notice that both $*$ -pullbacks and $*$ -pushforwards commute with Tate and usual shifts. Since $-\otimes_{\mathbb{Q}_{\ell}} \mathbb{Q}_{\ell}(\beta)$ commutes with $*$ -pullbacks, we have

$$i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \otimes_{\mathbb{Q}_{\ell, \pi(X_0)}} \mathbb{Q}_{\ell, \pi(X_0)}(\beta) \simeq i^* s_X^* (j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \otimes_{\mathbb{Q}_{\ell, \mathbb{V}_X}} \mathbb{Q}_{\ell, \mathbb{V}_X}(\beta)), \quad (5.1.4)$$

$$i^* j_* \mathbb{Q}_{\ell, X_{\mathcal{U}}} \otimes_{\mathbb{Q}_{\ell, \pi_0(X)}} \mathbb{Q}_{\ell, \pi_0(X)}(\beta) \simeq i^* (j_* \mathbb{Q}_{\ell, X_{\mathcal{U}}} \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta)). \quad (5.1.5)$$

Using the equivalence $\mathbb{Q}_{\ell, \mathbb{V}_X}(\beta) \simeq \bigoplus_{i \in \mathbb{Z}} \mathbb{Q}_{\ell, \mathbb{V}_X}(i)[2i]$, we see that

$$j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \otimes_{\mathbb{Q}_{\ell, \mathbb{V}_X}} \mathbb{Q}_{\ell, \mathbb{V}_X}(\beta) \simeq \bigoplus_{i \in \mathbb{Z}} (j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(i)[2i]). \quad (5.1.6)$$

We shall show that the canonical map

$$\bigoplus_{i \in \mathbb{Z}} (j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(i)[2i]) \rightarrow j_{0*} \left(\bigoplus_{i \in \mathbb{Z}} \mathbb{Q}_{\ell, \mathcal{U}_X}(i)[2i] \right) \quad (5.1.7)$$

is an equivalence. This follows immediately from the fact that the $*$ -pushforward commutes with filtered colimits and from the equivalence $\bigoplus_{i \in \mathbb{Z}} \mathbb{Q}_{\ell, \mathcal{U}_X}(i)[2i] \simeq \text{colim}_{i \geq 0} \bigoplus_{k=-i}^i \mathbb{Q}_{\ell, \mathcal{U}_X}(k)[2k]$. This shows that

$$i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X} \otimes_{\mathbb{Q}_{\ell, \pi(X_0)}} \mathbb{Q}_{\ell, \pi(X_0)}(\beta) \simeq i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta). \quad (5.1.8)$$

The same argument applies to show that

$$i^* j_* \mathbb{Q}_{\ell, X_{\mathcal{U}}} \otimes_{\mathbb{Q}_{\ell, \pi_0(X)}} \mathbb{Q}_{\ell, \pi_0(X)}(\beta) \simeq i^* j_* \mathbb{Q}_{\ell, X_{\mathcal{U}}}(\beta). \quad (5.1.9)$$

To show that the two maps are homotopic, it suffices to show that they are so before applying i^* . Notice that $(\mathbb{Q}_{\ell, \mathcal{U}_X} \rightarrow s_{\mathcal{U}*} s_{\mathcal{U}}^* \mathbb{Q}_{\ell, \mathcal{U}_X}) \otimes_{\mathbb{Q}_{\ell, \mathcal{U}_X}} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)$ is homotopic to $\mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \rightarrow s_{\mathcal{U}*} s_{\mathcal{U}}^* \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)$, as $-\otimes_{\mathbb{Q}_{\ell, \mathcal{U}_X}} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)$ is compatible with the $*$ -pullback and with the $!$ -pushforward, which coincides with the $*$ -pushforward for $s_{\mathcal{U}}$ as it is a closed morphism. For what we have said above,

$$\begin{aligned} j_{0*} ((\mathbb{Q}_{\ell, \mathcal{U}_X} \rightarrow s_{\mathcal{U}*} s_{\mathcal{U}}^* \mathbb{Q}_{\ell, \mathcal{U}_X}) \\ \otimes_{\mathbb{Q}_{\ell, \mathcal{U}_X}} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)) \simeq j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \rightarrow j_{0*} s_{\mathcal{U}*} s_{\mathcal{U}}^* \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \end{aligned} \quad (5.1.10)$$

and $-\otimes_{\mathbb{Q}_{\ell}} \mathbb{Q}_{\ell}(\beta)$ is compatible with $*$ -pullbacks. The last assertion follows as $-\otimes_{\mathbb{Q}_{\ell}} \mathbb{Q}_{\ell}(\beta)$ is exact. \square

Corollary 5.1.3 *Let $(X, s_X) \in \mathbf{LG}_{(S, \mathcal{L}_S)}$ and assume that X is a regular scheme. The following equivalence holds in $\mathbf{Mod}_{\mathbb{Q}_{\ell, X}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X))$*

$$i_0^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta) \simeq \text{cofiber}(c_1(\mathcal{L}_X) \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta) : \mathbb{Q}_{\ell, X}(\beta) \rightarrow \mathbb{Q}_{\ell, X}(\beta)). \quad (5.1.11)$$

Proof This follows immediately from the previous Proposition, Proposition 4.2.7 and from algebraic Bott periodicity $\mathbb{Q}_{\ell, X}(\beta)(-1)[-2] \simeq \mathbb{Q}_{\ell, X}(\beta)$. \square

Remark 5.1.4 Notice that $i_0^* j_{0*}$ and $i^* j_*$ are lax monoidal functors. In particular, $s_0^* i_0^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)$ and $i^* j_* \mathbb{Q}_{\ell, X_{\mathcal{U}}}(\beta)$ have commutative algebra structures. The map $sp_{(X, s_X)}^{\text{mi}} \otimes_{\mathbb{Q}_{\ell, X_0}} \mathbb{Q}_{\ell, X_0}(\beta)$ is a map of commutative algebras as $s_X^* j_{0*} \mathbb{Q}_{\ell, X_{\mathcal{U}_X}}(\beta) \rightarrow j_* s_{\mathcal{U}}^* \mathbb{Q}_{\ell, X_{\mathcal{U}_X}}(\beta)$ is so. In particular, the ℓ -adic sheaf $\Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}(\beta))$ lives in $\mathbf{Mod}_{i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X_0))$.

5.2 The main theorem

Proposition 5.2.1 *Let X be a regular scheme and let $\mathcal{L} \in \text{Pic}(X)$. Then*

$$\begin{aligned} \text{cofib}(1 - \mathcal{R}_X^{\ell}(m_{\mathcal{L}}) : \mathcal{R}_X^{\ell}(\text{BU}_X) \rightarrow \mathcal{R}_X^{\ell}(\text{BU}_X)) \\ \simeq \text{cofib}(c_1(\mathcal{L}) \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta) : \mathbb{Q}_{\ell, X}(\beta) \rightarrow \mathbb{Q}_{\ell, X}(\beta)), \end{aligned} \quad (5.2.1)$$

where $c_1(\mathcal{L})$ is the first Chern class of \mathcal{L} .

Moreover,

$$\text{cofib}(c_1(\mathcal{L}) \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta)) \simeq \text{cofib}(c_1(\mathcal{L}^{\vee}) \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta)). \quad (5.2.2)$$

Proof We start by noticing that the composition

$$\text{HK}(X) \otimes_{\mathbb{Z}} \mathbb{Q} \simeq H_{\mathcal{M}}^{2*,*}(X, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q} \simeq H_{\mathcal{M}}^{2*,*}(X, \mathbb{Q}), \quad (5.2.3)$$

where $H_{\mathcal{M}}^{2*,*}$ denotes motivic cohomology, coincides with the Chern character (see [15, §11.3.6]) which is compatible with the ℓ -adic Chern character (see [9, §3.6 - §3.7] and [15]). Now, according to [15, Proposition 12.2.9] and [20, Proposition 3.8], $c_1(\mathcal{L})$ is nilpotent and therefore the Chern character of \mathcal{L} can be written as $1 + c_1(\mathcal{L}) + \frac{1}{2}c_1(\mathcal{L})^2 + \dots + \frac{1}{m!}c_1(\mathcal{L})^m$ for some $m \geq 1$. Then

$$1 - \mathcal{R}_X^{\ell}(m_{\mathcal{L}}) = -c_1(\mathcal{L})(1 + \frac{1}{2}c_1(\mathcal{L}) + \dots + \frac{1}{m!}c_1(\mathcal{L})^{m-1}).$$

As $c_1(\mathcal{L})$ is nilpotent, so is $\frac{1}{2}c_1(\mathcal{L}) + \dots + \frac{1}{m!}c_1(\mathcal{L})^{m-1}$, whence $(1 + \frac{1}{2}c_1(\mathcal{L}) + \dots + \frac{1}{m!}c_1(\mathcal{L})^{m-1})$ is invertible. This shows that

$$\begin{aligned} \text{cofiber}(1 - \mathcal{R}_X^{\ell}(m_{\mathcal{L}})) &\simeq \text{cofiber}(-c_1(\mathcal{L}) \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta)) \\ &\simeq \text{cofiber}(c_1(\mathcal{L}) \otimes_{\mathbb{Q}_{\ell, X}} \mathbb{Q}_{\ell, X}(\beta)). \end{aligned}$$

The last equivalence follows from the fact that, by [15, Proposition 12.2.9 and Remark 13.2.2], we have

$$0 = c_1(\mathcal{L} \otimes \mathcal{L}^\vee) \simeq F(c_1(\mathcal{L}), c_1(\mathcal{L}^\vee)) = c_1(\mathcal{L}) + c_1(\mathcal{L}^\vee) + \beta^{-1} \cdot c_1(\mathcal{L}) \cdot c_1(\mathcal{L}^\vee). \quad (5.2.4)$$

In particular,

$$c_1(\mathcal{L}^\vee) \simeq -c_1(\mathcal{L})(1 + \beta^{-1} \cdot c_1(\mathcal{L}^\vee)) \quad (5.2.5)$$

and, as we have remarked above, $-(1 + \beta^{-1} \cdot c_1(\mathcal{L}^\vee))$ is invertible as $c_1(\mathcal{L}^\vee)$ is nilpotent. \square

We are finally ready to state our main theorem:

Theorem 5.2.2 *Let (X, s_X) be a twisted LG model over (S, \mathcal{L}_S) . Assume that X is regular and that $0, s_X : X \rightarrow \mathbb{V}(\mathcal{L}_X)$ are Tor-independent. The following equivalence holds in $\mathbf{Mod}_{i^*s_X^*j_{0*}\mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_\ell}(X_0))$:*

$$i^*\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, s_X)) \simeq \Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_\ell(\beta))[-1], \quad (5.2.6)$$

where the $i^*s_X^*j_{0*}\mathbb{Q}_{\ell, \mathcal{U}_X}(\beta)$ -module structure on the l.h.s. is the one induced by the equivalence of commutative algebras

$$i^*\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0)) \simeq i^*s_X^*j_{0*}\mathbb{Q}_{\ell, \mathcal{U}_X}(\beta). \quad (5.2.7)$$

Proof We start by noticing that $i = s_0 : X_0 \rightarrow X$. Indeed, $i_0 \circ s_0 = s_X \circ i$ and both i_0 and s_X are sections of the canonical morphism $\mathbb{V}_X = \mathbb{V}(\mathcal{L}_X) \rightarrow X$. In particular

$$i^*\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0)) \simeq s_0^*\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0)) \underset{\text{Proposition 3.3.10}}{\simeq} i^*s_X^*j_{0*}\mathbb{Q}_{\ell, \mathcal{U}_X}(\beta). \quad (5.2.8)$$

Since X is supposed to be regular, $\mathbf{Sing}(X, s_X) \simeq \mathbf{Sing}(X_0)$. By Propositions 3.3.8 and 5.2.1 we have a fiber-cofiber sequence

$$\begin{array}{ccc} i^*\mathcal{R}_X^{\ell}(\mathbf{Sing}(X_0)) & & \\ \downarrow & & \\ \text{cofiber}(c_1(\mathcal{L}_{X_0}) \otimes_{\mathbb{Q}_{\ell, X_0}} \mathbb{Q}_{\ell, X_0}(\beta) : \mathbb{Q}_{\ell, X_0}(\beta) \rightarrow \mathbb{Q}_{\ell, X_0}(\beta)) & & (5.2.9) \\ \downarrow f & & \\ i^*j_*\mathbb{Q}_{\ell, X_{\mathcal{U}}}(\beta). & & \end{array}$$

in $\mathbf{Mod}_{i^*s_X^*j_{0*}\mathbb{Q}_{\ell,\mathcal{U}_X}(\beta)}(\mathbf{Shv}_{\mathbb{Q}_{\ell}}(X_0))$. On the other hand, by Proposition 5.1.2, we have another fiber-cofiber sequence

$$\begin{array}{c} \Phi_{(X,s_X)}^{\mathrm{mi}}(\mathbb{Q}_{\ell}(\beta))[-1] \\ \downarrow \\ \mathrm{cofiber}(c_1(\mathcal{L}_{X_0}) \otimes_{\mathbb{Q}_{\ell,X_0}} \mathbb{Q}_{\ell,X_0}(\beta) : \mathbb{Q}_{\ell,X_0}(\beta) \rightarrow \mathbb{Q}_{\ell,X_0}(\beta)) \\ \downarrow g \\ i^*j_*\mathbb{Q}_{\ell,X_{\mathcal{U}}}(\beta). \end{array} \quad (5.2.10)$$

Both f and g in these two fiber-cofiber sequences are defined by the universal property of the object in the middle. Therefore, if we consider the diagram

$$\begin{array}{ccccc} \mathbb{Q}_{\ell,X_0}(\beta) & \xrightarrow{c_1(\mathcal{L}_{X_0}^{\vee}) \otimes_{\mathbb{Q}_{\ell,X_0}} \mathbb{Q}_{\ell,X_0}(\beta)} & \mathbb{Q}_{\ell,X_0}(\beta) & \longrightarrow & \mathrm{cofiber}(c_1(\mathcal{L}_{X_0}^{\vee}) \otimes_{\mathbb{Q}_{\ell,X_0}} \mathbb{Q}_{\ell,X_0}(\beta)) \\ & \searrow \sim 0 & \downarrow \tilde{f} \quad \downarrow \tilde{g} & \nearrow f & \\ & & i^*j_*\mathbb{Q}_{\ell,X_{\mathcal{U}}}(\beta), & & \end{array} \quad (5.2.11)$$

where the middle vertical arrows induce the morphism f and g respectively, it suffices to show $\tilde{f} \sim \tilde{g}$. By definition, \tilde{f} is induced by the counit $\mathbb{Q}_{\ell,X}(\beta) \rightarrow j_*j^*\mathbb{Q}_{\ell,X}(\beta)$, while \tilde{g} is induced by the base change morphism $b.c. : s_X^*j_{0*}\mathbb{Q}_{\ell,\mathcal{U}_X}(\beta) \rightarrow j_*s_{\mathcal{U}}^*\mathbb{Q}_{\ell,\mathcal{U}_X}(\beta)$:

$$\tilde{g} : i^*s_X^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) \rightarrow i^*s_X^*j_{0*}j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) \rightarrow i^*j_*s_{\mathcal{U}}^*j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta). \quad (5.2.12)$$

It also suffices to show that the two arrows are homotopic before applying i^* . Consider the diagram

$$\begin{array}{ccccc} s_X^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) & \longrightarrow & s_X^*j_{0*}j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) & \xrightarrow{b.c.} & j_*s_{\mathcal{U}}^*j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) \simeq j_*j^*s_X^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) \\ & \searrow & \downarrow & & \nearrow \\ & & s_X^*s_{X*}j_*s_{\mathcal{U}}^*j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) & & \\ & & \simeq & & \\ & & s_X^*s_{X*}j_*j^*s_X^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta), & & \end{array} \quad (5.2.13)$$

If we analyse the commutative triangle on the left, we see that the oblique arrow corresponds to the unit of the adjunction $((s_X \circ j)^*, (s_X \circ j)_*)$. Indeed, by definition, the vertical arrow is

$$s_X^*j_{0*}(j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta)) \xrightarrow{\mathrm{unit}(s_{\mathcal{U}}^*,s_{\mathcal{U}*})} s_{\mathcal{U}*}s_{\mathcal{U}}^*j_0^*\mathbb{Q}_{\ell,\mathbb{V}_X}(\beta) \quad (5.2.14)$$

and the horizontal arrow is the one induced by the unit of (j_0^*, j_{0*}) . The composition is exactly the unit of the adjunction $((j_0 \circ s_{\mathcal{U}})^*, (j_0 \circ s_{\mathcal{U}})_*)$. Hence, the claim is proved as $j_0 \circ s_{\mathcal{U}} \simeq s_X \circ j$. Now notice that s_{X*} is conservative (s_X is a closed morphism), i.e. the counit $s_X^* s_{X*} \rightarrow 1$ is a natural equivalence. If we compose the oblique arrow with it, we obtain the unit of (j^*, j_*) evaluated in $s_X^* \mathbb{Q}_{\ell, \mathbb{V}_X}(\beta)$. The statement about the module structures is clear. \square

Remark 5.2.3 Notice that our main theorem provides a generalization of the formula proved in [9]: assume that we are given a proper flat morphism $p : X \rightarrow S$ from a regular scheme to an excellent strictly henselian trait.⁸ Let $s_X : X \rightarrow \mathbb{A}_S^1$ be the pullback of the section $S \rightarrow \mathbb{A}_S^1$ given by the uniformizer π . Then we can consider the diagram

$$\begin{array}{ccccc}
 \sigma & \xrightarrow{i_\sigma} & S & \xleftarrow{j_\eta} & \eta \\
 p_\sigma \uparrow & & p \uparrow & & \uparrow p_\eta \\
 X_0 & \xrightarrow{i} & X & \xleftarrow{j} & X_\eta \\
 s_0 \downarrow & & \downarrow s_X & & \downarrow s_{\mathbb{G}_{m,X}} \\
 X & \xrightarrow{i_0} & \mathbb{A}_X^1 & \xleftarrow{j_0} & \mathbb{G}_{m,X}
 \end{array} \tag{5.2.15}$$

where all squares are cartesian. The theorem we have just proved tells us that

$$i^* \mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, s_X)) \simeq \Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}(\beta))[-1].$$

By the regularity assumption on X , $\mathbf{Sing}(X, s_X) \simeq \mathbf{Sing}(X_0)$. If we apply $p_{\sigma*}$ to the formula above we find:

$$\begin{aligned}
 p_{\sigma*} i^* \mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X_0)) &\simeq i_\sigma^* \mathcal{R}_S^{\ell, \vee}(\mathbf{Sing}(X_0)) \\
 &\simeq p_{\sigma*} \Phi_{(X, s_X)}^{\text{mi}}(\mathbb{Q}_{\ell}(\beta))[-1] \stackrel{\text{Lemma 4.2.4}}{\simeq} (p_{\sigma*} \Phi_p(\mathbb{Q}_{\ell, X}(\beta))[-1])^{\text{hl}}
 \end{aligned} \tag{5.2.16}$$

which is exactly the content of [9, Theorem 4.39]. Also notice that in this case

$$i^* s_X^* j_{0*} \mathbb{Q}_{\ell, \mathbb{G}_{m,X}}(\beta) \simeq \mathbb{Q}_{\ell, X_0}(\beta) \oplus \mathbb{Q}_{\ell, X_0}(\beta)[1], \tag{5.2.17}$$

as the first Chern class of the trivial line bundle is zero. This recovers the equivalence

$$(\mathbb{Q}_{\ell, \sigma}(\beta))^{\text{hl}} \simeq \mathbb{Q}_{\ell, \sigma}(\beta) \oplus \mathbb{Q}_{\ell, \sigma}(\beta)[1] \tag{5.2.18}$$

proved in *loc.cit.*

⁸ We shall recycle the notation that we have introduced in § 4.1.

6 The ℓ -adic realization of the dg category of singularities of a twisted LG model of rank r

In this section we will explain how, by means of a theorem due to D. Orlov and J. Burke–M. Walker, the main theorem we proved in the previous section allows us to compute the ℓ -adic realization of the dg category of singularities of the zero locus of a global section of any vector bundle on a regular scheme. This is the reason that led us to consider twisted LG model, as defined in §3.1, as the author was initially interested in computing the ℓ -adic realization of the zero locus of a multifunction $X \rightarrow \mathbb{A}_S^n$ (with X regular).

6.1 Reduction of codimension

In this section we will provide an ∞ -functorial lax monoidal enhancement of the “reduction of codimension” equivalence proved by Orlov ([42, Theorem 2.1]) and by Burke–Walker ([11, Theorem A.4]).

Context 6.1.1 Let S be a regular noetherian scheme of finite Krull dimension.

Definition 6.1.2 Notice that all we said in section Sect. 3.1 can be generalised *mutatis mutandis* to the situation where line bundles \mathcal{L}_S are replaced by vector bundles of a fixed rank r \mathcal{E}_S . In particular, for a fixed vector bundle \mathcal{E}_S on S , we can define a symmetric monoidal (ordinary) category $\mathrm{LG}_{(S, \mathcal{E}_S)}^{\boxplus}$, analogous to the one we defined in section Sect. 3.1.

Notation 6.1.3 Let \mathcal{E}_S be a vector bundle of rank r over S . We will denote $\mathbb{P}(\mathcal{E}_S) = \mathrm{Proj}_S(\mathrm{Sym}_{\mathcal{O}_S}(\mathcal{E}_S^\vee))$ and $\pi_S : \mathbb{P}(\mathcal{E}_S) \rightarrow S$ the associate projective bundle and projection. Moreover, we will denote by $\mathcal{O}(1)$ the twisting sheaf on $\mathbb{P}(\mathcal{E}_S)$.

Construction 6.1.4 We will construct an (ordinary) symmetric monoidal functor

$$\Xi^{\boxplus} : \mathrm{LG}_{(S, \mathcal{E}_S)}^{\boxplus} \rightarrow \mathrm{LG}_{(\mathbb{P}(\mathcal{E}_S^\vee), \mathcal{O}(1))}^{\boxplus} \quad (6.1.1)$$

following the lead of [11] and [42]. Let $(X, s) \in \mathrm{LG}_{(S, \mathcal{E}_S)}$. Consider $W_s \in \Gamma(\mathbb{P}(\mathcal{E}_X^\vee), \mathcal{O}(1))$ defined as the morphism

$$W_s : \mathcal{O}_{\mathbb{P}(\mathcal{E}_X^\vee)} \simeq \widetilde{\mathrm{Sym}_{\mathcal{O}_X}(\mathcal{E}_X)} \rightarrow \widetilde{\mathrm{Sym}_{\mathcal{O}_X}(\mathcal{E}_X)}(1) \simeq \mathcal{O}(1) \quad (6.1.2)$$

induced by the morphism of modules

$$\mathrm{Sym}_{\mathcal{O}_X}(\mathcal{E}_X) \rightarrow \mathrm{Sym}_{\mathcal{O}_X}(\mathcal{E}_X)(1) \quad (6.1.3)$$

induced by $s : \mathcal{O}_X \rightarrow \mathcal{E}_X^\vee$. Here $\widetilde{(-)}$ is the functor that associates a quasi-coherent module on $\mathbb{P}(\mathcal{E}_X^\vee)$ to a graded $\mathrm{Sym}_{\mathcal{O}_X}(\mathcal{E}_X)$ -module. The assignment

$$(X, s) \mapsto (\mathbb{P}(\mathcal{E}_X^\vee), W_s),$$

together with the obvious law for morphisms defines a functor

$$\Xi : \mathbf{LG}_{(S, \mathcal{E}_S)} \rightarrow \mathbf{LG}_{(\mathbb{P}(\mathcal{E}_S^\vee), \mathcal{O}(1))}. \quad (6.1.4)$$

It is immediate to observe that $(S, \mathcal{O}) \mapsto (\mathbb{P}(\mathcal{E}_S^\vee), \mathcal{O})$, i.e. the functor is compatible with the units of the two symmetric monoidal structures. It remains to show that

$$\Xi((X, s) \boxplus (Y, t)) \simeq \Xi((X, s)) \boxplus \Xi((Y, t)).$$

On the left hand side we have

$$(\mathbb{P}(\mathcal{E}_{X \times_S Y}^\vee), W_{s \boxplus t}),$$

while on the right hand side we have

$$(\mathbb{P}(\mathcal{E}_X^\vee) \times_{\mathbb{P}(\mathcal{E}_S^\vee)} \mathbb{P}(\mathcal{E}_Y^\vee), W_s \boxplus W_t).$$

Since $\mathbb{P}(\mathcal{E}_X^\vee) \times_{\mathbb{P}(\mathcal{E}_S^\vee)} \mathbb{P}(\mathcal{E}_Y^\vee) \simeq (X \times_S \mathbb{P}(\mathcal{E}_S^\vee)) \times_{\mathbb{P}(\mathcal{E}_S^\vee)} (Y \times_S \mathbb{P}(\mathcal{E}_S^\vee)) \simeq (X \times_S Y) \times_S \mathbb{P}(\mathcal{E}_S^\vee) \simeq \mathbb{P}(\mathcal{E}_{X \times_S Y}^\vee)$, it suffices to show that $W_{s \boxplus t} = W_s \boxplus W_t$. It is enough to do it Zariski-locally ($\mathcal{O}(1)$ is a sheaf). Locally, $\mathbb{P}(\mathcal{E}_{X \times_S Y}^\vee)$ is isomorphic to a projective space. We may consider the covering of $\mathbb{P}(\mathcal{E}_{X \times_S Y}^\vee)$ consisting of open affine subschemes $\text{Spec}(B \otimes_A C[\frac{T_1}{T_j}, \dots, \frac{T_n}{T_j}])$, where $\text{Spec}(A)$ (resp. $\text{Spec}(B)$, resp. $\text{Spec}(C)$) is an open affine subscheme of S (resp. X , resp. Y). The restriction to this open subset of $W_{s \boxplus t}$ is of the form

$$(f_1 \otimes 1 + 1 \otimes g_1) \cdot \frac{T_1}{T_j} + \dots + (f_n \otimes 1 + 1 \otimes g_n) \cdot \frac{T_n}{T_j},$$

while that of $W_s \boxplus W_t$ is

$$(f_1 \cdot \frac{T_1}{T_j} \otimes 1 + 1 \otimes g_1 \cdot \frac{T_1}{T_j}) + \dots + (f_n \cdot \frac{T_n}{T_j} \otimes 1 + 1 \otimes g_n \cdot \frac{T_n}{T_j}).$$

The claim follows and thus we obtain the desired symmetric monoidal functor.

If we compose the symmetric monoidal functor (6.1.1) with the lax monoidal ∞ -functor defined in Sect. 3.2 we get

$$\mathbf{Sing}(\mathbb{P}(\mathcal{E}_\bullet^\vee), W_\bullet)^\otimes : \mathbf{LG}_{(S, \mathcal{E}_S)}^{\text{op}, \boxplus} \rightarrow \mathbf{Mod}_{\mathbf{Sing}(\mathbb{P}(\mathcal{E}_S^\vee), \mathcal{O})}(\mathbf{dgCat}_S^{\text{idm}})^\otimes \quad (6.1.5)$$

which, at the level of objects, corresponds to the assignment

$$(X, s) \mapsto \mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), W_s).$$

We define the dg category of singularities of a twisted LG model (X, s) of rank r over (S, \mathcal{E}_S) in the following way: consider the derived zero locus of s , defined as the homotopy pullback of s along the zero section

$$\begin{array}{ccc} X_0 & \xrightarrow{i} & X \\ \downarrow & & \downarrow s \\ S & \xrightarrow{0} & \mathbb{V}(\mathcal{E}_S). \end{array} \quad (6.1.6)$$

As $0 : S \rightarrow \mathbb{V}(\mathcal{E}_S)$ is a closed lci morphism (locally, is of the form $U \rightarrow \mathbb{A}_S^r$), so is $i : X_0 \rightarrow X$. Thus the pushforward induces a dg functor

$$i_* : \mathbf{Sing}(X_0) \rightarrow \mathbf{Sing}(X). \quad (6.1.7)$$

Definition 6.1.5 We define

$$\mathbf{Sing}(X, s) := \text{Ker}(i_* : \mathbf{Sing}(X_0) \rightarrow \mathbf{Sing}(X)). \quad (6.1.8)$$

Similarly to what we did in Sect. 3.2, we can define a lax monoidal ∞ -functor

$$\mathbf{Sing}(\bullet, \bullet)^{\otimes} : \text{LG}_{(S, \mathcal{E}_S)}^{\text{op}, \boxplus} \rightarrow \mathbf{Mod}_{\mathbf{Sing}(S, 0)}(\mathbf{dgCat}_S^{\text{idm}})^{\otimes} \quad (6.1.9)$$

At the level of objects, it is defined by $(X, s) \mapsto \mathbf{Sing}(X, s)$.

Remark 6.1.6 Let $\text{Spec}(B)$ be an affine scheme and let E_B be a projective B -module of constant rank r and $s \in E_B$. The derived zero locus of s is the spectrum of $B \otimes_{\text{Sym}_B(E_B^\vee)}^\mathbb{L} B$, where the two augmentations $\text{Sym}_B(E_B^\vee) \rightarrow B$ are determined by s and 0. The dg algebra associated to the simplicial ring $B \otimes_{\text{Sym}_B(E_B^\vee)}^\mathbb{L} B$ via the Dold-Kan correspondence is the Koszul dg algebra associated to (B, E_B^\vee, s)

$$K(B, E_B^\vee, s) := 0 \rightarrow \bigwedge^r E_B^\vee \rightarrow \bigwedge^{r-1} E_B^\vee \rightarrow \cdots \rightarrow \bigwedge^2 E_B^\vee \rightarrow E_B^\vee \xrightarrow{s} B \rightarrow 0 \quad (6.1.10)$$

concentrated in degrees $[-r, 0]$. This is locally true (see for example [43, Remark 1.22] or [31]) and the global statement follows from the existence of a morphism of dg algebras

$$K(B, E_B^\vee, s) \rightarrow N(B \otimes_{\text{Sym}_B(E_B^\vee)}^\mathbb{L} B), \quad (6.1.11)$$

where N is the normalized Moore complex functor.

Construction 6.1.7 Similarly to what we said in Construction 3.2.12, there is an equivalence between $\mathbf{QCoh}(B \otimes_{\mathbb{L}_{\mathrm{Sym}_B(E_B^\vee)}} B)$ and the category of cofibrant $K(B, E_B^\vee, s)$ -dg modules, denoted by $\widehat{K(B, E_B^\vee, s)}$. Under this equivalence, $\mathbf{Coh}^b(B \otimes_{\mathbb{L}_{\mathrm{Sym}_B(E_B^\vee)}} B)$ corresponds to the full subcategory of $\widehat{K(B, E_B^\vee, s)}$ spanned by $K(B, E_B^\vee, s)$ -dg modules that are cohomologically bounded and whose cohomology is coherent over $\mathrm{coker}(E_B^\vee \xrightarrow{s} B)$.

Similarly, $\mathbf{Perf}(B \otimes_{\mathbb{L}_{\mathrm{Sym}_B(E_B^\vee)}} B)$ corresponds to the full subcategory of $\widehat{K(B, E_B^\vee, s)}$ spanned by homotopically finitely presented dg modules. We consider the full subcategory $\mathrm{Coh}^s(B, E_B^\vee, s)$ spanned by $K(B, E_B^\vee, s)$ -dg modules that are cohomologically bounded, whose cohomology is coherent over $\mathrm{coker}(E_B^\vee \xrightarrow{s} B)$ and whose underlying complex of B -modules is strictly perfect.

Lemma 6.1.8 [9, Lemma 2.33] *Let $\mathrm{Coh}^s(B, E_B^\vee, s)^{\mathrm{acy}}$ be the full sub-category of $\mathrm{Coh}^s(B, E_B^\vee, s)$ spanned by acyclic dg modules. Then the cofibrant replacement induces an equivalence of dg categories*

$$\mathrm{Coh}^s(B, E_B^\vee, s)[\mathrm{q.iso}^{-1}] \simeq \mathrm{Coh}^s(B, E_B^\vee, s) / \mathrm{Coh}^s(B, E_B^\vee, s)^{\mathrm{acy}} \quad (6.1.12)$$

$$\simeq \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}.$$

If we label $\mathbf{Perf}^s(B, E_B^\vee, s)$ the full subcategory of $\mathrm{Coh}^s(B, E_B^\vee, s)$ spanned by perfect $K(B, E_B^\vee, s)$ -dg modules, there are equivalences of dg categories

$$\mathrm{Coh}^s(B, E_B^\vee, s) / \mathbf{Perf}^s(B, E_B^\vee, s) \simeq \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} / \mathbf{Perf}(X_0) \simeq \mathbf{Sing}(X, s). \quad (6.1.13)$$

Proof The proof of [9, Lemma 2.33] applies, *mutatis mutandis*. \square

In the same way as in Construction 3.2.14 and in the following discussion, we can construct a lax monoidal ∞ -functor

$$\mathbf{Coh}^b(\bullet)_{\mathbf{Perf}(\bullet)}^{\otimes} : \mathrm{LG}_{(S, \mathcal{E}_S)}^{op, \boxplus} \rightarrow \mathbf{dgCat}_S^{\mathrm{idm}, \otimes} \quad (6.1.14)$$

Moreover, using the same arguments as in the discussion following Remark 1.27 in [43], we get the lax monoidal ∞ -functor (6.1.9).

At this point, the reader might complain that there are too many **Sing**'s involved, and this may cause confusion. As a partial justification to the choice of notation we made, let us show that these different ∞ -functors are closely related.

Definition 6.1.9 Let $(X, s) \in \mathrm{LG}_{(S, \mathcal{E}_S)}$. We define the projective bundle over X_0 associated to $\mathcal{E}_{X_0} = i^* \mathcal{E}_X$ as the derived pullback

$$\begin{array}{ccc} \mathbb{P}(\mathcal{E}_{X_0}^\vee) & \longrightarrow & \mathbb{P}(\mathcal{E}_X^\vee) \\ p_0 \downarrow & & \downarrow p \\ X_0 & \xrightarrow{i} & X. \end{array} \quad (6.1.15)$$

Recall that, given a global section of a line bundle on a scheme, we define its (derived) zero locus as in Definition 3.2.1. In particular, for $(X, s) \in \mathrm{LG}_{(S, \mathcal{E}_S)}$ we have a global section W_s of the line bundle $\mathcal{O}(1)$ on $\mathbb{P}(\mathcal{E}_X^\vee)$ and thus we have a (derived) pullback square

$$\begin{array}{ccc} V(W_s) & \xrightarrow{k} & \mathbb{P}(\mathcal{E}_X^\vee) \\ \downarrow k & & \downarrow W_s \\ \mathbb{P}(\mathcal{E}_X^\vee) & \xrightarrow{0} & \mathbb{V}(\mathcal{O}(1)). \end{array} \quad (6.1.16)$$

Let $(X, s) \in \mathrm{LG}_{(S, \mathcal{E}_S)}$ and consider the following diagram, where both squares are homotopy cartesian:

$$\begin{array}{ccccc} & & \mathbb{P}(\mathcal{E}_X^\vee) & \xrightarrow{W_s} & \mathbb{V}(\mathcal{O}(1)) \\ & & \uparrow k & & \uparrow 0 \\ \mathbb{P}(\mathcal{E}_{X_0}^\vee) & \xrightarrow{j} & V(W_s) & \xrightarrow{k} & \mathbb{P}(\mathcal{E}_X^\vee) \\ \downarrow p_0 & & & & \downarrow p \\ X_0 & \xrightarrow{i} & & & X. \end{array} \quad (6.1.17)$$

By [28, Chapter 4, Lemma 3.1.3, Lemma 5.1.4], $j_* p_0^*$ preserves complexes with coherent bounded cohomology. The derived proper base change equivalence $k_* j_* p_0^* \simeq p^* i_*$ implies that we have a dg functor

$$\Upsilon_{(X, s)} = j_* p_0^* : \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \rightarrow \mathbf{Coh}^b(V(W_s))_{\mathbf{Perf}(\mathbb{P}(\mathcal{E}_X^\vee))}. \quad (6.1.18)$$

and it is functorial in (X, s) . We will need to enhance the assignment $(X, s) \mapsto \Upsilon_{(X, s)}$ with a lax monoidal structure. We will use strict models.

Assume that $S = \mathrm{Spec}(A)$ is affine and let E be a projective A module of rank r . Since E is an A -module of finite type, there exists a surjection

$$A^n \rightarrow E. \quad (6.1.19)$$

Denote by t_i ($i = 1, \dots, n$) the images in E of the elements $(1, 0, \dots, 0)$, \dots , $(0, \dots, 0, 1) \in A^n$.

We will define a lax monoidal ∞ -natural transformation

$$\Upsilon^{\otimes} : \mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes} \rightarrow \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(E_{\bullet}^{\vee}))}^{\otimes}, \quad (6.1.20)$$

where

$$\mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes}, \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(E_{\bullet}^{\vee}))}^{\otimes} : \mathbf{LG}_{(S,E)}^{\boxplus, \text{aff}, \text{op}} \rightarrow \mathbf{dgCat}_S^{\text{idm}, \otimes}. \quad (6.1.21)$$

For every $(\text{Spec}(B), s) \in \mathbf{LG}_{(S,E)}^{\boxplus, \text{aff}, \text{op}}$,

$$\mathbb{P}(E_B^{\vee}) = \text{Proj}(\text{Sym}_B(E_B))$$

has a Zariski affine covering $\{D_+(t_i)\}_{i=1}^n$, where

$$D_+(t_i) = \text{Spec}(\text{Sym}_B(E_B^{\vee})_{(t_i)}). \quad (6.1.22)$$

Here $\text{Sym}_B(E_B^{\vee})_{(t_i)}$ denotes the ring of degree 0 elements of the graded ring $\text{Sym}_B(E_B^{\vee})[t_i^{-1}]$. Since

$$\mathbf{Coh}^b(V(W_s))_{\mathbf{Perf}(\mathbb{P}(E_B^{\vee}))}^{\otimes} \simeq \varprojlim \mathbf{Coh}^b(V(W_{s|U_{\bullet}}))_{\mathbf{Perf}(U_{\bullet})}^{\otimes}, \quad (6.1.23)$$

where U_{\bullet} is the Čech hypercover of $\{D_+(t_i)\}_{i=1}^n$, it will suffice to define an homotopy coherent diagram of lax monoidal ∞ -natural transformations

$$\mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes} \rightarrow \mathbf{Coh}^b(V(W_{\bullet|U_{\bullet}}))_{\mathbf{Perf}(U_{\bullet})}^{\otimes}. \quad (6.1.24)$$

Recall that $\mathbf{Coh}^s(B, E_B^{\vee}, s_B)$ is the A -dg category of $K(B, E_B^{\vee}, s_B)$ -dg modules whose underlying B -dg module is strictly perfect. In other words, it is the dg category of B -dg modules (M, d) endowed with a B -linear map

$$h : M \rightarrow M \otimes_B E_B[-1] \quad (6.1.25)$$

such that

$$h^2 = 0, \quad [d, h] = id_M \otimes s_B : M \rightarrow M \otimes_B E_B. \quad (6.1.26)$$

Construction 6.1.10 For every $(B, s_B) \in \mathbf{LG}_{(S,E)}^{\boxplus, \text{aff}, \text{op}}$ and every $i = 1, \dots, n$, define a pseudo-functor

$$\Upsilon_i : \mathbf{Coh}^s(B, E_B^{\vee}, s_B) \rightarrow \mathbf{Coh}^s(\text{Sym}_B(E_B)_{(t_i)}, t_i^{-1} \cdot \text{Sym}_B(E_B)_{(t_i)}, W_{s_B|D_+(t_i)}) \quad (6.1.27)$$

as follows. For $(M, d, h) \in \text{Coh}^s(B, E_B^\vee, s_B)$, let

$$\Upsilon_i(M, d, h) = (M \otimes_B \text{Sym}_B(E_B), d \otimes id, \chi_{i,h}). \quad (6.1.28)$$

The map $\chi_{i,h} : M \otimes_B \text{Sym}_B(E_B) \rightarrow M \otimes_B t_i \cdot \text{Sym}_B(E_B)$ is defined as the restriction to $D_+(t_i)$ of the composition

$$\begin{array}{ccc} M \otimes_B \text{Sym}_B(E_B) & \xrightarrow{h \otimes id} & M \otimes_B E_B \otimes_B \text{Sym}_B(E_B)[-1] \\ & \searrow & \downarrow \\ & & M \otimes_B \text{Sym}_B(E_B)(1)[-1], \end{array} \quad (6.1.29)$$

where $\text{Sym}_B(E_B)(1)$ is the $\text{Sym}_B(E_B)$ module $\text{Sym}_B(E_B)$ with the grading shifted by 1 and the second morphism is induced by multiplication. Notice that $\text{Sym}_B(E_B)(1) \otimes_{\text{Sym}_B(E_B)} \otimes_{\text{Sym}_B(E_B)(t_i)} = t_i \cdot \text{Sym}_B(E_B)(t_i)$. The desired properties that

$$\chi_{i,h}^2 = 0, \quad [d \otimes id, \chi_{i,h}] = W_{s_B|D_+(t_i)} \otimes id \quad (6.1.30)$$

are local (on $\text{Spec}(B)$) and if $E_B = B^r$, $s_B = (s_{B,1}, \dots, s_{B,n})$, then Υ_i admits the explicit description

$$\Upsilon_i(M, d, h=\{h_j\}_{j=1,\dots,r}) = (M \otimes_B B[x_1, \dots, x_r], d \otimes id, x_1 \cdot h_1 + \dots + x_r \cdot h_r) \quad (6.1.31)$$

and $W_{s_B|D_+(t_i)} = x_1 \cdot s_{B,1} + \dots + x_r \cdot s_{B,r}$.

It follows immediately from the definitions of the pseudo functorial structures of $(B, s_B) \mapsto \text{Coh}^s(B, E_B^\vee, s_B)$ and $(B, s_B) \mapsto \text{Coh}^s(\text{Sym}_B(E_B)(t_i), t_i^{-1} \cdot \text{Sym}_B(E_B)(t_i), W_{s_B|D_+(t_i)})$ that Υ_i is a pseudo natural transformation.

Remark 6.1.11 Let $X = \text{Spec}(B)$ and $X_0 = \text{Spec}(K(B, E_B^\vee, s_B))$. Notice that Υ_i models the dg functor

$$j_* p_0^* : \mathbf{Coh}^b(X_0) \mathbf{Perf}(X) \rightarrow \mathbf{Coh}^b(V(W_{s_B}) \cap D_+(t_i)) \mathbf{Perf}(D_+(t_i)), \quad (6.1.32)$$

where

$$p_0 : D_+(t_i) \times_{\mathbb{P}(E_X^\vee)} \mathbb{P}(E_{X_0}^\vee) \rightarrow X_0$$

and

$$j : D_+(t_i) \times_{\mathbb{P}(E_X^\vee)} \mathbb{P}(E_{X_0}^\vee) \rightarrow V(W_{s_B}) \cap D_+(t_i).$$

Construction 6.1.12 We will now endow Υ_i with a pseudo lax monoidal structure. If (B, s_B) and (C, s_C) are affine twisted LG models of rank r over (A, E) , then

$$\mu : \text{Coh}^s(B, E_B^\vee, s_B) \otimes \text{Coh}^s(C, E_C^\vee, s_C) \rightarrow \text{Coh}^s(B \otimes_A C, E_{B \otimes_A C}^\vee, s_B \boxplus s_C) \quad (6.1.33)$$

is defined on objects by

$$((M, d_M, h_M), (N, d_N, h_N)) \mapsto (M \otimes_A N, d_{M \otimes_A N}, h_M \boxplus h_N), \quad (6.1.34)$$

where $h_M \boxplus h_N$ is the map

$$\begin{array}{ccc} M \otimes_A N & \xrightarrow{\begin{bmatrix} h_M \otimes C \\ B \otimes h_N \end{bmatrix}} & (M \otimes_A N) \otimes_{B \otimes_A C} E_{B \otimes_A C}^{\oplus 2}[-1] \\ & \searrow & \downarrow [1 \ 1] \\ & & (M \otimes_A N) \otimes_{B \otimes_A C} E_{B \otimes_A C}[-1]. \end{array} \quad (6.1.35)$$

It is defined on morphisms in the obvious way.

The dg functor

$$\begin{array}{c} \text{Coh}^s(\text{Sym}_B(E_B)_{(t_i)}, t_i^{-1} \cdot \text{Sym}_B(E_B)_{(t_i)}, W_{s_B|D_+(t_i)}) \\ \otimes \\ \text{Coh}^s(\text{Sym}_C(E_C)_{(t_i)}, t_i^{-1} \cdot \text{Sym}_C(E_C)_{(t_i)}, W_{s_C|D_+(t_i)}) \\ \mu \downarrow \\ \text{Coh}^s(\text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, t_i^{-1} \cdot \text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, W_{s_B \boxplus s_C|D_+(t_i)}) \end{array} \quad (6.1.36)$$

is defined similarly.

We need to verify that the two compositions

$$\begin{array}{ccc} \text{Coh}^s(B, E_B^\vee, s_B) \otimes \text{Coh}^s(C, E_C^\vee, s_C) & & \\ \downarrow \mu \circ (\Upsilon_{(B, s_B)} \otimes \Upsilon_{(C, s_C)}) & \downarrow \Upsilon_{(B \otimes_A C, s_B \boxplus s_C)} \circ \mu & \\ \text{Coh}^s(\text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, t_i^{-1} \cdot \text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, W_{(s_B \boxplus s_C)|D_+(t_i)}) & & \end{array} \quad (6.1.37)$$

$$\text{Coh}^s(\text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, t_i^{-1} \cdot \text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, W_{(s_B \boxplus s_C)|D_+(t_i)})$$

are isomorphic up to natural isomorphism. The composition $\Upsilon_{(B \otimes_A C, s_B \boxplus s_C)} \circ \mu$ is

$$\begin{array}{l} ((M, d_M, h_M), (N, d_N, h_N)) \mapsto \\ (M \otimes_A N \otimes_{B \otimes_A C} \text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, d \otimes id, \chi_{i, h_M \boxplus h_N}), \end{array} \quad (6.1.38)$$

while the composition $\mu \circ (\Upsilon_{(B, s_B)} \otimes \Upsilon_{(C, s_C)})$ is

$$\begin{aligned} ((M, d_M, h_M), (N, d_N, h_N)) &\mapsto \\ &((M \otimes_B \text{Sym}_B(E_B)_{(t_i)}) \otimes_{\text{Sym}_A(E)_{(t_i)}} \\ &(N \otimes_C \text{Sym}_C(E_C)_{(t_i)}), d \otimes id, \chi_{i, h_M} \boxplus \chi_{i, h_N}). \end{aligned} \quad (6.1.39)$$

It is clear that there is a canonical isomorphism of $B \otimes_A C$ -dg modules

$$\begin{aligned} (M \otimes_A N \otimes_{B \otimes_A C} \text{Sym}_{B \otimes_A C}(E_{B \otimes_A C})_{(t_i)}, d \otimes id) \\ \simeq ((M \otimes_B \text{Sym}_B(E_B)_{(t_i)}) \otimes_{\text{Sym}_A(E)_{(t_i)}} (N \otimes_C \text{Sym}_C(E_C)_{(t_i)}), d \otimes id). \end{aligned} \quad (6.1.40)$$

Then we only need to show that

$$\chi_{i, h_M} \boxplus h_N = \chi_{i, h_M} \boxplus \chi_{i, h_N} \quad (6.1.41)$$

under this natural isomorphism. This follows from the definitions.

The units of the lax monoidal pseudo functors $(B, s_B) \mapsto \text{Coh}^s(B, E_B^\vee, s_B)$ and $(B, s_B) \mapsto \text{Coh}^s(\text{Sym}_B(E_B)_{(t_i)}, t_i^{-1} \cdot \text{Sym}_B(E_B)_{(t_i)}, W_{s_B})$ are compatible: the diagram

$$\begin{array}{ccc} & \text{Coh}^s(A, E^\vee, 0) & \\ \underline{A} \swarrow & \downarrow & \searrow \\ & \text{Coh}^s(\text{Sym}_A(E)_{(t_i)}, t_i^{-1} \text{Sym}_A(E)_{(t_i)}, 0) & \end{array} \quad (6.1.42)$$

commutes. Here \underline{A} is the dg category with one object $*$ and $\text{End}(*) = A$. The diagonal arrows are determined by

$$* \mapsto A, \quad * \mapsto \text{Sym}_A(E)_{(t_i)} \quad (6.1.43)$$

where A (resp. $\text{Sym}_A(E)_{(t_i)}$) is concentrated in degree 0 and the map $A \rightarrow E[-1]$ (resp. $\text{Sym}_A(E)_{(t_i)} \rightarrow t_i \text{Sym}_A(E)_{(t_i)}$) is zero.

The previous constructions provide us with pseudo lax monoidal natural transformations

$$\Upsilon_i^\otimes : \text{Coh}^s(\bullet, E_\bullet^\vee, s_\bullet)^\otimes \rightarrow \text{Coh}^s(\text{Sym}_\bullet(E_\bullet)_{(t_i)}, t_i^{-1} \text{Sym}_\bullet(E_\bullet), W_{s_\bullet})^\otimes. \quad (6.1.44)$$

It is obvious that the Υ_i^\otimes preserve quasi-isomorphisms and that they are compatible with one another. In other words, they can be used to define the homotopy coherent diagram of lax monoidal ∞ -natural transformations (6.1.24).

Therefore, we get a lax monoidal ∞ -natural transformation

$$\Upsilon^{\otimes} : \mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes} \rightarrow \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(E_{\bullet}^{\vee}))}^{\otimes} \quad (6.1.45)$$

between the ∞ -functors

$$\mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes}, \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(E_{\bullet}^{\vee}))}^{\otimes} : \mathbf{LG}_{(A,E)}^{\text{aff}, \boxtimes, \text{op}} \rightarrow \mathbf{dgCat}_S^{\text{idm}, \otimes}. \quad (6.1.46)$$

By Kan extension and descent, we extend Υ^{\otimes} to a lax monoidal ∞ -natural transformation between

$$\mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes}, \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(E_{\bullet}^{\vee}))}^{\otimes} : \mathbf{LG}_{(A,E)}^{\boxtimes, \text{op}} \rightarrow \mathbf{dgCat}_S^{\text{idm}, \otimes} \quad (6.1.47)$$

i.e. we extend to all twisted LG models of rank r over (A, E) .

Finally, if S is not affine, we define the lax monoidal ∞ -natural transformation

$$\Upsilon^{\otimes} : \mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)} \rightarrow \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(\mathcal{E}_{\bullet}^{\vee}))} \quad (6.1.48)$$

of lax monoidal ∞ -functors $\mathbf{LG}_{(S,\mathcal{E})}^{\boxplus, \text{op}} \rightarrow \mathbf{dgCat}_S^{\text{idm}, \otimes}$ as

$$\varprojlim_{\text{Spec}(A) \rightarrow S} (\Upsilon_{(A,\mathcal{E}_A)}^{\otimes} : \mathbf{Coh}^b(V(\bullet))_{\mathbf{Perf}(\bullet)}^{\otimes} \rightarrow \mathbf{Coh}^b(V(W_{\bullet}))_{\mathbf{Perf}(\mathbb{P}(E_{\bullet}^{\vee}))}^{\otimes}), \quad (6.1.49)$$

where the limit is taken over Zariski open subschemes $\text{Spec}(A) \rightarrow S$. Here $\Upsilon_{(A,\mathcal{E}_A)}^{\otimes}$ is the lax monoidal ∞ -natural transformation we defined above. Once again, we have used that

$$\mathbf{dgCat}_S^{\text{idm}, \otimes} = \varprojlim_{\text{Spec}(A) \rightarrow S} \mathbf{dgCat}_A^{\text{idm}, \otimes} \quad (6.1.50)$$

and that

$$\mathbf{LG}_{(S,\mathcal{E})}^{\boxplus} \simeq \varprojlim_{\text{Spec}(A) \rightarrow S} \mathbf{LG}_{(A,\mathcal{E}_A)}^{\boxplus} \quad (6.1.51)$$

(this is an analogue of Lemma 3.1.9 for twisted LG models of rank r , which can be proved *mutatis mutandis*).

Remark 6.1.13 Let (X, s_X) be a twisted LG model of rank r over (S, \mathcal{E}) . Then

$$\Upsilon_{(X,s_X)} = j_* p_0^* : \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \rightarrow \mathbf{Coh}^b(V(W_{s_X}))_{\mathbf{Perf}(\mathbb{P}(\mathcal{E}_X^{\vee}))}, \quad (6.1.52)$$

where we have used the notation of diagram (6.1.17)

Lemma 6.1.14 $j : \mathbb{P}(\mathcal{E}_X^\vee) \rightarrow V(W_s)$ is an lci morphism of derived schemes.

Proof Since the property of being lci is local, we can assume that $\mathcal{E}_X \simeq \mathcal{O}_X^r$ and $s_X = \underline{f} \in \mathcal{O}_X^r(X)$. Since j is clearly of finite presentation, we will only need to show that the relative cotangent complex is of Tor amplitude $[-1, 0]$. Recall that we have the fundamental fiber-cofiber sequence

$$j^* \mathbb{L}_{V(\underline{f})/\mathbb{P}_X^{r-1}} \rightarrow \mathbb{L}_{\mathbb{P}_{X_0}^{r-1}/\mathbb{P}_X^{r-1}} \rightarrow \mathbb{L}_{\mathbb{P}_{X_0}^{r-1}/V(\underline{f})} \quad (6.1.53)$$

at our disposal. Since $\mathbb{P}_{X_0}^{r-1} \simeq S \times_{\mathbb{A}_S^r}^h \mathbb{P}_X^{r-1}$, $\mathbb{L}_{\mathbb{P}_{X_0}^{r-1}/\mathbb{P}_X^{r-1}}$ is equivalent to the pullback of $\mathbb{L}_{S/\mathbb{A}_S^r} \simeq \mathcal{O}_S^r[1]$ along $\mathbb{P}_{X_0}^{r-1} \rightarrow S$, i.e. $\mathbb{L}_{\mathbb{P}_{X_0}^{r-1}/\mathbb{P}_X^{r-1}} \simeq \mathcal{O}_{\mathbb{P}_{X_0}^{r-1}}^r[1]$. On the other hand, $\mathbb{L}_{V(\underline{f})/\mathbb{P}_X^{r-1}} \simeq k^* \mathbb{L}_{\mathbb{P}_X^{r-1}/V(\mathcal{O}(-1))} \simeq k^*(\mathcal{O}(-1)[1])$. Therefore, $j^* \mathbb{L}_{V(\underline{f})/\mathbb{P}_X^{r-1}} \simeq \mathcal{O}_{\mathbb{P}_{X_0}^{r-1}}^r \otimes_{\mathcal{O}_{\mathbb{P}_X^{r-1}}} \mathcal{O}(-1)$. We just need to identify the morphism $j^* \mathbb{L}_{V(\underline{f})/\mathbb{P}_X^{r-1}} \simeq \mathcal{O}_{\mathbb{P}_{X_0}^{r-1}}^r \otimes_{\mathcal{O}_{\mathbb{P}_X^{r-1}}} \mathcal{O}(-1)[1] \rightarrow \mathcal{O}_{\mathbb{P}_{X_0}^{r-1}}^r[1] \simeq \mathbb{L}_{\mathbb{P}_{X_0}^{r-1}}$. It coincides with the morphism (T_1, \dots, T_r) . In particular, the cofiber of this morphism, i.e. $\mathbb{L}_{\mathbb{P}_{X_0}^{r-1}/V(\underline{f})}$, is equivalent to $\mathcal{O}_{\mathbb{P}_{X_0}^{r-1}}^{r-1}[1]$. \square

Construction 6.1.15 Since j is an lci morphism, the dg functor (6.1.48) preserves perfect complexes. Therefore, for every $(X, s_X) \in \mathbf{LG}_{(S, \mathcal{E}_S)}$, we have an induced dg functor

$$\Upsilon_{(X, s_X)} := j_* p_0^* : \mathbf{Sing}(X, s_X) \rightarrow \mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X}) \quad (6.1.54)$$

Starting from (6.1.48), by the usual standard arguments we thus obtain a lax monoidal ∞ -natural transformation

$$\Upsilon^\otimes : \mathbf{Sing}(\bullet, \bullet)^\otimes \rightarrow \mathbf{Sing}(\mathbb{P}(\mathcal{E}_\bullet^\vee), W_\bullet)^\otimes \quad (6.1.55)$$

of functors

$$\mathbf{LG}_{(S, \mathcal{E}_S)}^{\text{op}, \boxplus} \rightarrow \mathbf{Mod}_{\mathbf{Sing}(S, 0)}(\mathbf{dgCat}_S^{\text{idm}})^\otimes,$$

where $\mathbf{Sing}(\mathbb{P}(\mathcal{E}_\bullet^\vee), W_\bullet)^\otimes$ denotes the composition of (6.1.5) with

$$\mathbf{Mod}_{\mathbf{Sing}(\mathbb{P}(\mathcal{E}_S), 0)}(\mathbf{dgCat}_S^{\text{idm}})^\otimes \rightarrow \mathbf{Mod}_{\mathbf{Sing}(S, 0)}(\mathbf{dgCat}_S^{\text{idm}})^\otimes.$$

Theorem 6.1.16 ([11, 42]) *If s_X is a regular section, the lax monoidal ∞ -natural transformation (6.1.55) induces an equivalence of dg categories*

$$\Upsilon_{(X, s_X)} : \mathbf{Sing}(X, s_X) \simeq \mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X}). \quad (6.1.56)$$

Proof This is an immediate consequence of [11, Theorem A.4]. Indeed, as the dg categories are triangulated, it suffices to show that the statement is true on the induced functor of triangulated categories. This coincides by construction with that of *loc. cit.* \square

Remark 6.1.17 We actually believe that the statement above remains true even if s_X is not assumed to be regular, as long as one considers the derived zero loci instead of the classical ones.

6.2 The ℓ -adic realization of the dg category of singularities of a twisted LG model of rank r

It is now easy to obtain the following computation:

Theorem 6.2.1 *Let (X, s_X) be a twisted LG model of rank r over (S, \mathcal{E}_S) . Assume that X is a regular scheme and that s_X is a regular global section of \mathcal{E}_X . The following equivalence holds in $\mathbf{Mod}_{\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0))}(\mathbf{Shv}_{\mathbb{Q}_\ell}(X))$*

$$\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, s_X)) \simeq p_* i_* \Phi_{(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X})}^{\mathrm{mi}}(\mathbb{Q}_\ell(\beta))[-1], \quad (6.2.1)$$

where $i : V(W_{s_X}) \rightarrow \mathbb{P}(\mathcal{E}_X^\vee)$ is the closed embedding of the zero locus of W_{s_X} and $p : \mathbb{P}(\mathcal{E}_X^\vee) \rightarrow X$ is the canonical projection.

Proof As X is regular and s_X is a regular section, we have an equivalence $\mathbf{Sing}(X, s_X) \simeq \mathbf{Sing}(V(s_X))$. Notice that in this situation $V(s_X)$ coincides with the underived zero locus of s_X . As $\mathbb{P}(\mathcal{E}_X^\vee)$ is regular, by Theorem 6.1.16, we have that

$$\mathbf{Sing}(X, s_X) \simeq \mathbf{Sing}(V(W_{s_X})) \simeq \mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X}).$$

Then

$$\begin{aligned} \mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, s_X)) &\simeq p_* \mathcal{R}_{\mathbb{P}(\mathcal{E}_X^\vee)}^{\ell, \vee}(\mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X})) \\ &\stackrel{\text{Theorem 5.2.2}}{\simeq} p_* i_* \Phi_{(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X})}^{\mathrm{mi}}(\mathbb{Q}_\ell(\beta))[-1]. \end{aligned} \quad (6.2.2)$$

Notice that since s_X is a regular global section of \mathcal{E}_X , W_{s_X} is a regular global section of $\mathcal{O}_{\mathbb{P}(\mathcal{E}_X^\vee)}$ and the hypothesis of Theorem 5.2.2 apply to the pair $(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X})$.

The fact that this equivalence holds in $\mathbf{Mod}_{\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0))}(\mathbf{Shv}_{\mathbb{Q}_\ell}(X))$ follows immediately from the fact that

$$\mathcal{R}_{\mathbb{P}(\mathcal{E}_X^\vee)}^{\ell, \vee}(\mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X})) \simeq i_* \Phi_{(\mathbb{P}(\mathcal{E}_X^\vee), W_{s_X})}^{\mathrm{mi}}(\mathbb{Q}_\ell(\beta))[-1]$$

is an equivalence of $\mathcal{R}_{\mathbb{P}(\mathcal{E}_X^\vee)}^{\ell, \vee}(\mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), 0))$ -modules and from the fact that

$$\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0)) \rightarrow p_* \mathcal{R}_{\mathbb{P}(\mathcal{E}_X^\vee)}^{\ell, \vee}(\mathbf{Sing}(\mathbb{P}(\mathcal{E}_X^\vee), 0)) \quad (6.2.3)$$

is a morphism of commutative algebras. \square

Corollary 6.2.2 *Assume that $S = \text{Spec}(A)$ is a noetherian regular local ring of dimension n and let π_1, \dots, π_n be generators of the maximal ideal. Let $p : X \rightarrow S = \text{Spec}(A)$ be a regular, flat S -scheme of finite type. Let $\underline{\pi} : S \rightarrow \mathbb{A}_S^n$ be the closed embedding associated to π_1, \dots, π_n . Then $\underline{\pi} \circ p$ is a regular global section of \mathcal{O}_X^n . Then the equivalence*

$$\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, \underline{\pi} \circ p)) \simeq q_* i_* \Phi_{(\mathbb{P}_X^{n-1}, W_{\underline{\pi} \circ p})}^{\text{mi}}(\mathbb{Q}_\ell(\beta))[-1]$$

holds in $\mathbf{Mod}_{\mathcal{R}_X^{\ell, \vee}(\mathbf{Sing}(X, 0))}(\mathbf{Shv}_{\mathbb{Q}_\ell}(X))$.

Here $q : \mathbb{P}_X^{n-1} = \text{Proj}_X(\mathcal{O}_X[T_1, \dots, T_n]) \rightarrow X$ is the canonical projection and $i : V(W_{\underline{\pi} \circ p}) \rightarrow \mathbb{P}_X^{n-1}$ is the closed embedding determined by the equation

$$W_{\underline{\pi} \circ p} = p^*(\pi_1) \cdot T_1 + \dots + p^*(\pi_n) \cdot T_n = 0.$$

7 Towards a vanishing cycles formalism over $\mathbb{A}_S^1/\mathbb{G}_{m,S}$

The discussion in Sect. 4 suggests that it should be possible to construct a vanishing cycles formalism where the role of the base henselian trait is played by some more general geometric object. Moreover, the monodromy-invariant part of this construction should recover the sheaf of Definition 5.1.1.

Notice that similar ideas, i.e. to develop a formalism of vanishing cycles over a base scheme B with a closed-open decomposition (σ, η) , already appeared in J. Ayoub's works [4–6].

We will give a complete account on the formalism of tame vanishing cycles over $\mathbb{A}_S^1/\mathbb{G}_{m,S}$ in a forthcoming paper in collaboration with D.-C. Cisinski.

7.1 Tame vanishing cycles over \mathbb{A}_S^1

It is well known that even if tame vanishing cycles were first defined for schemes over an henselian trait, what one actually uses is the geometry of \mathbb{A}_S^1 . Indeed, one can consider the following diagram

$$S \xrightarrow{i_0} \mathbb{A}_S^1 = \text{Spec}_S(\mathcal{O}_S[t]) \xleftarrow{j_0} \mathbb{G}_{m,S} \longleftarrow \varprojlim_{n \in \mathbb{G}_{m,S}(S)} \mathbb{G}_{m,S}[x]/(x^n - t) \quad (7.1.1)$$

and use it to replace the one considered in [54, 55] when $S = \text{Spec}(A)$ is a strictly henselian trait

$$\sigma \hookrightarrow S \xleftarrow{\quad \eta \quad} \eta^t. \quad (7.1.2)$$

If we pullback the first diagram along the morphism $S \rightarrow \mathbb{A}_S^1$ given by an uniformizer, we recover the second one. It is not surprising that in this way we are only able to recover the so called tame vanishing cycles. Indeed, the wild inertia group is of arithmetic nature. One can define nearby cycles for schemes over \mathbb{A}_S^1 in the usual way: for $f : X \rightarrow \mathbb{A}_S^1$ consider the diagram

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{i} & X & \xleftarrow{j_n} & X_{\mathcal{U}_n} & \xleftarrow{\phi_n} & X_{\mathcal{U}_\infty} \\
 f_0 \downarrow & & f \downarrow & & \downarrow f_n & & \downarrow f_\infty \\
 S & \xrightarrow{i_0} & \mathbb{A}_S^1 & \xleftarrow{j_{0,n}} & \mathbb{G}_{m,S}[x]/(x^n - t) = \mathcal{U}_n & \xleftarrow{\phi_{0,n}} & \mathcal{U}_\infty = \varprojlim_{n \in \mathcal{O}_S^\times} \mathcal{U}_n.
 \end{array}$$

(7.1.3)

Definition 7.1.1 Let $\mathcal{F} \in \mathbf{Shv}_{\mathbb{Q}_\ell}(X)$.

- The ℓ -adic sheaf of nearby cycles of \mathcal{F} is defined as

$$\Psi_f(\mathcal{F}) := i^* j_{\infty*} j_\infty^* \mathcal{F} \simeq \varinjlim_{n \in \mathcal{O}_S^\times} i^* j_n^* j_n^* \mathcal{F}. \quad (7.1.4)$$

Notice that, since $\mu_{n,S} := \mathcal{O}_S[x]/(x^n - 1)$ naturally acts on $X_{\mathcal{U}_n}$, then $i^* j_n^* j_n^* \mathcal{F}$ has a natural induced action. Then $\Psi_f(\mathcal{F})$ has a natural action of $\varprojlim_{n \in \mathcal{O}_S^\times} \mu_{n,S} =: \mu_{\infty,S}$.

- There is a natural morphism $i^* \mathcal{F} \rightarrow \Psi_f(\mathcal{F})$, that we can see as a $\mu_{\infty,S}$ -equivariant morphism if we endow $i^* \mathcal{F}$ with the trivial action. The sheaf of vanishing cycles $\Phi_f(\mathcal{F})$ of \mathcal{F} is then defined as the cofiber of this morphism, with the induced $\mu_{\infty,S}$ -action.

7.2 Tame vanishing cycles over $\mathbb{A}_S^1/\mathbb{G}_{m,S}$

We can reproduce the situation described above for schemes over $\mathbb{A}_S^1/\mathbb{G}_{m,S}$. In this case, the role of the zero section is played by $B\mathbb{G}_{m,S} \rightarrow \mathbb{A}_S^1/\mathbb{G}_{m,S}$, and that of the open complementary by $S \simeq \mathbb{G}_{m,S}/\mathbb{G}_{m,S} \rightarrow \mathbb{A}_S^1/\mathbb{G}_{m,S}$. We can consider elevation to the n^{th} power in this case too:

$$\Theta_n : \mathbb{A}_S^1/\mathbb{G}_{m,S} \rightarrow \mathbb{A}_S^1/\mathbb{G}_{m,S} \quad (7.2.1)$$

which can be described as the functor

$$(p : X \rightarrow S, \mathcal{L}_X, s_X) \mapsto (p : X \rightarrow S, \mathcal{L}_X^{\otimes n}, s_X^{\otimes n}). \quad (7.2.2)$$

See [13] and [2].

Remark 7.2.1 Notice that the Θ_n are compatible, i.e.

$$\Theta_n \circ \Theta_m \simeq \Theta_{mn} \simeq \Theta_m \circ \Theta_n \quad (7.2.3)$$

for any $n, m \in \mathbb{N}$.

We can therefore consider the diagrams (for each $n \in \mathcal{O}_S^\times$)

$$\begin{array}{ccccccc} & & i_{0,n} & & & & \\ & \swarrow & & \searrow & & & \\ \mathrm{B}\mathbb{G}_{m,S} & \xrightarrow{t_{0,n}} & \mathrm{B}\mathbb{G}_{m,S} \times_{\mathbb{A}_S^1/\mathbb{G}_{m,S}} \mathbb{A}_S^1/\mathbb{G}_{m,S} & \rightarrow & \mathbb{A}_S^1/\mathbb{G}_{m,S} & \xleftarrow{j_{0,n}} & S \\ & \searrow \Theta_{0,n} & \downarrow & & \downarrow \Theta_n & & \downarrow id \\ & & \mathrm{B}\mathbb{G}_{m,S} & \xrightarrow{i_0} & \mathbb{A}_S^1/\mathbb{G}_{m,S} & \xleftarrow{j_0} & S. \end{array} \quad (7.2.4)$$

Notice that even if $t_{0,n} : \mathrm{B}\mathbb{G}_{m,S} \rightarrow \mathrm{B}\mathbb{G}_{m,S} \times_{\mathbb{A}_S^1/\mathbb{G}_{m,S}} \mathbb{A}_S^1/\mathbb{G}_{m,S}$ is not an equivalence, it shows $\mathrm{B}\mathbb{G}_{m,S} \times_{\mathbb{A}_S^1/\mathbb{G}_{m,S}} \mathbb{A}_S^1/\mathbb{G}_{m,S}$ as a nilpotent thickening of $\mathrm{B}\mathbb{G}_{m,S}$. More explicitly, objects of $\mathrm{B}\mathbb{G}_{m,S}$ are pairs $(p : X \rightarrow S, \mathcal{L}_X)$, while those of $\mathrm{B}\mathbb{G}_{m,S} \times_{\mathbb{A}_S^1/\mathbb{G}_{m,S}} \mathbb{A}_S^1/\mathbb{G}_{m,S}$ are triplets $(p : X \rightarrow S, \mathcal{L}_X, s_X)$, where s_X is a n -torsion global section of \mathcal{L}_X . What is important for us is that pullback along $t_{0,n}$ induces an equivalence in étale cohomology. Moreover, Θ_n is a finite morphism of stacks. In particular, the base change theorem should be valid for the cartesian squares above. However, even if the 6-functors formalism for the étale cohomology of stacks has been developed [37–39], the proper base change theorem has been proved in the case of representable morphisms. The morphisms Θ_n are not representable. For example, the pullback of Θ_n along the canonical atlas $\mathbb{A}_n^1 \rightarrow \mathbb{A}_S^1/\mathbb{G}_{m,S}$ is \mathbb{A}_S^1/μ_n .

For any X -point $(p : X \rightarrow S, \mathcal{L}_X, s_X)$ of $\mathbb{A}_S^1/\mathbb{G}_{m,S}$ (e.g. for any twisted LG model over S), we can consider the following diagram, cartesian over (7.2.4)

$$\begin{array}{ccccccc} & & i_n & & & & \\ & \swarrow & & \searrow & & & \\ X_{0,n} & \xrightarrow{t_n} & X_0 \times_X X_n & \longrightarrow & X_n & \xleftarrow{j_n} & X_{\mathcal{U}} \\ & \searrow \Omega_{0,n} & \downarrow & & \downarrow \Omega_n & & \downarrow id \\ & & X_0 & \xrightarrow{i} & X & \xleftarrow{j} & X_{\mathcal{U}}. \end{array} \quad (7.2.5)$$

The same observations we made for the base diagram remain valid in this case too.

Notation 7.2.2 For a stack Y , and a torsion abelian group Λ (we will only be interested in the case $\Lambda = \mathbb{Z}/\ell^n\mathbb{Z}$, $\ell \in \mathcal{O}_S^\times$) let $\mathbf{Shv}(Y, \Lambda)$ the ∞ -category of \mathbf{Mod}_Λ -valued étale sheaves on Y .

We can then propose the following definition.

Definition 7.2.3 Let $\mathcal{F} \in \mathbf{Shv}(X, \Lambda)$. Then

$$\Psi_n(\mathcal{F}) := i_n^* j_{n*} j_n^* \mathcal{F} \simeq i_n^* j_{n*} j_n^* \Omega_n^* \mathcal{F} \in \mathbf{Shv}(X_{0,n}, \Lambda). \quad (7.2.6)$$

Moreover, let

$$sp_n : i_n^* \Omega_n^* \mathcal{F} \rightarrow \Psi_n(\mathcal{F}) \quad (7.2.7)$$

be the morphism induced by the unit of (j_n^*, j_{n*}) .

We can then consider the images of these morphisms in $\mathbf{Shv}(X_0, \Lambda)$

$$\Omega_{0,n*}(sp_n : i_n^* \Omega_n^* \mathcal{F} \rightarrow \Psi_n(\mathcal{F})). \quad (7.2.8)$$

Consider the diagram

$$\begin{array}{ccccccc}
 & & i_{nm} & & & & \\
 & \swarrow & & \searrow & & & \\
 X_{0,nm} & \xrightarrow{\quad} & X_0 \times_X X_{nm} & \xrightarrow{\quad} & X_{nm} & \xleftarrow{j_{nm}} & X_{\mathcal{U}} \\
 \downarrow \Omega_{0,m}^n & & \downarrow i_n & & \downarrow \Omega_m^n & & \downarrow id \\
 X_{0,n} & \xrightarrow{\quad} & X_0 \times_X X_n & \xrightarrow{\quad} & X_n & \xleftarrow{j_n} & X_{\mathcal{U}} \\
 \downarrow \Omega_{0,n}^n & & \downarrow i_0 & & \downarrow \Omega_n & & \downarrow id \\
 X_0 & \xrightarrow{\quad} & X & \xleftarrow{j_0} & X_{\mathcal{U}} & & \\
 \Omega_{0,nm} \nearrow & & & & & &
 \end{array} \quad (7.2.9)$$

Then we have

$$\begin{array}{ccc}
 sp_n : i_n^* \Omega_n^* \mathcal{F} & \xrightarrow{\quad} & i_n^* j_{n*} j_n^* \Omega_n^* \mathcal{F} \\
 \downarrow 1 \rightarrow \Omega_{m*}^n \Omega_m^{n*} & & \downarrow \simeq \\
 \Omega_{0,m*}^n (sp_{nm}) : \Omega_{0,m*}^n i_{nm}^* \Omega_{nm}^* \mathcal{F} & \xrightarrow{\quad} & \Omega_{0,m*}^n i_{nm}^* j_{nm*} j_{nm}^* \Omega_{nm}^* \mathcal{F} \\
 \simeq i_n^* \Omega_{m*}^n \Omega_m^{n*} \Omega_n^* \mathcal{F} & & \simeq i_n^* j_{n*} j_n^* \Omega_n^* \mathcal{F}.
 \end{array} \quad (7.2.10)$$

We can then consider

$$\lim_{n \in \mathbb{N}^\times} \Omega_{0,n*} i_n^* \Omega_n^* \mathcal{F} \rightarrow i_0^* j_{0*} j_0^* \mathcal{F} \in \mathbf{Shv}(X_0, \Lambda). \quad (7.2.11)$$

For $\Lambda = \mathbb{Z}/\ell^d \mathbb{Z}$ one can then consider the induced morphism on ℓ -adic sheaves obtained by taking the limit over d and then tensor with \mathbb{Q}_ℓ . It is expected that in this

way one is able to recover monodromy-invariant vanishing cycles. In order to explain why the first Chern class of the line bundle appears in the computation, it might be useful to look at the base diagram. The ℓ -adic cohomology of $B\mathbb{G}_{m,S}$ is $\mathbb{Q}_\ell[c_1]$, where c_1 is the universal first Chern class and lies in degree 2. We expect that

$$\mathbb{Q}_\ell \otimes_{\mathbb{Z}_\ell} \left(\varprojlim_d \varinjlim_{n \in \mathbb{N}^\times} \Theta_{0,n*} i_n^* \Theta_n^* \mathbb{Z}/\ell^d \mathbb{Z}(\beta) \right) \simeq \text{cofib}(\mathbb{Q}_\ell(\beta) \xrightarrow{c_1} \mathbb{Q}_\ell(\beta)) \quad (7.2.12)$$

and one recovers $\text{cofib}(\mathbb{Q}_{\ell,X_0}(\beta) \xrightarrow{c_1(\mathcal{L}_{X_0})} \mathbb{Q}_{\ell,X_0}(\beta))$ by a formula of the kind

$$\begin{aligned} \mathbb{Q}_\ell[c_1] &\simeq H^*(B\mathbb{G}_{m,S}, \mathbb{Q}_\ell) \rightarrow H^*(X_0, \mathbb{Q}_\ell) \\ c_1 &\mapsto c_1(\mathcal{L}_{X_0}), \end{aligned} \quad (7.2.13)$$

where \mathcal{L}_{X_0} is the line bundle which determines the morphism $X_0 \rightarrow B\mathbb{G}_{m,S}$.

8 Some remarks on the regularity hypothesis

In the theorems about the ℓ -adic realization of the dg category of a (twisted, n -dimensional) LG model the regularity assumption on the ambient scheme is crucial. Indeed, we are not able to compute the motivic realization (and hence, the ℓ -adic one) of $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$. However, this dg category sits in the following pullback diagram

$$\begin{array}{ccc} \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} & \xrightarrow{i_*} & \mathbf{Perf}(X)_{X_0} \\ \downarrow & & \downarrow \\ \mathbf{Coh}^b(X_0) & \xrightarrow{i_*} & \mathbf{Coh}^b(X)_{X_0}. \end{array} \quad (8.0.1)$$

It is well known after Quillen's dévissage for G-theory that

$$\mathcal{M}^\vee(\mathbf{Coh}^b(X_0)) \simeq \mathcal{M}^\vee(\mathbf{Coh}^b(X)_{X_0}). \quad (8.0.2)$$

Therefore, to say that $\mathcal{M}^\vee(i_* : \mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)} \rightarrow \mathbf{Perf}(X)_{X_0})$ is an equivalence or to say that the image of square (8.0.1) via \mathcal{M}^\vee is still a pullback square are equivalent statements. We believe that this is the case, even though this matter will be investigated elsewhere. Notice that to know that such an equivalence holds true would allow to compute the motivic realization of $\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}$ (under the additional hypothesis $X_0 \simeq \pi_0(X_0)$). Indeed, the localization sequence for K-theory would allow us to

compute it as the fiber

$$\begin{aligned}
 \mathcal{M}_X^\vee(\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}) &\simeq \mathcal{M}_X^\vee(\mathbf{Perf}(X)_{X_0}) \\
 &\downarrow \\
 \mathcal{M}_X^\vee(\mathbf{Perf}(X)) & \\
 &\downarrow \\
 \mathcal{M}_X^\vee(\mathbf{Perf}(X - X_0)). &
 \end{aligned} \tag{8.0.3}$$

In other words,

$$\mathcal{M}_X^\vee(\mathbf{Coh}^b(X_0)_{\mathbf{Perf}(X)}) \simeq i_* i^! \mathbf{BU}_X. \tag{8.0.4}$$

In this case, the proof of Theorem 5.2.2 would work without any changes.

Acknowledgements This paper is part of the author's PhD thesis written under the supervision of B. Toën and G. Vezzosi, and to them owes a lot. Their generosity in sharing their ideas and insights is the backbone upon which this work relies on.

The author wishes to thank F. Déglise, T. Dyckerhoff, B. Keller, S. Scherotzke and J. Tapia for their comments and remarks on the work presented in this article.

Many thanks to D. Beraldo, T. Moulinos and M. Robalo for many useful conversations.

The author thanks an anonymous referee for many useful remarks and comments.

This project has received funding from the European Research Council (ERC) under the European Union Horizon 2020 research and innovation programme (Grant agreement NEDAG ADG-741501).

This project has received funding from the European Research Council (ERC) under the European Union Horizon 2020 research and innovation programme (Grant agreement No.725010).

More precisely, this work was carried out while the author was supported by the NEDAG PhD grant ERC-2016-ADG-741501 as a “doctorant contractuel employé par le CNRS à l'unité UMR5219 IMT”, and partially written while he was supported by the BG-BB-AS ERC-2016-COG-725010 as a Research Fellow in the UCL Department of Mathematics in the service of University College London 'UCL'.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Auslander, M., Buchsbaum, D.A.: Homological dimension in Noetherian rings. *Proc. Nat. Acad. Sci. U.S.A.* **42**, 36–38 (1956)
2. Abramovich, D., Graber, T., Vistoli, A.: Gromov–Witten theory of Deligne–Mumford stacks. *Am. J. Math.* **130**(5), 1337–1398 (2008)
3. André, M.: *Homologie des Algèbres Commutatives*. Grundlehren der Mathematischen Wissenschaften. Springer, Berlin (1974)
4. Ayoub, J.: Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique, I. *Astérisque*, Vol. 314, Société Math. France (2007)

5. Ayoub, J.: Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique, II. *Astérisque*, Vol. 315, Société Math. France (2007)
6. Ayoub, J.: La réalisation étale et les opérations de Grothendieck. *Ann. Sci. éc. Norm. Supér. (4)* **47**(1), 1–145 (2014)
7. Barwick, C.: On exact ∞ -categories and the theorem of the heart. In: *Compositio Mathematica*. 151, 11, p. 2160–2186 (2015)
8. Bloch, S.: Cycles on arithmetic schemes and Euler characteristics of curves. *Algebraic geometry (Bowdoin, 1985)*, *Proc. Symp. Pure Math.*, vol. 46, part II, American Mathematical Society, Providence, pp. 421–450 (1985)
9. Blanc, A., Robalo, M., Toën, B., Vezzosi, G.: Motivic realizations of singularity categories and vanishing cycles. *J. de l'École Polytechnique* **5**, 651–747 (2018)
10. Bondal, A., Van den Bergh, M.: Generators and representability of functors in commutative and non-commutative geometry. *Mosc. Math. J.* **3**, 1–36 (2003)
11. Burke, J., Walker, M.: Matrix factorizations in higher codimension. *Trans. Am. Math. Soc.* **367**, 3323–3370 (2015)
12. Ben-Zvi, D., Francis, J., Nadler, D.: Integral transforms and Drinfeld centers in derived algebraic geometry. *J. Am. Math. Soc.* **23**(4), 909–966 (2010)
13. Cadman, C.: Using stacks to impose tangency conditions on curves. *Am. J. Math.* **129**(2), 405–427 (2007)
14. Cisinski, D.-C., Déglise, F.: étale motives. *Compos. Math.* **152**, 556–666 (2016)
15. Cisinski, D.-C., Déglise, F.: *Triangulated Categories of Mixed Motives*. Springer Monographs in Mathematics, Springer, Berlin (2019)
16. Cisinski, D.-C.: Descente par éclatements en K-théorie invariante par homotopie. *Ann. Math.* **177**, 425–448 (2013)
17. Cisinski, D.-C., Tabuada, G.: Non-connective K-theory via universal invariants. *Compos. Math.* **147**(4), 1281–1320 (2011)
18. Cisinski, D.-C., Tabuada, G.: Symmetric monoidal structure on noncommutative motives. *J. K-Theory* **9**(2), 201–268 (2012)
19. Cohn, L.: Differential graded categories are k-linear stable infinity categories. [arXiv:1308.2587](https://arxiv.org/abs/1308.2587) (2013)
20. Déglise, F.: Around the Gysin triangle II. *Doc. Math.* **13**, 613–675 (2008)
21. Deligne, P.: *Catégories Tannakiennes*. Grothendieck Festschrift Vol.II, Progress in Math.87, Birkhäuser, Boston (1990)
22. Drinfeld, V.: Dg quotients of dg categories. *J. Algebra* **272**, 643–691 (2004)
23. Dyckerhoff, T.: Compact generators in categories of matrix factorizations. *Duke Math. J.* **159**(2), 223–274 (2011)
24. Efimov, A.I.: Cyclic homology of categories of matrix factorizations. *Internat. Math. Res. Not.* (12) 3834–3869 (2018)
25. Fujiwara, K.: A proof of the absolute purity conjecture (after Gabber) *Advanced Studies in Pure Mathematics* 36, 2002 Algebraic Geometry 2000, Azumino pp. 153–183
26. Gaiitsgory, D.: Sheaves of categories and the notion of l-affineness. *Contemp. Math.* **643**, 1–99 (2015)
27. Gaiitsgory, D., Lurie, J.: *Weil's Conjectures for Function Fields*. Annals of Mathematics Studies, vol. 199. Princeton University Press, Princeton (2019)
28. Gaiitsgory, D., Rozenblyum, N.: *A study in derived algebraic geometry*. Vol. 1. Correspondences and duality, volume 221 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI (2017)
29. Illusie, L., Laszlo, Y., Orgogozo, F.: *Travaux de Gabber sur l-uniformisation locale et la cohomologie étale des schémas quasi-excellents*, Société Mathématique de France, Paris, *Astérisque* vol. 363–364 (2014)
30. Illusie, L.: *Complexe Cotangent et Déformations I*. Lecture Notes in Mathematics vol. 239, Springer, Berlin (2009) [1971]
31. Khan, A., Rydh, D.: Virtual Cartier divisors and blow-ups. [arXiv:1802.05702](https://arxiv.org/abs/1802.05702) (2019)
32. Keller, B.: On differential graded categories. In *International Congress of Mathematicians*. Vol. II, pp. 151–190. Eur. Math. Soc., Zurich (2006)
33. Laumon, G.: Vanishing cycles over a base of dimension ≥ 1 . *Algebraic geometry (Tokyo/Kyoto, 1982)*, volume 1016 of *Lecture Notes in Math*, pp. 143–150. Springer, Berlin (1983)
34. Lurie, J.: *Higher Algebra*, available for download on the author's webpage (2017)

35. Lurie, J.: Higher Topos Theory. Annals of Mathematics Studies, vol. 170. Princeton University Press, Princeton (2009)
36. Lurie, J.: Spectral Algebraic Geometry, available for download on the author's webpage (2018)
37. Liu, Y., Zheng, W.: Gluing restricted nerves of ∞ -categories. [arXiv:1211.5294](#) (2015)
38. Liu, Y., Zheng, W.: Enhanced six operations and base change theorem for higher Artin stacks. [arXiv:1211.5948](#) (2017)
39. Liu, Y., Zheng, W.: Enhanced adic formalism and perverse t-structures for higher Artin stacks. [arXiv:1404.1128](#) (2017)
40. Morel, F., Voevodsky, V.: \mathbb{A}^1 -homotopy theory of schemes. Publ. Math. Inst. Hautes études Sci. (1999), no. 90, pp. 45–143 (2001)
41. Orgogozo, F.: Modifications et cycles proches sur une base générale. International Mathematical Research Notices (2006)
42. Orlov, D.: Triangulated categories of singularities and equivalences between Landau–Ginzburg models. Mat. Sb. **197**(12), 117–132 (2006)
43. Pippi, M.: On the structure of dg-categories of relative singularities. [arXiv:1911.01332](#) (2019)
44. Pippi, M.: Catégories des singularités, factorisations matricielles et cycles évanescents. Ph.D. Thesis, Université Toulouse III, Paul Sabatier (2020). [arXiv:2006.06301](#)
45. Preygel, A.: Thom-Sebastiani and duality for Matrix factorizations. [arXiv:1101.5834](#) (2011)
46. Quillen, D.: On the (co-)homology of commutative rings. Proc. Symp. Pure Mat. XVII, American Mathematical Society (1970)
47. Riou, J.: Algebraic K-theory, \mathbb{A}^1 -homotopy and Riemann-Roch theorems. J. Topol. **3**(2), 229–264 (2010)
48. Robalo, M.: Motivic homotopy theory of non-commutative spaces. PhD Thesis, Université Montpellier 2, available for download on the author's webpage (2014)
49. Robalo, M.: K-theory and the bridge from motives to noncommutative motives. Adv. Math. **269**, 399–550 (2015)
50. Serre, J.-P.: Sur la dimension homologique des anneaux et des modules noethériens. Proceedings of the international symposium on algebraic number theory, Tokyo and Nikko, 1955, pp. 175–189. Science Council of Japan, Tokyo (1956)
51. Serre, J.-P.: Corps locaux. Publications de l'Institut de Mathématique de l'Université de Nancago, VIII, Actualités Sci. Indust., vol. 1296, Hermann, Paris (1962)
52. Serre, J.-P.: Algèbre locale. Multiplicités. Lecture Notes in Mathematics, 11 Springer Berlin, vii+188 pp (1965)
53. Séminaire de Géométrie Algébrique du Bois Marie - 1966-67 - Théorie des intersections et théorème de Riemann-Roch - (SGA 6), *Lecture notes in mathematics* 225, Berlin; New York: Springer-Verlag, xii+700, 1971, dirigé par P. Berthelot, A. Grothendieck et L. Illusie. Avec la collaboration de D. Ferrand, J.-P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J.-P. Serre
54. Groupes de monodromie en géométrie algébrique. I, Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 I). - Lect. Notes in Math., vol. 288, Springer-Verlag, Berlin, 1972, dirigé par A. Grothendieck. Avec la collaboration de M. Raynaud et D. S. Rim
55. Groupes de monodromie en géométrie algébrique. I, Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 II). - Lect. Notes in Math., vol. 340, Springer, Berlin, 1973, dirigé par P. Deligne et N. Katz
56. Tabuada, G.: Une structure de catégorie de modèles de Quillen sur la catégorie des dg-catégories. Comptes Rendus Mathématique **340**(1), 15–19 (2005)
57. Tabuada, G.: Higher K-theory via universal invariants. Duke Math. J. **145**(1), 121–206 (2008)
58. Thomason, R.W., Trobaugh, T.: Higher algebraic K-theory of schemes and of derived categories. The Grothendieck Festschrift, Vol. III. Progress in math., vol. 88, Birkhäuser Boston, Boston, MA, pp. 247–435 (1990)
59. Toën, B.: The homotopy theory of dg-categories and derived Morita theory. Invent. Math. **167**(3), 615–667 (2007)
60. Toën, B.: Lectures on dg-categories. In: Topics in algebraic and topological K-theory. Lect. Notes in Math., vol. 2008, Springer, Berlin, pp. 243–302 (2011)
61. Toën, B.: Proper local complete intersection morphisms preserve perfect complexes. [arXiv:1210.2827](#) (2012)
62. Toën, B.: Derived Azumaya algebras and generators for twisted derived categories. Invent. Math. **189**(3), 581–652 (2012)

63. Toën, B.: Derived algebraic geometry. *EMS Surv. Math. Sci.* **1**(2), 153–245 (2014)
64. Toën, B., Vaquie, M.: Moduli of objects in dg-categories. *Ann. Sci. école Norm. Sup. (4)* **40**(3), 387–444 (2007)
65. Toën, B., Vezzosi, G.: Homotopical algebraic geometry I: Topos theory. *Adv. Math.* **193**, 257–372 (2005)
66. Toën, B., Vezzosi, G.: Homotopical algebraic geometry II: geometric stacks and applications. *Mem. Am. Math. Soc.* **193**(902), x+224 (2008)
67. Toën, B., Vezzosi, G.: Géométrie non-commutative, formule des traces et conducteur de Bloch. *Actes du 1er Congrès de la SMF-Tours, 2016*, 77–107, Sémin. Congr., 31, SMF, Paris (2017)
68. Toën, B., Vezzosi, G.: The l-adic trace formula for dg-categories and Bloch’s conductor conjecture. *Boll. Unione Mat. Ital.* **12**(1–2), 3–17 (2019)
69. Toën, B., Vezzosi, G.: Trace and Künneth formulas for singularity categories and applications. [arXiv:1710.05902](https://arxiv.org/abs/1710.05902) (2019)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.